

Mechanical Denaturation of DNA: existence of a low temperature denaturation

Enzo Orlandini¹, Somendra M. Bhattacharjee^{2 3}, Davide Marenduzzo⁴, Amos Maritan^{2 4}
and Flavio Seno¹

1 INFN - Dipartimento di Fisica "G. Galilei" , Via Marzolo 8, 35100 Padova, Italy,

2 Abdus Salam International Center for Theoretical Physics, Via Beirut 2, 34014 Trieste Italy,

3 Institute of Physics, Bhubaneswar 751 005 India,

4 International School for Advanced Studies, INFN, Via Beirut 2, 34014 Trieste Italy

Abstract

Recent theoretical predictions on DNA mechanical separation induced by pulling forces are numerically tested within a model in which self-avoidance for DNA strands is fully taken into account. DNA strands are described by interacting pairs of self avoiding walks (SAW) which are pulled apart by a force applied at the two extremities. The whole phase diagram is spanned by extensive Monte Carlo (MC) simulations and the existence of a low temperature denaturation is confirmed. The basic features of the phase diagram and the re-entrant phase boundary are also obtained with a simple heuristic argument based on an energy-entropy estimate.

05.10.Ln, 64.60.Cn, 87.15.La, 87.14.Gy

In the last few years, enormous technical improvements of experimental devices such as atomic force microscopes^{1,2}, optical tweezers^{3,4} and glass microneedles^{5,6} have allowed micromanipulation of single biological systems, and the determination and characterization of new, in part unexpected, behaviours of biomolecules⁷. These developments have been particularly interesting and innovative for DNA molecules. DNA micromanipulation techniques involve handles attached to the two ends of the molecule which serve as anchoring points to manipulable physical supports⁸. In this way it has been possible to study the response of DNA to external torques^{9,10} and its mechanical unzipping in the absence of enzymes^{11,12}. Nevertheless, only very recently theoretical investigations have started to consider the denaturation of DNA under the presence of forces or torques^{13–15}. On the other hand theoretical studies of thermal denaturation (or melting) of DNA have a much longer history^{16,17} (for more recent studies see^{18,19})

Recently, Bhattacharjee²⁰ has extended a minimal model¹⁶ in which the two strands of DNA are ideal polymer chains, introducing a pulling force applied to two extremities^{21,22}. By mapping into a non-hermitian quantum mechanics problem he has shown that the relative polymer can be pulled out (i.e. the DNA can be unzipped) only if the applied force exceeds a critical value.

These results have been further developed^{23,24} by considering analytically solvable models in which DNA strands are represented by directed walks. Surprisingly the force vs temperature phase diagram revealed the presence^{23,24} of a novel re-entrant unzipping transition at low temperature (see Fig. 1). These results have been subsequently confirmed by other authors^{25,26} within ideal chain models of DNA.

The aim of our paper is to study the phase diagram of DNA in the presence of a pulling force with a more realistic model in which self-avoidance for the DNA strands is explicitly taken into account to verify the robustness of the low temperature re-entrance.

We model^{16,19} the two strands of a $2N$ base-pair DNA by two interacting N steps SAW, on a cubic lattice, $\Gamma^1 = \{\Gamma_0^1, \Gamma_1^1, \dots, \Gamma_N^1\}$ and $\Gamma^2 = \{\Gamma_0^2, \Gamma_1^2, \dots, \Gamma_N^2\}$, with Γ_p^i as the lattice site occupied by the p th monomer of the i th chain ($i = 1, 2$). The first monomers Γ_0^1, Γ_0^2

are kept fixed at one lattice unit apart. Each monomer (site) on a strand corresponds to a base and it is supposed to have its complementary at the same contour position in the other strand. Two complementary monomers are favoured to be in contact (i. e., if they occupy two nearest neighbour sites of the lattice) by an energetic gain ϵ representing the binding energy.

In this approach^{16,19–21} the sequence of bases is not explicitly considered since the model is coarse grained in character. In this respect each monomer is not to be thought of as a single base, but as a group of bases (block). Consequently a mismatch between corresponding blocks has to be very disfavoured with respect to a correct matching. This aspect is adequately treated by our model.

To describe the mechanical unzipping of DNA an energetic term $-\vec{f} \cdot \vec{r}$ is added where $\vec{r} = \Gamma_N^1 - \Gamma_N^2$ is the vector joining the two ends of the strands and \vec{f} is an external force taken, without loss of generality, along the z axis, e.g. $\vec{f} = f\hat{e}_z$ (see Fig. 2). Note that at $f = 0$, this model is in the same universality class as that of Ref.¹⁹.

The Hamiltonian of the problem is then given by:

$$H_N = -\epsilon \left(\sum_{i=0}^N \delta(|\Gamma_i^1 - \Gamma_i^2|, 1) \right) - f\hat{e}_z \cdot \vec{r}, \quad (1)$$

where $\delta(a, b)$ is the Kronecker delta and $|x - y|$ is the Euclidean distance between x and y in lattice units.

Throughout all the calculations $\epsilon = +1$ and therefore all the thermodynamic properties of the systems depend on the temperature $T = \beta^{-1}$ and on the force f . The free energy per monomer $\mathcal{F}_N(\beta, f)$, is

$$\mathcal{F}_N(T, f) = -\frac{1}{\beta N} \log \left(\sum \exp(-\beta H_N) \right) \quad (2)$$

where the sum is over all the possible pairs of N -step *SAW*'s Γ^1 and Γ^2 .

The phase diagram can be studied by looking at the average fraction number of contacts¹⁶ Θ and at the z -component of the average end point separation $\langle r^z \rangle$ defined as:

$$\Theta = \frac{\langle N_c \rangle}{N} = -\frac{\partial}{\partial \epsilon} \mathcal{F}_N, \quad \langle r^z \rangle \sim \frac{\partial}{\partial f} (\mathcal{F}_N N) \quad (3)$$

In the denaturated regime Θ should vanish whereas it should be nonzero in the zipped phase. To compute averages, for fixed values of f and N , Monte-Carlo simulations of the model have been performed by a Metropolis based hybrid algorithm^{27,28} that consists of local and global (pivot moves) deformations attempted on each strand of the polymer. In addition a move that tries to interchange the position of pairs of zipped (or unzipped) portions of the double chain has been considered.

The efficiency of the algorithm has been further enhanced by the multiple Markov chain sampling scheme^{29,28}, a method that recently has been shown to be quite effective in exploring the low temperature phase diagrams of interacting polymers. All the runs have been performed by covering, with at most 30 different Markov chains, a large portion of the temperature space ranging from $T = 3$ down to $T = 0.1$. The values of N considered range from $2N = 50$ up to $2N = 150$ and the values of f go from $f = 0$ up to $f = 1$.

In Fig.2 and Fig. 3 the numerical results respectively for Θ and $\langle r^z \rangle / (2N)$ are plotted as a function of T for different values of the force strength f and for systems of $2N = 150$ monomers. The data signal clearly the presence of three distinct regimes.

a) For $f < f_0 \sim 0.5$ the system undergoes a transition from the denaturated to the zipped regime. In particular at $f = 0$ results similar to those found in ref.¹⁹ are re-obtained, namely a first order like transition at $T_D = 0.58 \pm 0.02$, where however the fluctuation in the size of the “bubbles” diverges. Interestingly, our data support the prediction that the critical force near the melting point (T_D) is $f \sim |T_D - T|^a$ with $a < 1$ ^{30,26} (see also below). This is different from the 3-dimensional Gaussian chain case²⁰ or the directed case of Fig. 1²³ (in $d = 1 + 1$) where one finds $a = 1$. In these two cases the zero-force melting transition at T_D is continuous.

b) For $f_0 < f < f_t \sim 0.70$ the unzipped/zipped transition is still present but if the temperature is further lowered the two strands separate again through a low temperature denaturation or unzipping induced by the presence of the force. This is established by the peaks in Fig. 3 and by the minima in Fig. 4 for $f = 0.625$ and $f = 0.65$.

Analysis of the specific heat peaks for different chain lengths support the fact that both

these transitions are first order. One might think of the existence of a subtle difference between the low and the high temperature denaturation. To explore this possibility we studied the size of the molten “bubbles”. For $f = 0$ a simple argument¹⁹ predicts that the size fluctuation of this “bubble” diverges when the thermal denaturation temperature T_D is approached from below. As already mentioned we observed this effect for $f = 0$, but not for $f \neq 0$, both approaching the low temperature transition from above or the high temperature transition from below. We explain this fact by noticing that for $f \neq 0$ the transition temperature is less than T_D and the part of the two strands containing the “bubbles” is not influenced by the presence of the external force (the external force act as a boundary effect). Thus, by approaching the transition line from the zipped phase, bubbles can never become critical if $f \neq 0$. Therefore we conclude that from a statistical point of view low and high temperature denaturation transition are equivalent.

c) For $f > f_t$ the pulling force is always dominating over the base-base interaction and the system is always in the denaturated state.

The value of f_0 can be exactly calculated and the basic features of the phase diagram and the re-entrant phase boundary can be obtained with a simple heuristic argument based on an energy-entropy estimate. Indeed at $T = 0$, when thermal fluctuations are absent, the only allowed configurations are those in which the first l monomers of the strands are parallel and form l contacts, whereas the remaining $N - l$ are pointing in opposite directions parallel to the force. Hence, the resulting energy is a simple function of l :

$$E(l) = -l(1 - 2f) - f(2N + 1) \quad (4)$$

For $f > 1/2$ the energy minimum occurs for $l = 0$ (complete stretching) whereas for $f < 1/2$ it occurs for $l = N$ (complete zipping). This argument states definitively that $f_0 = 1/2$ in full agreement with our numerical findings. Notice that for $f = f_0 = 1/2$, $E(l)$ is independent of l , indicating coexistence.

For low enough temperatures we expect that the dominant configurations are still of Y-like configurations with $2(N - l)$ monomers zipped and two “free” strands each of length

l whose extrema are kept at a relative position \vec{r} by the force \vec{f} . In this Y model the energy is thus given by:

$$E = -(N - l)\epsilon - \vec{r} \cdot \vec{f} \quad (5)$$

whereas the entropy is

$$S = \log \left[P_{2l}(\vec{r}) \mu_u^{2l} \mu_z^{N-l} \right] \quad (6)$$

where $P_N(\vec{r})$ is the probability that an N -step strand has an end-to-end displacement \vec{r} and μ_u and μ_z are the effective coordination of the single strand and of the double strand respectively. For the simple model introduced before $\mu_z = \mu_u$ but in general this need not be true.

It is well known³¹ that

$$P_{2l}(\vec{r}) \sim \exp \left[- \left(\frac{r}{l^\nu} \right)^\delta c \right] \quad l^\nu \ll r \ll l \quad (7)$$

where $r = |\vec{r}|$, $\delta = (1 - \nu)^{-1}$, ν is the usual polymer critical exponent³¹ and c is a constant. For an ideal polymer $\nu = 1/2$ and one recovers the usual Gaussian distribution.

In the last equation power law corrections lead to subleading terms in the entropy estimate. From the last three equations the following free energy is obtained:

$$F(l, \vec{r}) = l(\epsilon - T \log \frac{\mu_u^2}{\mu_z}) - \vec{r} \cdot \vec{f} + cT \left(\frac{r}{l^\nu} \right)^\delta + \text{const} \quad (8)$$

Minimization of F with respect to \vec{r} yields

$$r(l, f) = l \left(\frac{f}{c\delta T} \right)^{\frac{1}{\delta-1}}, \quad (9)$$

where r is the magnitude of \vec{r} in the direction of \vec{f} . This r gives a free energy

$$F(l) \equiv F(l, r(l, f)) = l \left[\epsilon - T \log \frac{\mu_u^2}{\mu_z} - \frac{\delta - 1}{\delta^{\frac{\delta}{\delta-1}}} (cT)^{-\frac{1}{\delta-1}} f^{\frac{1}{\nu}} \right] \quad (10)$$

implying that $F(l)$ gets its minimum at $l = 0$ (zipped phase) and $l = N$ (unzipped phase) when $f < f_c(T)$ and $f > f_c(T)$ respectively with

$$f_c(T) \sim T^{1-\nu} \left(1 - \frac{T}{T_D}\right)^\nu, \quad (11)$$

$T_D = \epsilon / \log\left(\frac{\mu_u^2}{\mu_z}\right)$ being the thermal denaturation temperature for this Y model. Thus at $f_c(T)$ a first order phase transition occurs where the fraction of the bounded bases has a jump. The last equation predicts a re-entrance with $f_c \rightarrow 0$ when $T \rightarrow 0$. This limit is given by a spurious effect due to the assumption that Eq. 7 applies also at very low T where eq. 9 predicts $r \rightarrow \infty$. Indeed if Eq. 7 is assumed to hold till $P_{2l}(r)\mu_u^{2l} \sim 1$, e.g. $r \leq r_M \sim 2l$, and after that r remains constant, then we easily find that at low T

$$f_c(T) = \frac{1}{2}(\epsilon + T \log \mu_z). \quad (12)$$

Eq. (11) near the denaturation temperature and eq. (12) at the lowest T predict a behaviour qualitatively in agreement with the one described in fig. 1³² and confirmed by our simulation analysis.

In conclusion we have shown how a pulling force applied to the extrema of a model of double stranded DNA molecule could introduce non trivial effects in the phase diagram. The results obtained on MC simulations in which, unlike previous studies, excluded volume effects are fully taken into account, confirm the existence of a low temperature denaturation. Although in our approach the intrinsic helicity of the DNA and the randomness of the bases sequences have not been incorporated, we believe, on the basis of preliminary numerical results, that they should not affect the main features of the phase diagram and in particular the existence of a re-entrance region. Therefore we expect that such an effect could be pinpointed by high precision measurements of DNA denaturation under a pulling force.

Acknowledgments: We thank A. Trovato for stimulating discussions. FS acknowledges hospitality by SISSA and ICTP. This work was supported by MURST (COFIN 99).

REFERENCES

- ¹ H.G. Hansma, J. Vac. Sci. Technol. **B14**, 1390 (1996).
- ² C. Bustamante & D. Keller, Phys. Today **48**(12), 32 (1995).
- ³ K. Svoboda & S.M. Block, Annu. Rev. Biophysics. Biomol. Struct. **23**, 247 (1994).
- ⁴ A. Ashkin, Proc. Natl. Acad. Sci. USA **94**, 4853 (1997).
- ⁵ A. Kishino & T. Yanagida, Nature **34**, 74 (1988).
- ⁶ P. Cluzel, A. Lebrun, C. Heller, R. Lavery, J.L. Viovy, D. Chatenay & F. Caron, Science **271**, 792 (1996).
- ⁷ V. Moy , E.L. Florin & H.E. Gaub , Science **264**, 257 (1994); R. Merkel, A. Leung, K. Ritchie & E. Evan, Nature **397**, 50 (1999); M. Rief, F. Oesterhelt, B. Heymann & H.E. Gaub, Science **276**,1109 (1997).
- ⁸ T.R. Strick, J.F. Allemand, V. Croquette & D. Bensimon, Journal of Statistical Physics **93**, 647 (1998).
- ⁹ T.R. Strick, J.F. Allemand, V. Croquette & D. Bensimon, Physica **263A**, 392 (1999).
- ¹⁰ J.F. Leger, G. Romano, A. Sarkar, J. Robert, I. bordieu, D Chatenay & J.F. Marko, Phys. Rev. Lett. **83**, 1066 (1999).
- ¹¹ B. Essevaz-Roulet, U. Bockelmann & F. Heslot, Proc. Natl. Acad. Sci. USA **94**, 11935 (1997).
- ¹² U. Bockelmann, B. Essevaz-Roulet & F. Heslot, Phys. Rev. Lett. **79**, 4489 (1997); Phys. Rev. **E58**, 2386 (1998).
- ¹³ S. Cocco & R. Monasson, Phys. Rev. Lett. **83** 5178 (1999).
- ¹⁴ M. Peyrard, Europhys. Lett. **44**, 271 (1998).
- ¹⁵ R.E. Thompson & E. Siggia, Europhys. Lett. **31**, 335 (1995).

- ¹⁶ D. Poland & H.A. Scheraga. J. of Chem. Phys. **45**, 1464-1469 (1966).
- ¹⁷ M.E. Fisher, J. Chem. Phys. **45**, 1469 (1966).
- ¹⁸ D. Cule & T. Hwa, Phys. Rev. Lett. **79**, 2375 (1997).
- ¹⁹ M.S. Causo, B. Coluzzi & P. Grassberger, Phys. Rev. **E62**, 3958 (2000).
- ²⁰ S.M. Bhattacharjee, cond-mat/9912297, J. Phys. **A 33** L423 (2000); J. Phys. **A 33**, 9003 (2000); cond-mat/0010132.
- ²¹ More recently a similar model which takes into account also the presence of sequence disorder has been studied by D. K. Lubensky & D. R. Nelson, Phys. Rev. Lett. **85**, 1572 (2000). See also S. M. Bhattacharjee and D. Marenduzzo, cond-mat/0106110.
- ²² The dynamics of unzipping has also been studied recently: see²³, K. L. Sebastian, Phys. Rev. E **62**, 1128 (2000); S. Cocco, R. Monasson and J. F. Marko, Proc. Natl. Acad. Sci. USA **98**, 8908 (2001).
- ²³ D. Marenduzzo, S.M. Bhattacharjee, A. Maritan, E. Orlandini and F. Seno, cond-mat/0103142.
- ²⁴ D. Marenduzzo, A. Trovato and A. Maritan, Phys. Rev. E **64**, 031901 (2001).
- ²⁵ D. K. Lubensky, D. R. Nelson, cond-mat/0107423.
- ²⁶ E. A. Mukamel, E. I. Shakhnovich, cond-mat/0108447.
- ²⁷ N. Madras & A. Sokal, J. Stat. Phys. **47**, 573 (1987).
- ²⁸ E. Orlandini, F. Seno & A.L. Stella, Phys. Rev. Lett. **84**, 294 (2000).
- ²⁹ M.C. Tesi, E.J. van Rensburg, E. Orlandini & S.G. Whittington, J. Stat. Phys. **29**, 2451 (1996).
- ³⁰ Y. Kafri, D. Mukamel, L. Peliti, cond-mat/0108323.

³¹ M. E. Fisher, J. Chem. Phys. **44**, 616 (1966), see also P.G. de Gennes, *Scaling concepts in polymer physics* (Cornell Univ. Press, Ithaca, 1979) sec. 1.2.

³² In the critical case of fig. 1 ν is the critical correlation length exponent for melting as in Ref.²⁰

FIGURES




FIG. 1. The exact force (f) versus temperature (T) phase diagram as obtained in ref.²³ for a directed model ($d=2$) of DNA. See eq. (1) for definition of force. The values of f_t , f_0 and T_D are model dependent.




FIG. 2. A typical configuration of a double stranded DNA chains ($2N = 200$) obtained with the model described in the text. The different colours indicate the two different strands.




FIG. 3. Average fraction of bound pairs Θ versus T , for several values of the force f and for $2N = 150$. Error bars are of the size of the symbols. The two peaks ($f = 0.625$ and $f = 0.65$) signal a zipped phase between two unzipped regions (re-entrance).




FIG. 4. End to end end separation in the force direction versus T , for different values of the force f for $2N = 150$. Error bars correspond to one standard deviation. The two minima ($f = 0.625$ and $f = 0.65$) signal a zipped phase between two unzipped regions (re-entrance).