

Fractal Von Neumann Entropy

Wellington da Cruz*

Departamento de Física,

Universidade Estadual de Londrina, Caixa Postal 6001,

Cep 86051-970 Londrina, PR, Brazil

(October 26, 2018)

Abstract

We consider the *fractal von Neumann entropy* associated with the *fractal distribution function* and we obtain for some *universal classes h of fractons* their entropies. We obtain also for each of these classes a *fractal-deformed Heisenberg algebra*. This one takes into account the braid group structure of these objects which live in two-dimensional multiply connected space.

PACS numbers: 05.30.-d; 05.70.Ce; 11.25.Hf

Keywords: Fractal von Neumann entropy; Fractal distribution function; Fractons; Fractal-deformed Heisenberg algebras

Typeset using REVTeX

*E-mail: wdacruz@exatas.uel.br

The *fractal von Neumann entropy* associated with the *fractal distribution function* was introduced in Ref. [1]. We have defined *universal classes h of particles or quasiparticles* which carry rational or irrational spin quantum number s . The fractal parameter or Hausdorff dimension h runs within the interval $1 < h < 2$. *Fractons* as charge-flux systems are such particles which live in two-dimensional multiply connected space and satisfy specific fractal distribution function. The particles of each class are collected taking into account the *fractal spectrum*

$$h - 1 = 1 - \nu, \quad 0 < \nu < 1; \quad h - 1 = \nu - 1, \quad 1 < \nu < 2; \quad (1)$$

etc.

and the spin-statistics relation $\nu = 2s$, which is valid for such fractons. For example, consider the universal classes with distinct values of spin $\left\{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\right\}_{h=1}$, $\left\{0, 1, 2, \dots\right\}_{h=2}$ and $\left\{\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, \dots\right\}_{h=\frac{3}{2}}$, then we have the Fermi-Dirac distribution

$$n[1] = \frac{1}{\xi + 1} \quad (2)$$

the Bose-Einstein distribution

$$n[2] = \frac{1}{\xi - 1} \quad (3)$$

and the Fractal distribution

$$n\left[\frac{3}{2}\right] = \frac{1}{\sqrt{\frac{1}{4} + \xi^2}} \quad (4)$$

with $\xi = \exp\{(\epsilon - \mu)/KT\}$.

Another result of our approach to fractional spin particles is that the classes h satisfy a *duality symmetry* defined by $\tilde{h} = 3 - h$, and as a consequence we extract a *fractal supersymmetry* which defines pairs of particles $(s, s + \frac{1}{2})$. Thus, *the fractal distribution function is a quantum-geometrical description of the statistical laws of Nature, since the quantum path is a fractal curve and this reflects the Heisenberg uncertainty principle.*

The **Fractal von Neumann entropy** per state in terms of the average occupation number is given as [1]

$$\mathcal{S}_G[h] = n K \left[(\mathcal{Y}[\xi] - 1) \ln(\mathcal{Y}[\xi] - 1) - (\mathcal{Y}[\xi] - 2) \ln(\mathcal{Y}[\xi] - 2) \right] \quad (5)$$

$$= K \left[[1 + (h - 1)n] \ln \left\{ \frac{1 + (h - 1)n}{n} \right\} - [1 + (h - 2)n] \ln \left\{ \frac{1 + (h - 2)n}{n} \right\} \right] \quad (6)$$

where

$$n = \frac{1}{\mathcal{Y}[\xi] - h} \quad (7)$$

is the **Fractal distribution function** which appears as a simple and elegant generalization of the fermionic and bosonic distributions for particles with braiding properties. The function $\mathcal{Y}[\xi]$ satisfies the equation

$$\xi = \left\{ \mathcal{Y}[\xi] - 1 \right\}^{h-1} \left\{ \mathcal{Y}[\xi] - 2 \right\}^{2-h}. \quad (8)$$

We also have

$$\xi^{-1} = \left\{ \Theta[\mathcal{Y}] \right\}^{h-2} - \left\{ \Theta[\mathcal{Y}] \right\}^{h-1} \quad (9)$$

where

$$\Theta[\mathcal{Y}] = \frac{\mathcal{Y}[\xi] - 2}{\mathcal{Y}[\xi] - 1} \quad (10)$$

is the *single-particle partition function*. The free energy for the particles in a given quantum state is expressed as

$$\mathcal{F}[h] = KT \ln \Theta[\mathcal{Y}], \quad (11)$$

such that for fermions

$$\mathcal{F}[1] = -KT \ln \left\{ 1 + \xi^{-1} \right\} \quad (12)$$

and for bosons

$$\mathcal{F}[2] = KT \ln \{1 - \xi^{-1}\}. \quad (13)$$

Hence, we find the average occupation number

$$n[h] = \xi \frac{\partial}{\partial \xi} \ln \Theta[\mathcal{Y}] = \frac{1}{KT} \xi \frac{\partial \mathcal{F}}{\partial \xi}. \quad (14)$$

Using the thermodynamic relation $\mathcal{S} = -\left(\frac{\partial \mathcal{F}}{\partial T}\right)_{V,N}$, we obtain another general expression for the entropy

$$\mathcal{S}_G[h] = -K \ln \Theta[\mathcal{Y}] - \frac{KT}{\Theta[\mathcal{Y}]} \frac{\partial \Theta[\mathcal{Y}]}{\partial T} \quad (15)$$

$$= -K \ln \Theta[\mathcal{Y}] - \frac{KT}{(\mathcal{Y} - 2)(\mathcal{Y} - 1)} \frac{\partial \mathcal{Y}}{\partial T}, \quad (16)$$

where the energy is given by

$$\mathcal{E} = -\frac{KT^2}{\Theta[\mathcal{Y}]} \frac{\partial \Theta[\mathcal{Y}]}{\partial T} \quad (17)$$

$$= -\frac{KT^2}{(\mathcal{Y} - 2)(\mathcal{Y} - 1)} \frac{\partial \mathcal{Y}}{\partial T}, \quad (18)$$

since we have $\mathcal{E} = \mathcal{F} + T\mathcal{S}$.

Now, if you consider the density matrix ρ , the entropy is found to be [2]

$$\frac{\mathcal{S}}{K} = -\text{Tr } \rho \ln \rho \quad (19)$$

$$= -\sum_n \mathcal{W}(n) \mathcal{P}(n) \ln \mathcal{P}(n) \quad (20)$$

where

$$\mathcal{W}(n) = \frac{[G + (nG - 1)(h - 1)]!}{[nG]! [G + (nG - 1)(h - 1) - nG]!} \quad (21)$$

is the statistical weight for the classes of fractons and $N = nG$ is the number of particles, G is the number of states, n is the average occupation number [1]. The microstate probability is given by

$$\mathcal{P}(n) = p^{nG} q^{[n(h-2)+1]G}, \quad (22)$$

with $p + q = 1$ and the total probability is unity

$$\sum_n \mathcal{W}(n) \mathcal{P}(n) = 1. \quad (23)$$

Differentiating this last expression with respect to p , we find

$$n \left[\frac{q}{p} + 2 \right] - nh = 1 \quad (24)$$

and defining $\left[\frac{q}{p} + 2 \right] \equiv \mathcal{Y}[\xi]$, we obtain again the fractal distribution function. Considering now the Eq.(20) plus the Eqs.(21) and (22) we have another derivation of the Eqs.(5) and (6) for the fractal von Neumann entropy.

The entropies for fermions $\left\{ \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots \right\}_{h=1}$ and bosons $\left\{ 0, 1, 2, \dots \right\}_{h=2}$, can be recovered promptly

$$\mathcal{S}_G[1] = -K \left\{ n \ln n + (1 - n) \ln(1 - n) \right\} \quad (25)$$

and

$$\mathcal{S}_G[2] = K \left\{ (1 + n) \ln(1 + n) - n \ln n \right\}. \quad (26)$$

For fractons of the self-dual class $\left\{ \frac{1}{4}, \frac{3}{4}, \frac{5}{4}, \dots \right\}_{h=\frac{3}{2}}$, we have

$$\mathcal{S}_G \left[\frac{3}{2} \right] = K \left\{ (2 + n) \ln \sqrt{\frac{2 + n}{2n}} - (2 - n) \ln \sqrt{\frac{2 - n}{2n}} \right\} \quad (27)$$

and for two more examples, the dual classes $\left\{ \frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \dots \right\}_{h=\frac{4}{3}}$ and $\left\{ \frac{1}{6}, \frac{5}{6}, \frac{7}{6}, \dots \right\}_{h=\frac{5}{3}}$, the entropies read as

$$\mathcal{S}_G \left[\frac{4}{3} \right] = K \left\{ (3 + n) \ln \sqrt[3]{\frac{3 + n}{3n}} - (3 - 2n) \ln \sqrt[3]{\frac{3 - 2n}{3n}} \right\} \quad (28)$$

and

$$\mathcal{S}_G \left[\frac{5}{3} \right] = K \left\{ (3 + 2n) \ln \sqrt[3]{\frac{3 + 2n}{3n}} - (3 - n) \ln \sqrt[3]{\frac{3 - n}{3n}} \right\}. \quad (29)$$

We have also introduced the topological concept of *fractal index* associated with each class (h is a geometrical parameter related to the quantum path of the particles) and defined by [3]

$$i_f[h] = \frac{6}{\pi^2} \int_{\infty(T=0)}^{1(T=\infty)} \frac{d\xi}{\xi} \ln \{ \Theta[\mathcal{Y}(\xi)] \} \quad (30)$$

so we obtain for the bosonic class $i_f[2] = 1$, for the fermionic class $i_f[1] = 0.5$ and for some classes of fractons we have $i_f[\frac{3}{2}] = 0.6$, $i_f[\frac{4}{3}] = 0.56$, $i_f[\frac{5}{3}] = 0.656$. Thus for the interval of definition $1 \leq h \leq 2$, there exists the correspondence $0.5 \leq i_f[h] \leq 1$, which signalizes the connection between fractons and quasiparticles of the conformal field theories, in accordance with the unitary $c < 1$ representations of the central charge.

On the other hand, fractons satisfy a *fractal-deformed Heisenberg algebra* which generalizes the fermionic and bosonic ones. The fractal-deformed Heisenberg algebra is obtained of the relation

$$\mathbf{a}(x)\mathbf{a}^\dagger(y) - f[\pm h]\mathbf{a}^\dagger(y)\mathbf{a}(x) = \delta(x - y), \quad (31)$$

between creation and annihilation operators. The factor of deformation is defined as

$$f[\pm h] = \exp(\pm ih\pi), \quad (32)$$

such that for $h = 1$ and $x = y$, we reobtain the fermionic anticommutation relations $\{\mathbf{a}(x), \mathbf{a}^\dagger(x)\} = 1$, and for $h = 2$ and $x = y$, we reobtain the bosonic commutation relations $[\mathbf{a}(x), \mathbf{a}^\dagger(x)] = 1$. If $x \neq y$ and $1 < h < 2$, we have nonlocal operators for fractons

$$\mathbf{a}(x)\mathbf{a}^\dagger(y) = f[\pm h]\mathbf{a}^\dagger(y)\mathbf{a}(x). \quad (33)$$

The braiding relations have the hermiticities

$$\mathbf{a}(x)\mathbf{a}^\dagger(y) = f[+h]\mathbf{a}^\dagger(y)\mathbf{a}(x) \quad (34)$$

$$\mathbf{a}^\dagger(x)\mathbf{a}(y) = f[+h]\mathbf{a}(y)\mathbf{a}^\dagger(x) \quad (35)$$

$$\mathbf{a}(x)\mathbf{a}^\dagger(y) = f[-h]\mathbf{a}^\dagger(y)\mathbf{a}(x) \quad (36)$$

$$\mathbf{a}^\dagger(x)\mathbf{a}(y) = f[-h]\mathbf{a}(y)\mathbf{a}^\dagger(x) \quad (37)$$

$$\mathbf{a}(x)\mathbf{a}(y) = f[+h]\mathbf{a}(y)\mathbf{a}(x) \quad (38)$$

$$\mathbf{a}^\dagger(x)\mathbf{a}^\dagger(y) = f[+h]\mathbf{a}^\dagger(y)\mathbf{a}^\dagger(x) \quad (39)$$

$$\mathbf{a}(x)\mathbf{a}(y) = f[-h]\mathbf{a}(y)\mathbf{a}(x) \quad (40)$$

$$\mathbf{a}^\dagger(x)\mathbf{a}^\dagger(y) = f[-h]\mathbf{a}^\dagger(y)\mathbf{a}^\dagger(x), \quad (41)$$

where the plus sign of h stands for anticlockwise exchange and the minus sign of h for clockwise exchange.

The statistics parameter ν for the plus sign has the pattern

$$\{-, +, -, +, \dots\}_h \quad (42)$$

and the minus sign, gives us another one

$$\{+, -, +, -, \dots\}_h. \quad (43)$$

For example, consider the class $h = \frac{3}{2}$, then

$$\left\{ \mp \frac{1}{2}, \pm \frac{3}{2}, \mp \frac{5}{2}, \pm \frac{7}{2}, \dots \right\}_{h=\frac{3}{2}}, \quad (44)$$

i.e.

$$f[+h] = e^{-i\frac{1}{2}\pi} = e^{+i\frac{3}{2}\pi} = e^{-i\frac{5}{2}\pi} = e^{+i\frac{7}{2}\pi} = \dots \quad (45)$$

and

$$f[-h] = e^{+i\frac{1}{2}\pi} = e^{-i\frac{3}{2}\pi} = e^{+i\frac{5}{2}\pi} = e^{-i\frac{7}{2}\pi} = \dots \quad (46)$$

As we can see, in each class, the particles with different values of statistics parameter ($\nu = 2s$) have unambiguously distinct braiding properties to obey the fractal distribution function determined by the parameter h .

The phase of the wave function (consider two-particle system) changes by $+\pi\nu$ (anti-clockwise exchange) and $-\pi\nu$ (clockwise exchange) in response to which way we braid in interchanging x and y . On the one hand, the violation of the discrete symmetries of parity and time reversal is verified for such fractons, i.e.

$$e^{\pm i\pi\nu} \longrightarrow e^{\mp i\pi\nu}. \quad (47)$$

We have applied these ideas in some systems of the condensed matter such as fractional quantum Hall effect [1] and Luttinger liquids [4]. Besides this we have also considered some connection with conformal field theories [3] . Another direction of research is the connection between **Number Theory and Physics** as explicated by our formulation [1,3]. There exists other possibilities of application, for example, we are considering the study of entanglement states for *fracton quantum computing* [5].

In conclusion, we have considered in this Letter, the fractal von Neumann entropy associated with the fractal distribution function for some universal classes h of fractons. The derivation of the Eqs.(5) and (6) obtained in Ref. [1] using the Eq.(20) suggested in Ref. [2] and adapted to our formulation is another result of this paper. We have also determined the fractal-deformed Heisenberg algebras for these classes of fractons which obey specific fractal distribution function. A next step is to consider the construction of the nonlocal operators for fractons taking into account an angle function as discussed in [6].

REFERENCES

- [1] W. da Cruz, Int. J. Mod. Phys. **A15** (2000), p.3805.
- [2] A. K. Rajagopal and S. Teitler, Physica **A147** (1988), p.627;
A. K. Rajagopal, Phys. Rev. Lett. **74** (1995), p.1048.
- [3] W. da Cruz and R. de Oliveira, Mod. Phys. Lett. **A15** (2000), p.1931.
- [4] W. da Cruz, J. Phys: Cond. Matter. **12** (2000), p.L673.
- [5] W. da Cruz, work in progress.
- [6] A. Lerda and S. Sciuto, Nucl. Phys. **B401** (1993), p.613 ;
M. Frau, A. Lerda and S. Sciuto, hep-th/9407161;
M. Lüscher, Nucl. Phys. **B326** (1989), p.557 ;
V. F. Müller, Z. Phys. **C47** (1990), p.301;
D. Eliezer and G. W. Semenoff, Ann. Phys. **217** (1992), p.66;
and references therein.