

# Theory of the Transient and Stationary Fluctuation Theorems for a Dragged Brownian Particle in a Fluid

R. van Zon and E.G.D. Cohen  
The Rockefeller University  
1230 York Ave., New York NY 10021-6399

22 October 2002

**Comment on “Experimental Demonstration of Violations of the Second Law of Thermodynamics for Small Systems and Short Time Scales” [1].**

In a recent experimental demonstration of the *transient fluctuation theorem* (TFT) by Wang *et al.*[1], a small latex bead, initially at rest in a harmonic trap and in equilibrium with a surrounding fluid, starts to be dragged at a velocity  $\mathbf{v}_{opt}$  at  $t = 0$  by means of the trap. Measurements of the entropy production over a time  $\tau$ ,

$$\Sigma_\tau \equiv \frac{1}{k_B T} \mathbf{v}_{opt} \cdot \int_0^\tau \mathbf{F}(t) dt, \quad (1)$$

were made, with  $T$  the temperature,  $k_B$  Boltzmann’s constant,  $\mathbf{F}(t) = -k(\mathbf{x}(t) - \mathbf{x}_0(t))$ ,  $\mathbf{x}(t)$  the bead’s position, and  $\mathbf{x}_0(t)$  zero for  $t < 0$ , and  $\mathbf{v}_{opt}t$  for  $t > 0$ . Thus,  $P_T(\Sigma_\tau < 0)$  and  $P_T(\Sigma_\tau > 0)$ , the probabilities to find a negative resp. positive values of  $\Sigma_\tau$ , were measured and shown to comply with the integrated TFT[1, 2]:

$$\frac{P_T(\Sigma_\tau < 0)}{P_T(\Sigma_\tau > 0)} = \langle \exp[-\Sigma_\tau] \rangle_{T+}, \quad (2)$$

where on the right-hand side (r.h.s.) the average is performed over transient ( $T$ ) trajectories with positive  $\Sigma_\tau$ .

Assuming the probability distribution  $P(\mathbf{x}, t)$  is described by the Smoluchowski equation, it satisfies

$$\partial_t P(\mathbf{x}, t) = \nabla \cdot \{D \nabla + \tau_r^{-1} [\mathbf{x} - \mathbf{x}_0(t)]\} P(\mathbf{x}, t). \quad (3)$$

with  $D$  is the diffusion constant and  $\tau_r$  the particle’s relaxation time in the harmonic trap. Since Eq. (1) is a linear combination of positions at different




Figure 1: l.h.s. and r.h.s. of Eq. (2) (solid line) and Eq. (6) (dotted line and dashed line, resp.), for  $\sigma\tau_r = 1$ .

times, and the Greens function and equilibrium state are Gaussian, so will the  $\Sigma_\tau$  distribution. It can then be characterized by its first two moments

$$\begin{aligned}\langle \Sigma_\tau \rangle_T &= \sigma\tau - \sigma\tau_r(1 - e^{-\tau/\tau_r}) \\ \langle \Delta\Sigma_\tau^2 \rangle_T &= \langle [\Sigma_\tau - \langle \Sigma_\tau \rangle]^2 \rangle_T = 2\langle \Sigma_\tau \rangle_T.\end{aligned}\quad (4)$$

Here  $\sigma = |\mathbf{v}_{opt}|^2/D$  is the entropy production rate in the non-equilibrium stationary state (NESS). Eq. (2) follows from Eq. (4) and the Gaussian distribution of  $\Sigma_\tau$ . Sevick *et al.* have independently developed an equivalent theory for the TFT from a Langevin approach[3].

Although the TFT looks similar to the *stationary state fluctuation theorem*[4] (SFT), it is quite different[5]. The TFT looks at variations in entropy production over a trajectory of length  $\tau$  starting from an initial equilibrium ensemble. The SFT deals with the distribution of entropy production  $\Sigma_\tau$  over trajectory segments of length  $\tau$  along a single trajectory in the NESS. As above, the SFT can be derived from Eq. (3) because the NESS is Gaussian, as is the stationary (*S*) distribution of  $\Sigma_\tau$ . One obtains

$$\langle \Sigma_\tau \rangle_S = \sigma\tau; \quad \langle \Delta\Sigma_\tau^2 \rangle_S = \langle \Delta\Sigma_\tau^2 \rangle_T \quad (5)$$

Now  $\langle \Delta\Sigma_\tau^2 \rangle_S \rightarrow 2\langle \Sigma_\tau \rangle_S$  only as  $\tau \rightarrow \infty$  [Eqs. (4,5)], so

$$\frac{P_S(\Sigma_\tau < 0)}{P_S(\Sigma_\tau > 0)} \stackrel{\tau \rightarrow \infty}{\cong} \langle \exp[-\Sigma_\tau] \rangle_{S+}, \quad (6)$$

Fig. 1 shows the left-hand side (l.h.s.) and r.h.s. of Eqs. (2), and (6); for  $\tau \rightarrow \infty$ , the curves of the SFT indeed approach each other [Eq. (6)], but the curves of the TFT and of the SFT approach a constant, non-trivial ratio

$$\frac{P_T(\Sigma_\tau < 0)}{P_T(\Sigma_\tau > 0)} \bigg/ \frac{P_S(\Sigma_\tau < 0)}{P_S(\Sigma_\tau > 0)} \stackrel{\tau \rightarrow \infty}{=} \exp \left[ \frac{\sigma\tau_r}{2} \right],$$

as follows from the asymptotics of the error functions in the explicit expressions for the left-hand sides of Eqs. (2) and (6). So a negative  $\Sigma_\tau$  is less likely in the NESS than in the transient state. In fact for large  $\sigma\tau_r$ , negative fluctuations in the NESS are exponentially suppressed. This is important if one wants to devise an experiment on the SFT. More details will be given in a future publication.

This work has been supported by the U.S. Department of Energy, under grant No. DE-FG-02-88-ER13847.

## References

- [1] G.M. Wang *et al.*, Phys. Rev. Lett. **89**, 050601 (2002).
- [2] D.J. Evans and D.J. Searles, Phys. Rev. E **50**, 1645 (1994).
- [3] E.M. Sevick, D.R.M. Williams, D.J. Searles and D.J. Evans, private communication.
- [4] D.J. Evans, E.G.D. Cohen and G.P. Morriss, Phys. Rev. Lett. **71**, 2401 (1993); G. Gallavotti and E.G.D. Cohen, J. Stat. Phys. **80**, 931 (1995).
- [5] E.G.D. Cohen and G. Gallavotti, J. Stat. Phys. **96**, 1343 (1999).