

The Anderson transition in quantum chaos

Antonio M. García-García

Physics Department, Princeton University, Princeton, New Jersey 08544, USA

Jiao Wang

Temasek Laboratories, National University of Singapore, 119260 Singapore

We investigate the effect of classical singularities in the quantum properties of non-random Hamiltonians. We present explicit results for the case of a kicked rotator with a non-analytical potential though extensions to higher dimensionality or conservative systems are straightforward. It is shown that classical singularities produce anomalous diffusion in the classical phase space. Quantum mechanically, the eigenstates of the evolution operator are power-law localized with an exponent universally given by the type of classical singularity. For logarithmic singularities, the classical motion presents $1/f$ noise and the quantum properties resemble those of an Anderson transition, namely, multifractal eigenstates and critical statistics. Neither the classical nor the quantum properties depend on the details of the potential but only on the type of singularity. It is thus possible to define new universality class in quantum chaos by the relation between classical singularities (anomalous diffusion) and quantum power-law localization.

PACS numbers: 72.15.Rn, 71.30.+h, 05.45.Df, 05.40.-a

The description of quantum features based on the underlying classical dynamics has been intensively investigated since the early days of quantum mechanics. For instance, the celebrated Bohigas-Giannoni-Schmit conjecture [1] states that the spectral correlations of a quantum system whose classical counterpart is fully chaotic depend only on the global symmetries of the system and are identical to those of a random matrix with the same symmetry (usually referred to as Wigner-Dyson statistics (WD)). However this conjecture is not always verified. A paradigmatic example is the kicked rotor: Although classically chaotic, its spectral correlations follow Poisson statistics and eigenfunctions are exponentially localized. This behavior was fully understood [2] after mapping the kicked rotator problem onto an 1D disordered system (free particle in a random potential). It is by now well established that, in less than three dimensions, all the eigenstates of a disordered system are exponentially localized in the thermodynamic limit for any amount of disorder. However, in three and higher dimensions, there exists a metal insulator transition (MIT), also called Anderson transition, for a critical amount of disorder. At the MIT, the wavefunction moments \mathcal{P}_q present anomalous scaling with respect to the sample size L , $\mathcal{P}_q = \int d^d r |\psi(\mathbf{r})|^{2q} \propto L^{-D_q(q-1)}$, where D_q is a set of exponents describing the transition. Wavefunctions with such a nontrivial scaling are said to be multifractal [3]. Spectral fluctuations at the MIT are commonly referred to as ‘critical statistics’ [4]. Typical features include: scale invariant spectrum [5], level repulsion and linear number variance [6]. Similar properties have also been observed in disordered systems with random power-law hopping [7] (linked to classical anomalous diffusion [8]) provided that the exponent of the hopping decay matches the dimension of the space. In the 1D case

($1/r$ decay) [9] the relation to a MIT has been analytically established. For *non-random* power-law hopping a MIT was recently reported [10] even for power-law exponents larger than the dimensionality of the space. Unlike the random case, critical states appear only in a certain energy window. A similar power-law behavior has also been found in certain non-random Hamiltonians: Fermi accelerator [11], Coulomb billiard [12], Anisotropic Kepler problem [13] and generalized Kicked rotors [14, 15]. In all of them the potential has a singularity and consequently the KAM theorem does not hold. The classical dynamics of these systems is intermediate between chaotic and integrable and the classical phase space is homogeneously filled with cantori. The goal of this letter is to show that, in certain cases, classical chaotic Hamiltonians with non-analytical potentials present classical anomalous diffusion and quantum power-law localization of the eigenstates. The case of logarithmic singularities, corresponding to a MIT, will be investigated in detail.

Classical mechanics—We study the Hamiltonian

$$\mathcal{H} = \frac{p^2}{2} + V(q) \sum_n \delta(t - nT) \quad (1)$$

with $V(q) = \epsilon|q|^\alpha$ and $V(q) = \epsilon \log(|q|)$; $q \in [-\pi, \pi]$, $\alpha \in [-1, 1]$ and ϵ a real number. The dynamical evolution over T is dictated by the map: $p_{n+1} = p_n - \frac{\partial V(q_n)}{\partial q_n}$, $q_{n+1} = q_n + T p_{n+1} \pmod{2\pi}$. Our aim is to show that, though obviously deterministic, the motion is accurately modeled by an anomalous diffusion process. In particular we compute the classical density of probability $P(p, t)$ for finding the particle at time t with a momentum p . Unlike the case of normal diffusion, the information obtained from the knowledge of a few moments of the distribution is not sufficient to fully characterize the classical motion. For instance, the second moment may be $\langle r^2 \rangle \sim t$ but




FIG. 1: $P(p, t)$ versus p (log scale). In all cases $t = 2000$ except for $\alpha = -0.5$ where $t = 40$. For $V(q) = |q|^\alpha$ with $\alpha = 0.9$ (solid) and $V(q) = (|q| + 0.005)^{0.4}$ (dash) the diffusion is normal (squares and circles are for best Gaussian fitting respectively). However for $\alpha = 0.4$ (dot) and $\alpha = -0.5$ (dash-dot), $P(p, t) \propto tp^{\gamma(\alpha)}$. Inset shows γ as a function of α . The star corresponds to $V(q) = \log(|q|)$ (see Fig.2).




FIG. 2: $P(p, t = 40)$ versus p (semi-log scale). For $V(q) = \log(|q|)$ (solid) $P(p, t)$ is very well fitted by the Lorentzian distribution $P(p, t) = \frac{1}{\pi} \frac{\nu(t)}{1 + p^2 \nu^2(t)}$ with $\nu(t) = 2/t$ (squares). As shown in the inset, $P(p, t)$ (solid) is not affected by adding a noise term $\cos(q)$ to $V(q)$ (circles).

this by no means assures that the density of probability is Gaussian like [16].

The classical density of probability $P(p, t)$ has been evaluated by evolving 10^8 initial conditions uniformly distributed in $(q, p) \in [-\pi, \pi) \times [-\pi, \pi)$. For $\alpha > 0.5$ (see Fig. 1), the diffusion is normal and $P(p, t) \sim e^{-cp^2/t}/\sqrt{t}$ (c a constant). However, for $-0.5 < \alpha < 0.5$ and $t > p$, $P(p, t) \sim tp^{\gamma(\alpha)}$ has power-law tails (see inset in Fig. 1 for a relation between γ and α). Such tails are considered a signature of anomalous diffusion. They are indeed a solution of the fractional diffusion equation, $(\frac{\partial}{\partial t} - \frac{\epsilon}{2} \frac{\partial^{-\gamma-1}}{\partial |p|^{-\gamma-1}})P(p, t) = \delta(p)\delta(t)$ (see [17] for the definition of fractional derivative). In order to explicitly show that the origin of the anomalous diffusion is ex-

clusively related to the classical singularity we have repeated the calculation with the singularity-free potential $V(q) = \epsilon(|q| + b)^\alpha$, $b > 0$. As expected, the diffusion goes back to normal as in the kicked rotor with a smooth potential (see Fig. 1). We have also checked that our results are generic, namely, they do not depend on the details of the potential but only on the kind of singularity. Thus $P(p, t)$ is not altered by adding a smooth perturbation V_{per} in Eq.(1) (see inset of Fig. 2). The potential $V(q) = \epsilon \log(|q|)$ has been studied in detail since its quantum counterpart (see below) resembles a disordered conductor at the MIT. In this case (see Fig. 2) $P(p, t)$ is accurately fitted by a Lorentzian. We recall this special form of $P(p, t)$ describes the dynamics of systems with $1/f$ noise [17]. In passing we mention that in this case the classical phase space is dominated exclusively by cantori with no coexistence of chaotic and integrable regions.

Quantum mechanics—We now discuss the quantum properties of Eq.(1). For the sake of clearness let us first state our main results:

1. The eigenstates of the evolution matrix associated to the Hamiltonian Eq. (1) are power-law localized with a exponent depending only on the singularity α .
2. For $V(q) = \epsilon \log(|q|)$, the eigenstates are multifractals and the spectral correlations are described by critical statistics as at the MIT.

The quantum dynamics of a periodically kicked system is governed by the quantum evolution operator \mathcal{U} over a period T . Thus, after a period T , an initial state ψ_0 evolves to $\psi(T) = \mathcal{U}\psi_0 = e^{-\frac{i\hat{p}^2 T}{4\hbar}} e^{-\frac{iV(\hat{q})}{\hbar}} e^{-\frac{i\hat{p}^2 T}{4\hbar}} \psi_0$ where \hat{p} and \hat{q} stand for the usual momentum and position operator. Our aim is to solve the eigenvalue problem $\mathcal{U}\Psi_n = e^{-i\kappa_n/\hbar}\Psi_n$ where Ψ_n is an eigenstate of \mathcal{U} with eigenvalue κ_n . In order to proceed we first express the evolution operator in a matrix form $\langle m|\mathcal{U}|n\rangle$ in the basis of the momentum eigenstates $\{|n\rangle = \frac{e^{in\theta}}{\sqrt{2\pi}}\}$. For practical calculations one option is to make the Hilbert space finite (i.e. $m, n = 1, \dots, N$) but still keep the matrix Unitary. This can be achieved by imposing a periodic condition (with period P) to the momentum. A large period makes the phase space effectively resemble a cylinder ($P = N$ in this paper; for more details see [18]). In order to keep the kinetic term of the evolution matrix also periodic, we take $T = 2\pi M/N$ with M an integer (not a divisor of N). M is specified to make T roughly constant (≈ 0.1 in this paper) for every N used. Based on the relation between the period in momentum P and the period in position Q ($PQ = 2\pi N\hbar$), $\hbar = 1$. The resulting evolution matrix (for N odd) then reads

$$\langle m|\mathcal{U}|n\rangle = \frac{1}{N} e^{-i2\pi M n^2/N} \sum_l e^{i\phi(l, m, n)} \quad (2)$$

where $\phi(l, m, n) = 2\pi(l + \theta_0)(m - n)/N - iV(2\pi(l + \theta_0)/N)$, $l = -(N - 1)/2, \dots, (N - 1)/2$ and $0 \leq \theta_0 \leq 1$;




FIG. 3: The level spacing distribution $P(s)$ in units of the mean level spacing $s = (E_i - E_{i-1})/\Delta$. A transition is observed from Poisson to WD statistics (keeping ϵ constant) as α goes from negative to positive. In both cases $N = 3100$.




FIG. 4: The spectral rigidity $\Delta_3(L)$ versus L . Symbols are the numerical calculation for $V(q) = \epsilon \log(q)$. Lines are the analytical prediction Eq.(3). The right (left) panel corresponds to the case of (broken) time reversal invariance.

θ_0 is a parameter depending on the boundary conditions ($\theta_0 = 0$ for periodic boundary conditions). The eigenvalues and eigenvectors of \mathcal{U} can now be computed by using standard diagonalization techniques. For $\theta_0 = 0$, parity is a good quantum number and consequently states with different parity must be treated separately.

We first study the spectral fluctuations of the unfolded spectrum. The analysis of short range spectral correlations as the level spacing distribution $P(s)$ indicates (see Fig. 3) that for $\alpha > 1/2$ ($\alpha < -1/2$), $P(s)$ tends to Poisson (WD) as the volume is increased. In the intermediate region $-1/2 < \alpha < 1/2$ long range correlators as the spectral rigidity $\Delta_3(L) = \frac{2}{L^2} \int_0^L (L^3 - 2L^2x + x^3) \Sigma^2(x) dx$ where $\Sigma^2(L) = \langle L^2 \rangle - \langle L \rangle^2 = L + 2 \int_0^L (L-s) R_2(s) ds$ is the number variance and $R_2(s)$ is the two level spectral correlations function, are more suitable for a qualitative analysis due to the possibility of performing a spectral average. We restrict ourselves to the potential $V(q) = \epsilon \log(q)$ related to classical $1/f$ noise. The $R_2(s)$ associated to this type of motion has been evaluated in

Ref.[8] for systems with broken time reversal invariance,

$$R_2(s) = -K^2(s) = -\frac{\pi^2 h^2}{4} \frac{\sin^2(\pi s)}{\sinh^2(\pi^2 h s/2)} \quad (3)$$

where $h \ll 1$ is related to ϵ by $h = 1/\epsilon \ll 1$. In Fig. 4 we show $\Delta_3(L)$ for different ϵ 's and $N = 3099$. The time reversal invariance in Eq.(2) is broken by setting $T = 2\pi\beta$ with β an irrational number. As observed in Fig. 4 (left), $\Delta_3(L)$ is asymptotically linear, this feature is typical of a disordered conductor at the MIT. Moreover, the agreement with the analytical prediction based on Eq. (3) is excellent. We remark that the value of h best fitting the numerical result is within 5% of the analytical estimate $h = 1/\epsilon$. We have repeated the calculation keeping the time reversal invariance ($T = 2\pi M/N$) of Eq.(3) (Fig. 4, right). We do not have in this case an analytical result to compare with, but $\Delta_3(L)$ is also asymptotically linear as in the previous case. The eigenstates $\Psi_n = \sum \psi_n(k)|k\rangle$ of the evolution operator \mathcal{U} are also strongly affected by the classical singularity α . Instead of the exponential localization typical of the standard kicked rotor, our numerical results indicate that $|\psi_n(k)| \sim |k - k_{max}|^{-(\alpha+1)}$ decays with power-law tails (k_{max} is where $|\psi_n(k)|$ takes the maximum). This is in agreement with the findings of [9]. The eigenvector calculation was carried out in the cylindrical phase space (i.e. we switch off the periodic condition in momentum). Fig. 5 shows the results after a proper average over all N eigenstates. Remarkably, both classical (anomalous) dynamics and quantum (power-law) localization are controlled by the classical singularity α . This fact allows us to define a new universality class in quantum chaos labeled by α . For $V(q) = \epsilon \log(|q|)$, $|\psi_n(k)| \sim 1/k$ and, according to [9], the system is at the MIT. We now investigate how transport properties are affected by the non-analytical potential. We compute the quantum density of probability $P_q(k, t) = |\langle k|\phi(t)\rangle|^2$ of finding a particle with momentum $k\hbar$ at time t for a given initial state $|\phi(0)\rangle$. We restrict ourselves to the case $V(q) = \epsilon \log(|q|)$ in order to compare our findings with the ones at the Anderson transition where it is reported [19] that, in the limit $k \ll t \gg 1$, $P_q(k, t) \sim t^{-D_2} k^{D_2-1}$ where D_2 is the multifractal dimension defined previously. As expected, we have observed a similar power-law behavior in our model (see Fig. 6). The numerical value of D_2 increases as ϵ does and it is within 10% of the one obtained by the inverse participation ratio (not shown). These results show that quantum mechanically the diffusion is also anomalous but, unlike the classical case, the decay of $P_q(k, t)$ depends not only on the classical singularity α but also on the coupling constant ϵ . The overall effect of the quantum corrections is also to suppress the classical (anomalous) diffusion at a rate that increases as ϵ gets smaller. To summarize, quantum 1+1D systems with classical log singularities share all the properties of a disordered conductor at the MIT: Multifractality, critical statistics and




FIG. 5: Averaged modulus of the eigenstates of the evolution operator for $N = 2100$. We set $\epsilon = 0.01$ to a small value in order to avoid the finite size effects. The best fitting power exponents -0.49 , -0.98 and -1.36 correspond to $\alpha = -0.5, 0(\log), 0.4$ respectively are very close to the analytical prediction $-(1 + \alpha)$ [9].




FIG. 6: Time averaged $P_q(k, t)$ over $4950 \leq t \leq 5050$ and $P_q(k = 0, t)$ versus t (inset) for $V(q) = 10 \log(|q|)$. The best fitting (dotted line) is of a slope -0.34 and -0.67 (inset) respectively, which correspond to a $D_2 \approx 0.67$.

quantum anomalous diffusion. Analytical results can in principle be obtained by mapping Eq.(1) onto an 1D Anderson model. This method was introduced in [2] for the case of a kicked rotor with a smooth potential. We do not repeat here the details of the calculation but just state how the 1D Anderson model is modified by the non-analytical potential. It turns out that classical non-analyticity induces long-range disorder in the associated 1D Anderson model, $T_m u_m + \sum_r U_r u_{m+r} = E u_m$ where $U_r \sim A_r / r^{1+\alpha}$ is the Fourier coefficient of $-\tan V(q)/2$, $T_m = \theta_0/2 - Tm^2/2$ and $E = -U_0$. For random Gaussian A_r this model corresponds to the one studied in [9] which is solved by using the supersymmetry method. For

the special case of a constant non-random A_r (closer to our case) it was found that [10] some of the eigenvectors remain critical even for $1 < \alpha$. This suggests that the results reported above for $\alpha = 0$ (log potential) may be extended to other α 's though clearly more work is needed to further clarify this issue. In conclusion, we have shown that classical singularities in chaotic Hamiltonians may lead to classical anomalous diffusion and quantum power-law localization. Both quantum and classical features are governed by the classical singularity. We thus suggest a new universality class in quantum chaos labeled by the type of classical singularity. In the case of logarithmic singularities the classical dynamics presents $1/f$ noise. Quantum mechanically the system possesses all the features of a MIT: multifractal wavefunctions, critical statistics and quantum anomalous diffusion.

AMG thanks Baowen Li's group at the National University of Singapore for his warm hospitality and financial support. We also thank Baowen Li for illuminating discussions. AMG is supported by a postdoctoral fellowship from the Spanish MEC. JW is supported by DSTA Singapore under Project Agreement POD0001821.

-
- [1] O. Bohigas, M.J. Gianonni and C. Schmit, Phys. Rev. Lett. **52**, 1 (1984)
 - [2] S. Fishman, D. R. Grempel, and R. E. Prange, Phys. Rev. Lett. **49**, 509 (1982).
 - [3] H. Aoki, J. Phys. **16C**, L205 (1983).
 - [4] V.E. Kravtsov and K.A. Muttalib, Phys. Rev. Lett. **79**, 1913 (1997); S. Nishigaki, Phys. Rev. E **59**, (1999) 2853.
 - [5] B.I. Shklovskii, B. Shapiro, B.R. Sears, P. Lambrianides and H.B. Shore, Phys. Rev. B **47**, (1993) 11487.
 - [6] B.L. Altshuler, I.K. Zharekeshev, S.A. Kotochigova and B.I. Shklovskii, JETP **67** (1988) 62.
 - [7] L.S. Levitov, Phys. Rev. Lett. **64** 547 (1990).
 - [8] A. M. Garcia-Garcia, Phys.Rev. **E69** (2004) 066216.
 - [9] A.D. Mirlin, Y.V. Fyodorov, F.-M. Dittes, J. Quezada, and T.H. Seligman, Phys. Rev. E **54** 3221 (1996); E. Cuevas, et.al., Phys. Rev. Lett. **88** 016401 (2002).
 - [10] A. Rodriguez, V. A. Malyshev, G. Sierra et.al. Phys. Rev. Lett. **90**, 027404 (2004).
 - [11] J. V. Jose and R. Cordery, Phys. Rev. Lett. **56**, 290 (1986).
 - [12] B.L. Altshuler, L. S. Levitov, Phys. Rep. **288**, 487 (1997).
 - [13] D. Wintgen and H. Marxer, Phys. Rev. Lett. **60**, 971 (1988).
 - [14] B. Hu, B. Li, J. Liu and Y. Gu, Phys. Rev. Lett. **82** (1999) 4224.
 - [15] J. Liu, W. T. Cheng, C.G. Cheng, Comm. Theor. Phys. **33** 15 (2000).
 - [16] J. Klafter and G. Zumofen, Phys. Rev. E **49**, 4873 (1994).
 - [17] G.M. Zaslavsky, Phys.Rep. **371**, (2002) 461.
 - [18] F.M. Izrailev. Phys. Rep. **196**, 299 (1990).
 - [19] B. Huckestein, R. Klesse, Phys. Rev. B **59**, 9714 (1999).