Ising model susceptibility: Fuchsian differential equation for $\chi^{(4)}$ and its factorization properties

N. Zenine[§], S. Boukraa[†], S. Hassani[§] and J.-M. Maillard[‡]

§Centre de Recherche Nucléaire d'Alger,
2 Bd. Frantz Fanon, BP 399, 16000 Alger, Algeria
†Université de Blida, Institut d'Aéronautique, Blida, Algeria
‡ LPTL, Université de Paris 6, Tour 24, 4ème étage, case 121,
4 Place Jussieu, 75252 Paris Cedex 05, France

 $E{-}mail{:}\ maillard@lptl.jussieu.fr, sboukraa@wissal.dz, njzenine@yahoo.com\\$

Abstract. We give the Fuchsian linear differential equation satisfied by $\chi^{(4)}$, the "four-particle" contribution to the susceptibility of the isotropic square lattice Ising model. This Fuchsian differential equation is deduced from a series expansion method introduced in two previous papers and is applied with some symmetries and tricks specific to $\chi^{(4)}$. The corresponding order ten linear differential operator exhibits a large set of factorization properties. Among these factorizations one is highly remarkable: it corresponds to the fact that the two-particle contribution $\chi^{(2)}$ is actually a solution of this order ten linear differential operator. This result, together with a similar one for the order seven differential operator corresponding to the three-particle contribution, $\chi^{(3)}$, leads us to a conjecture on the structure of all the n-particle contributions $\chi^{(n)}$.

PACS: 05.50.+q, 05.10.-a, 02.30.Hq, 02.30.Gp, 02.40.Xx**AMS Classification scheme numbers**: 34M55, 47E05, 81Qxx, 32G34, 34Lxx, 34Mxx, 14Kxx

Key-words: Susceptibility of the Ising model, series expansions, Fuchsian differential equations, factorization of linear differential operators, \mathcal{D} -module, equivalence of differential operators, apparent singularities, indicial equations.

1. Introduction

The magnetic susceptibility of square lattice Ising model, can be written [1] as an infinite sum

$$\chi(T) = \sum_{n=1}^{\infty} \chi^{(n)}(T) \tag{1}$$

of individual contributions, with the odd (respectively even) n corresponding to high (respectively low) temperature case. These individual contributions are (n-1)-dimensional integrals [2, 3, 4, 5, 6, 7], and are seen as successive n-particle contributions to the susceptibility [1].

To get an understanding of the analytical structure of χ , two approaches are usually taken nowadays. One approch taking into account a fundamental non-linear symmetry, namely non-linear Painlevé difference equations [8, 9, 10, 11, 12, 13]. provides a series expansion for the whole susceptibility χ . With this method coefficients of χ were recently generated [8].

The second approach considers the individual n-particle excitations as given by (n-1)-dimensional integrals. Isotropic series coefficients are generated [6, 7, 8]. This latter method allows one to seek the differential equations satisfied by the $\chi^{(n)}$, since they are D-finite contrary to the whole susceptibility χ for which there are strong indications that it has a natural boundary in the complex plane of the variable s = sh(2K), where K = J/kT is the usual Ising model coupling constant. This has been shown for the isotropic case [6] and for the anisotropic case [14, 15, 16]. Such a function cannot be D-finite.

The understanding of the magnetic susceptibility may then require the knowledge of each (or some) of the individual contributions. This knowledge can be in the form of a closed expression, as is the case for $\chi^{(1)}$ and $\chi^{(2)}$ or in the form of a differential equation as found for $\chi^{(3)}$ [17, 18]. The last case was far from being obvious and has required the building of an original method of expansion. The use of a remarkable formula allowed us to give the series expansion in the temperature variable (or a closely related variable) where the (n-1)-dimensional integrals have been fully performed.

In this paper, we continue to use this expansion method to tackle the next individual contribution, namely, $\chi^{(4)}$. We should note that, although, the method is general and applicable for high or low temperature and for any n, some of the tricks and tools used may be specific for a given $\chi^{(n)}$. In Section 2, we present the basic features of the expansion method that allow us to obtain the fully integrated $\chi^{(4)}$ as four sums of products of four hypergeometric functions, without any numerical approximation. In Section 3, we give the homogeneous Fuchsian linear differential equation satisfied by $\chi^{(4)}$. Section 4 contains some remarkable algebraic properties of this differential equation. Finally, Section 5 contains our conclusions.

2. Fully integrated $\tilde{\chi}^{(4)}$ expansion

2.1. The expansion method

Let us focus on the fourth contribution to the susceptibility χ defined by the triple integral as given in [7]

$$\chi^{(4)} = (1 - s^{-4})^{1/4} \cdot \tilde{\chi}^{(4)}$$

$$\tilde{\chi}^{(4)} = \int_{0}^{2\pi} \frac{d\phi_1}{2\pi} \int_{0}^{2\pi} \frac{d\phi_2}{2\pi} \int_{0}^{2\pi} \frac{d\phi_3}{2\pi} \cdot \tilde{y}_1 \tilde{y}_2 \tilde{y}_3 \tilde{y}_4 \cdot R^{(4)} \cdot H^{(4)}$$
(2)

with

$$R^{(4)} = \frac{1 + \tilde{x}_1 \tilde{x}_2 \tilde{x}_3 \tilde{x}_4}{1 - \tilde{x}_1 \tilde{x}_2 \tilde{x}_3 \tilde{x}_4} \tag{3}$$

$$H^{(4)} = \frac{1}{4!} \prod_{i < j} \frac{\tilde{x}_i \tilde{x}_j}{(1 - \tilde{x}_i \tilde{x}_j)^2} \cdot (Z_i - Z_j)^2$$
 (4)

$$Z_n = \exp(i\phi_n), \qquad n = 1, \dots, 4$$
 (5)

$$\phi_1 + \phi_2 + \phi_3 + \phi_4 = 0 \tag{6}$$

$$\tilde{x}_n = \frac{s}{1 + s^2 - s\cos\phi_n + \sqrt{(1 + s^2 - s\cos\phi_n)^2 - s^2}},$$

$$\tilde{y}_n = \frac{s}{\sqrt{(1 + s^2 - s\cos\phi_n)^2 - s^2}}, \qquad n = 1, \dots, 4$$
(7)

Instead of the variable s, we found it more suitable to use $w = \frac{1}{2}s/(1+s^2)$ which has, by construction, Kramers-Wannier duality invariance $(s \leftrightarrow 1/s)$ and thus allows us to deal with both limits (high and low temperature, small and large s) on an equal footing [17, 18]. It is also convenient to consider the scaled variables

$$x_n = \frac{\tilde{x}_n}{w} = \frac{2}{1 - 2w\cos\phi_n + \sqrt{(1 - 2w\cos\phi_n)^2 - 4w^2}},$$

$$y_n = \frac{\tilde{y}_n}{2w} = \frac{1}{\sqrt{(1 - 2w\cos\phi_n)^2 - 4w^2}}$$
(8)

which behave like 1 + O(w) at small w.

As performing the integrals in (2) is highly non trivial, we apply the "expansion method" previously described in [17, 18], where the key ingredient was the Fourier expansion of $y x^n$, a quantity appearing in any $\chi^{(p)}$.

This remarkable formula, for yx^n , that carries only one summation index reads

$$yx^{n} = a(0,n) + 2\sum_{k=1}^{\infty} w^{k} a(k,n) \cos k\phi$$
$$= \sum_{k=-\infty}^{\infty} A(k,n) Z^{k} = \sum_{k=-\infty}^{\infty} A(k,n) Z^{-k}$$
(9)

with

$$A(k,n) = A(-k,n) = w^{|k|} a(|k|,n)$$
(10)

where a(k, n) is a non terminating hypergeometric series that reads

$$a(k,n) = \binom{m}{k}. \tag{11}$$

$$_{4}F_{3}\left(\frac{(1+m)}{2},\frac{(1+m)}{2},\frac{(2+m)}{2},\frac{(2+m)}{2};1+k,1+n,1+m;16w^{2}\right)$$

where m = k + n. Note that a(k, n) = a(n, k).

The integrand of $\tilde{\chi}^{(4)}$ is expanded in the various variables x_j , instead of the variable w. In this framework, with the help of the Fourier expansion (9), the angular integration becomes straightforward as was shown in [18] for the $\tilde{\chi}^{(3)}$ case (see below for the $\tilde{\chi}^{(4)}$ case).

2.2. Calculation of $\tilde{\chi}^{(4)}$

With $H^{(4)}$ taken as in (4), $\tilde{\chi}^{(4)}$ will be given by *five* summations on products of four hypergeometric functions. This is shown in Appendix A. This route is feasible for any $\chi^{(n)}$ and does not use any symmetry or tricks specific to the considered $\chi^{(n)}$.

In the following, we use alternatively a simplified form of $H^{(4)}$ equivalent, for integration purposes, to (4) such that $\tilde{\chi}^{(4)}$ will be expressed by only four summations (i.e., one summation less compared to the expression (88) given in Appendix A). Let us just sketch the salient steps of this calculation. The details are left to Appendices B, C and D.

With the help of the key relation

$$(Z_i - Z_j) \frac{\tilde{x}_i \tilde{x}_j}{1 - \tilde{x}_i \tilde{x}_j} = -(\tilde{x}_i - \tilde{x}_j) \frac{Z_i Z_j}{1 - Z_i Z_j}$$

$$(12)$$

the quantity $H^{(4)}$ becomes †:

$$H^{(4)} = \frac{1}{4!} \prod_{i < j} \frac{\tilde{x}_i - \tilde{x}_j}{1 - \tilde{x}_i \tilde{x}_j} \cdot \prod_{i < j} \frac{Z_i - Z_j}{1 - Z_i Z_j}.$$
 (13)

Using the symmetry of the rest of the integrand in the angular variables, the quantity $H^{(4)}$ can be written (see Appendix B) as‡

$$H^{(4)} \equiv -\frac{1}{8}(Z_1 - Z_2)(Z_3 - Z_4) \frac{Z_1 Z_2}{(1 - Z_1 Z_2)^2} \cdot \left(P_{12-34}^{(x)} - (1 \leftrightarrow 3) - (2 \leftrightarrow 3)\right)$$
(14)

where:

$$P_{12-34}^{(x)} = (\tilde{x}_1 - \tilde{x}_2)(\tilde{x}_3 - \tilde{x}_4) \cdot \left(\frac{\tilde{x}_1 \tilde{x}_2}{1 - \tilde{x}_1 \tilde{x}_2} \frac{\tilde{x}_3 \tilde{x}_4}{1 - \tilde{x}_3 \tilde{x}_4} + \frac{(\tilde{x}_1 \tilde{x}_2)^2}{1 - \tilde{x}_1 \tilde{x}_2} + \frac{(\tilde{x}_3 \tilde{x}_4)^2}{1 - \tilde{x}_3 \tilde{x}_4}\right).$$
(15)

Taking the expression (14) for $H^{(4)}$, expanding the integrand in the x_j variables, one notes that the integrand depends only on combinations of the form

$$\left(\prod_{i=1}^{4} y_i \cdot x_i^{n_i}\right) \cdot (Z_1 - Z_2)(Z_3 - Z_4) \frac{Z_1 Z_2}{(1 - Z_1 Z_2)^2} \tag{16}$$

which have simple integration rules (see Appendix C) and, thus, the problem of angular integrations is settled.

Finally, the expansion method described above, together with the form (14) of $H^{(4)}$, allow us to obtain $\tilde{\chi}^{(4)}(w)$ as a fully integrated expansion in the form (see Appendix D)

$$\tilde{\chi}^{(4)} = 16 \, w^{16} \cdot \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} w^{8m+4k+4n+2j} \cdot (2m+1)(2m+2k+1)$$

$$(1+\theta(j-1)) \cdot (1+\theta(k-1)) \cdot \frac{1}{2} \Big(V(m,m+k,n,n+j) + V(m,m+k,n+j,n) \Big)$$
(17)

where $\theta(x)$ is the step function defined as

$$\theta(x) = \begin{cases} 1, & x \ge 0\\ 0, & x < 0 \end{cases} \tag{18}$$

and

$$V(m,k,n,j) = d(m,m+1;j+k+2) \cdot d(n+m+2,n+m+3;k)$$

$$+ d(m,m+1;k) \cdot d(n+m+2,n+m+3;j+k+2)$$

$$+ d(m,n+m+3;j+k+2) \cdot d(m+1,n+m+2;k)$$

$$+ d(m,n+m+3;k) \cdot d(m+1,n+m+2;j+k+2)$$

$$- d(m,n+m+2;j+k+2) \cdot d(m+1,n+m+3;k)$$

$$- d(m,n+m+2;k) \cdot d(m+1,n+m+3;j+k+2)$$

$$(19)$$

- † We have used the constraint (6) in the form $\prod_i Z_i = 1$.
- ‡ Throughout this paper the notation \equiv stands for equality for integration purposes.

with:

$$d(n_1, n_2; k) = a(n_1, k+1) a(n_2, k) - a(n_1, k) a(n_2, k+1).$$
(20)

2.3. Series generation

Note that the summand in equation (17) depends only on combinations of $C_d(n_1, n_2, k, j)$ which is the coefficient of w^{2j} in the expansion of $d(n_1, n_2; k)$. It is given by

$$C_d(n_1, n_2, k, j) = \sum_{i=0}^{j} \left(C_a(n_1, k+1, i) C_a(n_2, k, j-i) - C_a(n_1, k, i) C_a(n_2, k+1, j-i) \right)$$
(21)

where $C_a(n,k,i)$ is the coefficient of w^{2i} in the expansion of a(n,k) which reads:

$$C_a(n,k,i) = \binom{n+k+2i}{i} \binom{n+k+2i}{i+k}$$
(22)

From our integrated form of $\tilde{\chi}^{(4)}$, the generation of series coefficients becomes straightforward. Recall that (17) is already integrated, and, thus, the computing time to obtain the series coefficients comes from the evaluation of the sums. For the forms used, this time is of order N^7 . Improvements can be made by optimal data storage in order to avoid repeated summation evaluations. Actually, we have found it more efficient to store the coefficients C_d .

We have been able to generate, from formal calculations, a long series of coefficients from the expression (17) up to order§ 432:

$$\frac{\tilde{\chi}^{(4)}(w)}{16\,w^{16}} = 1 + 64\,w^2 + 2470\,w^4 + 74724\,w^6 + \dots + O(w^{434}) \quad (23)$$

Note that our formal calculation program has been rewritten by J. Dethridge into an optimized C++ program that can give the series in few hours using very little memory.

3. The Fuchsian differential equation satisfied by $\tilde{\chi}^{(4)}$

It is clear from the expression (17) that $\tilde{\chi}^{(4)}$ is even in w. We thus introduce, in the following, the variable $x=16\,w^2$. With our long series, and with a dedicated program, we have succeeded in obtaining the differential equation for $\tilde{\chi}^{(4)}$ that is given by (with $x=16\,w^2$)

$$\sum_{n=1}^{10} a_n(x) \cdot \frac{d^n}{dx^n} F(x) = 0 \tag{24}$$

[§] More precisely we obtained this series up to order 368 with a 2 Giga-memory computer, up to order 390 with a 3 Giga-memory computer, and up to order 432 with a 8 Giga-memory computer of the stix laboratory at the Ecole Polytechnique (medicis platform for formal calculations).

^{||} Private communication.

[§] This is also the case for any $\tilde{\chi}^{(2n)}$ (see for instance [7]).

with

$$a_{10} = -512 x^{6} (x - 4) (1 - x)^{6} P_{10}(x),$$

$$a_{9} = 256 (1 - x)^{5} x^{5} P_{9}(x), \quad a_{8} = -384 (1 - x)^{4} x^{4} P_{8}(x),$$

$$a_{7} = 192 (1 - x)^{3} x^{3} P_{7}(x), \quad a_{6} = -96 (1 - x)^{2} x^{2} P_{6}(x),$$

$$a_{5} = 144 (1 - x) x P_{5}(x), \quad a_{4} = -72 P_{4}(x), \quad a_{3} = -108 P_{3}(x)$$

$$a_{2} = -54 P_{2}(x) \quad a_{1} = -27 P_{1}(x)$$
(25)

where $P_{10}(x)$, $P_{9}(x)$..., $P_{1}(x)$ are polynomials of degree respectively 17, 19, 20, 21, 22, 23, 24, 23, 22 and 21 given in Appendix E.

With (25), the differential equation needs 242 unknowns to be found (counting the polynomial in front of the derivative of order 0, which is identically null). Our series expansion for $\tilde{\chi}^{(4)}/16w^{16}$, having only 217 terms in the variable $x=16w^2$, is, thus, not long enough to let the differential equation be found. This calls for some comments.

The differential equation (24) is of minimal order. It is obvious that one can obtain other differential equations of greater order. As explained in [18] (see Section 4), before the differential equation built from a series expansion pops out, the singularities computed as roots of the polynomial in front of the highest derivative reach stabilized numerical values as the degrees of the polynomials and/or the order get higher. Even the minimal multiplicity of the singularity can be seen. This leads us to take the following form for the $\tilde{\chi}^{(4)}$ differential equation

$$\sum_{i=0}^{q} R_i(x) \cdot x^i (1-x)^i \cdot \frac{d^i}{dx^i} F(x) = 0$$
 (26)

where only the physical singularities are explicitly included. Note that the other (non-apparent) singularity can also be included. If the differential equation can be identified with the number of series coefficients at hand, the unnecessary terms will factor out. A compromise has to be found with respect to how long the series is. Now, there are many ways to choose the degrees of the polynomials $R_i(x)$. We have taken

$$deg(R_i(x)) = \mu + q - i, \qquad i = 0, 1, \dots, q$$
(27)

in order to have the point at infinity as a regular singular point.

If the differential equation of order q and degree μ exists, the series of $\tilde{\chi}^{(4)}/16w^{16}$ should have at least N terms, with :

$$N = \mu \cdot (q+1) + \frac{1}{2} q \cdot (q+3). \tag{28}$$

Staying below the above hyperbola, we have obtained, at q=11 and $\mu=9$ which requires 185 terms in the series, two linearly independent differential equations that satisfy the remaining 30 terms of the series. The combination of these two differential equations gives (24) which has been checked to be of minimal order. Note the fact that besides the Fuchsian differential equation of minimal order with an apparent polynomial, there are other differential equations of higher order that require less terms in the series to be identified. We plan to report on this feature elsewhere.

The singularities of this differential equation x = 0, 1, 4 and $x = \infty$ are all regular singular points. The roots of the polynomial $P_{10}(x)$ are apparent singularities. One

|| This feature can easily be seen, making the change of variable t = 1/x and looking for the necessary condition for t = 0 to be a regular singular point from which (27) is deduced.

notes, contrary, to $\chi^{(3)}$ that no new singularity is found besides the known physical and non physical singularities (i.e., Nickel's [6, 7]). The critical exponents of all these singular points are given in Table 1 with the maximum power of logarithmic terms in the solutions. As was the case for $\chi^{(3)}$, these logarithmic terms appear due to the multiple roots of the indicial equation. A noteworthy remark is the occurrence of logarithmic terms up to the power 3 for $\chi^{(4)}$ to be compared with the power 2 for $\chi^{(3)}$ at the singular points $w=\pm 1/4$ and $w=\infty$.

x-singularity	s-singularity	Critical exponents in x	P
0	$0,\infty$	8, 3, 3, 2, 2, 1, 1, 0, 0, -1/2	1
1	±1	3, 2, 1, 1, 0, 0, 0, 0, -1, -3/2	3
4	$\pm \frac{1}{2} \pm i \frac{\sqrt{3}}{2}$	8, 7, 13/2, 6, 5, 4, 3, 2, 1, 0	0
∞	$\pm i$	5/2, 3/2, 3/2, 1/2, 1/2, 1/2, 1/2, 0, -1/2, -1/2	3
x_P , 17 roots	s_P , 68 roots	10, 8, 7, 6, 5, 4, 3, 2, 1, 0	0

Table 1: Critical exponents for each regular singular point. P is the maximum power of the logarithmic terms for each singularity. x_P is any of the 17 roots of $P_{10}(x)$.

It is worth recalling the Fuchsian relation on Fuchsian type equations. Denoting by $x_1, x_2, \dots, x_m, x_{m+1} = \infty$, the regular singular points of a Fuchsian type equation of order q and $\rho_{j,1}, \dots, \rho_{j,q}$ $(j = 1, \dots, m+1)$ the q roots of the indicial equation [28, 29] corresponding to each regular singular point w_j , the following Fuchsian relation [28, 29] holds:

$$\sum_{j=1}^{m+1} \sum_{k=1}^{q} \rho_{j,k} = \frac{(m-1) \ q \ (q-1)}{2}.$$
 (29)

The number of regular singular points here is m+1=21 corresponding respectively to the 17 roots of P_{10} , the x=0,1,4, regular singular points, and the point at infinity $x=\infty$. The Fuchsian relation (29) is actually satisfied here with q=10, m=20.

Considering the Fuchsian relation (29), but now in the variable s, one first remarks that the $\rho_{j,k}$ exponents are the same as the ones in the x variable for the 68 roots of the apparent polynomial, as well as for the four x=4 singularities (namely $s=\pm\frac{1}{2}\pm i\frac{\sqrt{3}}{2}$), but are multiplied by a factor two for all the other ones. This is obvious from the definition of $x=4s^2/(1+s^2)^2$ at x=0 and $x=\infty$, and this can be seen for x=1 from:

$$1 - x = \left(\frac{1 - s^2}{1 + s^2}\right)^2 \tag{30}$$

The Fuchsian relation is actually satisfied in the s variable, but now, with q=10, m=77.

Let us close this section by confirming the dominant singular behavior at x=4 and $x=\infty$ given in [7], since this can easily be done from the critical exponents given in Table 1. Near x=4, the solution behaves as $t^{13/2}$ and near $x=\infty$ as $t^{-1/2}\log(t)$, with respectively t=4-x and t=1/x. Fom Table 1, the subdominant singular behavior at $x=\infty$ is $t^{3/2}\log(t)$ and $t^{1/2}\log^k(t)$ (with k=1,2,3).

4. Properties of the Fuchsian differential equation (24)

From Table 1, and from the formal solutions around the singular points, it is easy to find the following simple solutions of (24):

$$S_0(x) = constant \tag{31}$$

$$S_1(x) = \frac{8 - 12x + 3x^2}{8(1 - x)^{3/2}},\tag{32}$$

$$S_2(x) = \frac{2 - 6x + x^2}{2(1 - x)\sqrt{x}}. (33)$$

These solutions correspond to solutions of some differential operators of order one. Let us call these order one differential operators respectively L_0 , L_1 and L_2 .

A remarkable finding is the following solution of the Fuchsian differential equation (24):

$$S_3(x) = \frac{1}{64} x^2 \cdot {}_2F_1\left(\frac{5}{2}, \frac{3}{2}; 3; x\right) \tag{34}$$

which is nothing but the two-particle contribution to the magnetic susceptibility, i.e., $\tilde{\chi}^{(2)}$ associated with an operator of order two, N_0 ($N_0(S_3) = 0$). We will come back to this point later.

The second solution of the order-two operator N_0 is given in terms of the MeijerG function [19]:

$$\tilde{S}_3(x) = \frac{\pi}{2} \text{MeijerG}([[], [1/2, 3/2]], [[2, 0], []], x)$$
 (35)

which can also be written as

$$\tilde{\mathcal{S}}_3(x) = \mathcal{S}_3(x) \log(x) + B(x) \tag{36}$$

with:

$$B(x) = \frac{1}{12\pi} \cdot \sum_{k=0}^{\infty} x^k \frac{d}{dk} \left(\frac{\Gamma(k-1/2)\Gamma(k+1/2)}{\Gamma(k-1)\Gamma(k+1)} \right)$$
(37)

With these five solutions corresponding to three differential operators of order one, and one differential operator of order two, it is easy to construct 24 factorizations of \mathcal{L}_{10} , the differential operator corresponding to the Fuchsian differential equation (24), which can be written¶ as:

$$\mathcal{L}_{10} = O_5 \cdot G(N). \tag{38}$$

G(N) is a shorthand notation of a differential operator of order 5, factorizable in one operator of order two and three operators of order 1. G(N) has 24 different factorizations involving eight differential operators of order two and 24 operators of

 \P All the operators are such that the coefficient in front of the highest derivative is +1.

order one. The differential operator O_5 factorizes as $M_1 \cdot L_{24}$, i.e., one operator of order four and one operator of order one. Let us give two examples of the 24 factorizations⁺:

$$\mathcal{L}_{10} = M_1 \cdot L_{24} \cdot N_4 \cdot L_{12} \cdot L_3 \cdot L_0 \tag{39}$$

$$\mathcal{L}_{10} = M_1 \cdot L_{24} \cdot L_{13} \cdot N_6 \cdot L_3 \cdot L_0 \tag{40}$$

This large number of factorizations [22, 31, 21] induces the occurrence of intertwiners \dagger . In the examples above, one has $N_4 \cdot L_{12} = L_{13} \cdot N_6$. Seeking a similar relation for $L_{24} \cdot N_4$, one finds $N_9 \cdot L_{25}$. This last factorization introduces six factorizations that we denote as:

$$\mathcal{L}_{10} = M_1 \cdot N_9 \cdot G(L). \tag{41}$$

G(L) is a notation for an operator of order four, factorizable as four operators of order one.

One factorization of G(L) reads:

$$G(L) = L_{25} \cdot L_{12} \cdot L_3 \cdot L_0. \tag{42}$$

This differential operator G(L) that factorizes \mathcal{L}_{10} at right, obviously has \mathcal{S}_0 , \mathcal{S}_1 and \mathcal{S}_2 as solutions. The fourth solution (of the order four differential operator G(L)) can be obtained by order reduction. It reads

$$S_4(x) = \frac{4(x-2)\sqrt{4-x}}{x-1} + 16\log\frac{x}{(2+\sqrt{4-x})^2} + 16S_1(x)\cdot\log g(x) - 16\sqrt{x}S_2(x)\cdot{}_2F_1\left(\frac{1}{2},\frac{1}{2};\frac{3}{2};\frac{x}{4}\right)$$
(43)

with:

$$g(x) = \frac{1}{x} \left((8 - 9x + 2x^2) + 2(2 - x) \cdot \sqrt{(1 - x)(4 - x)} \right)$$
 (44)

To get more factorizations, we need to obtain simple solutions of \mathcal{L}_{10}^* , the adjoint of the differential operator \mathcal{L}_{10} . There is no solution of \mathcal{L}_{10}^* corresponding to an order-one operator, however, we have been able to find a solution corresponding to an operator of order two (denoted N_8^*). This solution of N_8^* is a combination of elliptic integrals with polynomials of quite large degrees and reads

$$S_1^*(x) = \frac{x(1-x)^6(4-x)}{3840000 \cdot P_{10}(x)} \cdot \left(q_1(x)K(x) + q_2(x)E(x)\right) \tag{45}$$

where

$$K(x) = {}_{2}F_{1}\left(\frac{1}{2}, \frac{1}{2}; 1; x\right), \qquad E(x) = {}_{2}F_{1}\left(-\frac{1}{2}, \frac{1}{2}; 1; x\right)$$
 (46)

and:

$$q_1(x) = (1-x)(47352014438400 - 246257318625280 x + 275880211382272 x^2 + 68328139784192 x^3 + 645943284072448 x^4 - 2774821715853312 x^5 + 3217221650489344 x^6 - 683914539437568 x^7$$

 $^{^+}$ We denote by L the operators of order 1, by N, the operators of order 2 and by M the operators of order 4 (see Appendix F).

[†] We thank Jacques-Arthur Weil for usefull comments on the equivalence of linear differential operators [20] (see equation (5) in [21]).

[‡] We did not look for the second solution of N_8^* , our purpose being the factorization of \mathcal{L}_{10} .

$$-2042467767948624 x^{8} + 3083863919521506 x^{9}$$

$$-2746206480894969 x^{10} + 1558224994851490 x^{11}$$

$$-347705392468761 x^{12} - 145625559012638 x^{13}$$

$$+117842186745065 x^{14} - 30744722745590 x^{15}$$

$$+3089482306025 x^{16} + 18651488480 x^{17}$$

$$-21574317760 x^{18} + 821760000 x^{19})$$

$$q_{2}(x) = -47352014438400 + 269933325844480 x$$

$$-390130367987712 x^{2} + 27877970018304 x^{3}$$

$$-580571855978496 x^{4} + 3135069528473600 x^{5}$$

$$-4488375407386624 x^{6} + 1736922901371392 x^{7}$$

$$+2600277912748368 x^{8} - 5123144341863018 x^{9}$$

$$+5224617790090830 x^{10} - 3547418998359865 x^{11}$$

$$+1453586336314895 x^{12} - 273126255420088 x^{13}$$

$$-8194519962996 x^{14} + 11308926014655 x^{15}$$

$$-1235672485785 x^{16} - 60982101700 x^{17}$$

$$+15765744320 x^{18} - 607865600 x^{19}$$

$$-3840000 x^{20}$$

$$(48)$$

This order 2 operator N_8 completes the factorization scheme of \mathcal{L}_{10} , the differential operator of order ten. Let us recap the factorizations

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot G(L)$$

$$\mathcal{L}_{10} = M_1 \cdot N_9 \cdot G(L)$$

$$\mathcal{L}_{10} = M_1 \cdot L_{24} \cdot G(N)$$
(49)

G(L) is a differential operator of order four, factorizable in order one operators. It has six different factorizations involving thirteen differential operators of order one. G(N) is an operator of order five, factorizable in one operator of order two and three operators of order one. G(N) has 24 different factorizations involving eight differential operators of order two, and 24 order one operators with twelve appearing in G(L).

All these 36 factorizations (49) are given in Appendix F. Appendix F shows that these 36 factorizations can be considered as *only one* factorization up to a set of equivalence-symmetries [20, 21, 22, 23, 24, 25, 26].

From the ten solutions of the differential equation (24), six solutions are given explicity, S_0 , S_1 , S_2 , S_4 (this last being a solution of an order four operator) and two solutions (S_3 , \tilde{S}_3) corresponding to the differential equation of $\tilde{\chi}^{(2)}$. The remaining four solutions are those of the operator of order eight, $M_2 \cdot G(L)$.

From the 36 factorizations shown in Appendix F, those six of the form $N_8 \cdot M_2 \cdot G(L)$ are of the most importance. Their occurrence allows us to get the contribution α of $\tilde{\chi}^{(2)}$ in the physical solution $\tilde{\chi}^{(4)}$ from $M_2 \cdot G(L)(\tilde{\chi}^{(4)} - \alpha \tilde{\chi}^{(2)}) = 0$. This contribution is obtained easily and gives

$$\tilde{\chi}^{(4)} = \frac{1}{3}\tilde{\chi}^{(2)} + \Phi_4 \tag{50}$$

where Φ_4 is solution of the order eight differential operator $M_2 \cdot G(L)$.

Recall that the same situation occurred for the differential equation of order seven for $\tilde{\chi}^{(3)}$. One can see that the rational solution w/(1-4w) occurring§ in the differential equation of order seven for $\tilde{\chi}^{(3)}$ is nothing but $\tilde{\chi}^{(1)}$. With this remark we can rewrite a decomposition of $\tilde{\chi}^{(3)}$ we gave in [17, 18] as follows:

$$\tilde{\chi}^{(3)} = \frac{1}{6}\tilde{\chi}^{(1)} + \Phi_3. \tag{51}$$

where Φ_3 is solution of the order six differential operator (noted L_6 in [17, 18]).

5. Comments and speculations

Denoting \mathbf{L}_4 the order ten differential operator associated with the ordinary differential equation satisfied by $\tilde{\chi}^{(4)}$, and more generally, \mathbf{L}_n the differential operators associated with the ordinary differential equation satisfied by $\tilde{\chi}^{(n)}$, one has:

$$\mathbf{L}_4(\tilde{\chi}^{(4)}) = \mathbf{L}_4(\tilde{\chi}^{(2)}) = 0.$$
 (52)

Furthermore, one can see that the rational solution w/(1-4w), occurring in the differential equation of order seven for $\tilde{\chi}^{(3)}$, is nothing but $\tilde{\chi}^{(1)}$ and thus one has:

$$\mathbf{L}_{3}(\tilde{\chi}^{(3)}) = \mathbf{L}_{3}(\tilde{\chi}^{(1)}) = 0. \tag{53}$$

Both relations come from the factorizations of the differential operators corresponding to $\tilde{\chi}^{(3)}$ and $\tilde{\chi}^{(4)}$.

At this point, it is tempting to make some conjectures generalizing relations (52, 53) and relations (50,51).

From (52) and (53) the conjecture is

$$\mathbf{L}_{2n+1}(\tilde{\chi}^{(2n+1)}) = \mathbf{L}_{2n+1}(\tilde{\chi}^{(1)}) = 0, \mathbf{L}_{2n}(\tilde{\chi}^{(2n)}) = \mathbf{L}_{2n}(\tilde{\chi}^{(2)}) = 0$$
(54)

meaning that the differential operator of $\chi^{(1)}$ (respectively $\chi^{(2)}$) right divides the differential operator corresponding to $\chi^{(2n+1)}$ (respectively $\chi^{(2n)}$).

One stronger conjecture is to expect the same situation as in (50,51) occurring in the higher particle contributions, i.e.,

$$\tilde{\chi}^{(2n)} = \alpha_{2n} \,\tilde{\chi}^{(2)} + \Phi_{2n},
\tilde{\chi}^{(2n+1)} = \alpha_{2n+1} \,\tilde{\chi}^{(1)} + \Phi_{2n+1}$$
(55)

where Φ_{2n} (respectively Φ_{2n+1}) is solution of a differential operator that right divides the differential operator \mathbf{L}_{2n} (respectively \mathbf{L}_{2n+1}) and is not divisible by the differential operator \mathbf{L}_2 (respectively \mathbf{L}_1). In this situation, it is easy to obtain the numbers α_n which give the contribution of $\tilde{\chi}^{(1)}$ and $\tilde{\chi}^{(2)}$ in the higher $\tilde{\chi}^{(n)}$'s as explained for $\tilde{\chi}^{(4)}$ (see text before (50)).

Let us note that both conjectures are free from any constraint due to the singularities that occur in the differential equations, since any $\tilde{\chi}^{(2n+1)}$ (respectively $\tilde{\chi}^{(2n)}$) has the singularities occurring in $\tilde{\chi}^{(1)}$ (respectively $\tilde{\chi}^{(2)}$).

A much stronger conjecture is to expect any $\chi^{(m)}$ to be "embedded" in any $\chi^{(n)}$ where m divides n with same odd-even parity.

$$\mathbf{L}_{n}(\tilde{\chi}^{(n)}) = \mathbf{L}_{n}(\tilde{\chi}^{(m)}) = 0,$$

$$\mathbf{L}_{n} = \mathbf{L}_{n}^{(m)} \cdot \mathbf{L}_{m}.$$
(56)

§ This corresponds to the solution S_1 given in [17, 18] which is $S_1 = \tilde{\chi}^{(1)}/2$.

This means that \mathbf{L}_n might be built from the least common left multiple (lclm) of the differential operators associated with the \mathbf{L}_m 's where m divides the integer n respecting even-odd parity. The fact that n should be multiple of m is due to the non physical singularities appearing in the differential equations. For instance, no "embedding" like $\chi^{(3)}$ being solution of the differential equations of $\chi^{(5)}$ has to be expected. Whether $\chi^{(3)}$ is a solution of the differential equation satisfied by, e.g., $\chi^{(9)}$ is not ruled out, the (non-apparent) singularities of the first being (non-apparent) singularities of the last. But, then, the new singularities discovered [17] for $\chi^{(3)}$ have to occur in the differential equation of $\chi^{(9)}$.

Similarly one can expect, with this last conjecture, the following relations for \mathbf{L}_{12} and \mathbf{L}_{15} (associated with $\tilde{\chi}^{(12)}$ and $\tilde{\chi}^{(15)}$)

$$\mathbf{L}_{12}(\tilde{\chi}^{(12)}) = \mathbf{L}_{12}(\tilde{\chi}^{(6)}) = \mathbf{L}_{12}(\tilde{\chi}^{(4)}) = \mathbf{L}_{12}(\tilde{\chi}^{(2)}) = 0$$

$$\mathbf{L}_{15}(\tilde{\chi}^{(15)}) = \mathbf{L}_{15}(\tilde{\chi}^{(5)}) = \mathbf{L}_{15}(\tilde{\chi}^{(3)}) = \mathbf{L}_{15}(\tilde{\chi}^{(1)}) = 0$$
(57)

and thus:

$$\mathbf{L}_{12} = \mathbf{L}_{12}^{(6)} \cdot \mathbf{L}_{6} = \mathbf{L}_{12}^{(4)} \cdot \mathbf{L}_{4} = \mathbf{L}_{12}^{(2)} \cdot \mathbf{L}_{2}$$

$$\mathbf{L}_{15} = \mathbf{L}_{15}^{(5)} \cdot \mathbf{L}_{5} = \mathbf{L}_{15}^{(3)} \cdot \mathbf{L}_{3} = \mathbf{L}_{15}^{(1)} \cdot \mathbf{L}_{1}.$$
(58)

For instance, finding the differential operator satisfied by $\tilde{\chi}^{(15)}$ would enable us to see if the differential operator \mathbf{L}_{15} actually has some relation with the least common left multiple of \mathbf{L}_1 , \mathbf{L}_3 and \mathbf{L}_5 .

6. Conclusion

Considering the isotropic Ising square lattice model susceptibility, we extended our "expansion method" (that allowed us to find the differential equation satisfied by $\chi^{(3)}$), to the four-particle contribution to the susceptibility, namely $\chi^{(4)}$. We first obtained $\chi^{(4)}$ as a fully integrated multisum on products of four hypergeometric functions, and obtained a long series for $\chi^{(4)}$. From this long series, we gave the Fuchsian differential equation of order ten satisfied by $\chi^{(4)}$. This differential equation has a rich structure in terms of factorizations.

We have given in closed form six, of the ten solutions of the differential equation (24), namely, S_0 , S_1 , S_2 , S_4 , S_3 and \tilde{S}_3 . The remaining four solutions are those of an operator of order eight.

One of these solutions is highly remarkable: it is actually the two-particle contribution $\chi^{(2)}$. A similar situation also occured for the differential equation of the three particle contribution $\chi^{(3)}$, which actually had $\chi^{(1)}$ as solution.

In general, for all the $\tilde{\chi}^{(n)}$'s, it is tempting to expect $\tilde{\chi}^{(1)}$ to be a solution of the differential equation satisfied by the $\tilde{\chi}^{(2n+1)}$'s and $\tilde{\chi}^{(2)}$ to be a solution of the differential equation satisfied by the $\tilde{\chi}^{(2n)}$'s.

Beyond, one can contemplate much stronger conjectures corresponding to further "embedding", namely $\tilde{\chi}^{(m)}$ being solution of the differential equation of $\tilde{\chi}^{(n)}$, when n is a multiple of m, n and m having the same parity.

A confirmation of these various embeddings and conjectures is crucial because it corresponds to a *global structure* of the $\tilde{\chi}^{(n)}$'s and thus of χ itself. We will investigate this feature in a future publication.

Acknowledgments We thank Jacques-Arthur Weil for valuable comments on factorizations of differential operators. We would like to thank A. J. Guttmann and J.

Dethridge for extensive calculations on our series. We would like to thank particularly J. Dethridge for rewriting our formal programs into a masterpiece C++ program that confirmed our series and our final ODE result up to order 250. We thank the medicis formal calculation platform, at the stix laboratory of the Ecole Polytechnique, for enabling us to perform some formal calculations on their 8 Giga-memory machine. (S. B) and (S. H) acknowledge partial support from PNR3-Algeria. (N. Z.) is thankful to T. Hamidouche and H. Mazrou for letting him use their computer facilities to generate the first 130 terms of the series.

7. Appendix A

In this Appendix, we show that, with $H^{(4)}$ taken as in (4), i.e., not the alternative simplified form we use in this paper, our expansion method applied to $\tilde{\chi}^{(4)}$ produces five summations on products of four hypergeometric functions. Note that this method is applicable to any $\chi^{(n)}$.

Let us write the product $R^{(4)} \cdot H^{(4)}$ as

$$R^{(4)} \cdot H^{(4)} = T^{(4)} \cdot A^{(4)} \tag{59}$$

with:

$$T^{(4)} = R^{(4)} \cdot \prod_{i < j} \frac{\tilde{x}_i \tilde{x}_j}{(1 - \tilde{x}_i \tilde{x}_j)^2}$$
(60)

$$A^{(4)} = \frac{1}{4!} \cdot \prod_{i < j} (Z_i - Z_j)^2.$$
 (61)

By standard expansion of the quantity $T^{(4)}$, defined by (60), in the variables \tilde{x}_i , one obtains

$$T^{(4)} = \sum_{p=0}^{\infty} \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \sum_{i_3=0}^{\infty} \sum_{i_4=0}^{\infty} \sum_{i_5=0}^{\infty} \sum_{i_6=0}^{\infty} (1 + \theta(p-1)) \cdot \prod_{k=1}^{6} (i_k + 1) \cdot \tilde{x}_1^{p+i_1+i_2+i_3+3} \tilde{x}_2^{p+i_1+i_4+i_5+3} \tilde{x}_3^{p+i_2+i_4+i_6+3} \tilde{x}_4^{p+i_3+i_5+i_6+3}$$
(62)

where $\theta(x)$ is the step function given in (18). Defining

$$n_1 = p + i_1 + i_2 + i_3,$$
 $n_2 = p + i_1 + i_4 + i_5$
 $n_3 = p + i_2 + i_4 + i_6,$ $n_4 = p + i_3 + i_5 + i_6,$ (63)

one can solve (63) in the indices (i_3, i_4, i_5, i_6) , to obtain:

$$i_3 = -(p + i_1 + i_2) + n_1 (64)$$

$$i_4 = -(p + i_1 + i_2) + \frac{1}{2}(n_1 + n_2 + n_3 - n_4)$$
(65)

$$i_5 = i_2 - \frac{1}{2} \left(n_1 - n_2 + n_3 - n_4 \right) \tag{66}$$

$$i_6 = i_1 - \frac{1}{2} \left(n_1 + n_2 - n_3 - n_4 \right) \tag{67}$$

All these indices should be *integers*, inducing constraints on the n_i 's and limitations on the summations of the remaining indices (i.e., p, i_1 , i_2). With this change of summation indices, $T^{(4)}$ becomes

$$T^{(4)} = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \sum_{n_4=0}^{\infty} C(n_1, n_2, n_3, n_4) \cdot \tilde{x}_1^{n_1+3} \tilde{x}_2^{n_2+3} \tilde{x}_3^{n_3+3} \tilde{x}_4^{n_4+3}$$

$$(68)$$

and, since $T^{(4)}$ is completely symmetric in the variables \tilde{x}_i , the coefficient $C(n_1, n_2, n_3, n_4)$ is completely symmetric in the n_i indices and is given by summation on the remaining indices p, i_1 and i_2 :

$$C(n_1, n_2, n_3, n_4) = \sum_{p=0}^{\infty} \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} (1 + \theta(p-1)) \cdot \prod_{k=1}^{6} (i_k + 1) \times \theta(i_3) \cdot \theta(i_4) \cdot \theta(i_5) \cdot \theta(i_6) \cdot \sigma(n_1 + n_2 + n_3 + n_4) \times \theta\left(\frac{1}{2}(n_1 + n_2 + n_3 - n_4)\right)$$
(69)

where theta functions (18) of indices (i_3, i_4, i_5, i_6) take place to keep track of the fact that these indices should be positive integers. The symbol $\sigma(n)$ defined as:

$$\sigma(n) = \frac{1}{2} \left(1 + (-1)^n \right) \tag{70}$$

comes from the fact that i_3 , i_4 , i_5 , i_6 are integers, due to the right most terms at right-hand-side of (65-67), the n_i 's should verify $n_1 + n_2 + n_3 + n_4 =$ even integer. The argument in the last theta in (69) comes from (65).

Let us define the index q and its upper limit of summation (see (64,65)):

$$q = p + i_1 + i_2 (71)$$

$$q_0 = \min(n_1, \frac{n_1 + n_2 + n_3 - n_4}{2}) \tag{72}$$

Furthermore, the coefficient $C(n_1, n_2, n_3, n_4)$ being symmetric in all its arguments, it is sufficient to compute it in the case where:

$$n_1 \le n_2 \le n_3 \le n_4 \tag{73}$$

The coefficient $C(n_1, n_2, n_3, n_4)$ now becomes:

$$C(n_1, n_2, n_3, n_4) = \theta(\frac{n_1 + n_2 + n_3 - n_4}{2}) \cdot \sigma(n_1 + n_2 + n_3 + n_4)$$

$$\sum_{q=0}^{q_0} \sum_{p=0}^{q} \left((1 + \theta(p-1)) \cdot \sum_{i_1=0}^{q-p} \left(\prod_{k=1}^{6} (i_k + 1) \right) \right)$$
(74)

where the indices i_3, i_4, i_5, i_6 are given by (64-67) and $i_2 = q - p - i_1$. The three summations can be performed and one obtains

$$C(n_1, n_2, n_3, n_4) = \theta(\frac{n_1 + n_2 + n_3 - n_4}{2}) \cdot \sigma(n_1 + n_2 + n_3 + n_4) \cdot \left(\frac{q_0 + 4}{4}\right) Q(n_1, n_2, n_3, n_4, q_0)$$
(75)

with the polynomial $Q(n_1, n_2, n_3, n_4, q_0)$ given by

$$Q(n_1, n_2, n_3, n_4, q_0) = \frac{1}{8} Q_0(n_1, n_2, n_3, n_4) + \frac{q_0}{2520} Q_1(n_1, n_2, n_3, n_4) + \frac{q_0^2}{7560} Q_2(n_1, n_2, n_3, n_4) + \frac{q_0^3}{1890} Q_3(n_1, n_2, n_3, n_4) + \frac{q_0^4}{540} (92 - 81n_1 - 9n_2 - 9n_3 + 63n_4) + \frac{4q_0^5}{135}$$
(76)

with:

$$Q_0(n_1, n_2, n_3, n_4) = (1 + n_1)(n_1 + n_2 - n_3 - n_4 - 2)$$

$$(n_1 - n_2 + n_3 - n_4 - 2)(2 + n_1 + n_2 + n_3 - n_4)$$
(77)

$$Q_{1}(n_{1}, n_{2}, n_{3}, n_{4}) = -2216 + 2472n_{1} + 1194n_{1}^{2} - 1281n_{1}^{3} + 126n_{1}^{4}$$

$$-48n_{2} + 1392n_{1}n_{2} - 987n_{1}^{2}n_{2} + 126n_{1}^{3}n_{2} + 774n_{2}^{2} + 441n_{1}n_{2}^{2}$$

$$-126n_{1}^{2}n_{2}^{2} + 147n_{2}^{3} - 126n_{1}n_{2}^{3} - 48n_{3} + 1392n_{1}n_{3} - 987n_{1}^{2}n_{3}$$

$$+126n_{1}^{3}n_{3} - 1548n_{2}n_{3} - 882n_{1}n_{2}n_{3} + 252n_{1}^{2}n_{2}n_{3} - 147n_{2}^{2}n_{3}$$

$$+126n_{1}n_{2}^{2}n_{3} + 774n_{3}^{2} + 441n_{1}n_{3}^{2} - 126n_{1}^{2}n_{3}^{2} - 147n_{2}n_{3}^{2}$$

$$+126n_{1}n_{2}n_{3}^{2} + 147n_{3}^{3} - 126n_{1}n_{3}^{3} - 2568n_{4} - 708n_{1}n_{4}$$

$$+2709n_{1}^{2}n_{4} - 378n_{1}^{3}n_{4} - 288n_{2}n_{4} + 1134n_{1}n_{2}n_{4} - 252n_{1}^{2}n_{2}n_{4}$$

$$-147n_{2}^{2}n_{4} + 126n_{1}n_{2}^{2}n_{4} - 288n_{3}n_{4} + 1134n_{1}n_{3}n_{4} - 147n_{3}^{2}n_{4}$$

$$+294n_{2}n_{3}n_{4} - 252n_{1}n_{2}n_{3}n_{4} - 252n_{1}^{2}n_{3}n_{4} + 126n_{1}n_{2}n_{4}^{2}$$

$$+126n_{1}n_{3}^{2}n_{4} - 486n_{4}^{2} - 1575n_{1}n_{4}^{2} + 378n_{1}^{2}n_{4}^{2} - 147n_{2}n_{4}^{2}$$

$$-147n_{3}n_{4}^{2} + 126n_{1}n_{3}n_{4}^{2} + 147n_{4}^{3} - 126n_{1}n_{4}^{3}$$
(78)
$$Q_{2}(n_{1}, n_{2}, n_{3}, n_{4}) = -2992 - 4446n_{1} + 6534n_{1}^{2} - 1449n_{1}^{3} - 1926n_{2}$$

$$-819n_{1}^{2}n_{2} - 270n_{2}^{2} + 945n_{1}n_{2}^{2} + 315n_{2}^{3} - 1926n_{3} + 2808n_{1}n_{3}$$

$$-819n_{1}^{2}n_{3} + 540n_{2}n_{3} - 1890n_{1}n_{2}n_{3} - 315n_{2}^{2}n_{3} + 2808n_{1}n_{2}$$

$$-270n_{3}^{2} + 945n_{1}n_{3}^{2} - 315n_{2}n_{3}^{2} + 315n_{3}^{3} + 594n_{4} - 315n_{3}n_{4}^{2}$$

$$-8532n_{1}n_{4} + 3213n_{1}^{2}n_{4} - 1728n_{2}n_{4} + 1134n_{1}n_{2}n_{4} + 315n_{3}^{3}$$

$$-315n_{2}^{2}n_{4} - 1728n_{3}n_{4} + 1134n_{1}n_{3}n_{4} + 630n_{2}n_{3}n_{4}$$

$$-315n_{3}^{2}n_{4} + 1998n_{4}^{2} - 2079n_{1}n_{4}^{2} - 315n_{2}n_{4}^{2}$$
(79)
$$Q_{3}(n_{1}, n_{2}, n_{3}, n_{4}) = 520 - 1215n_{1} + 495n_{1}^{2} - 207n_{2} + 144n_{1}n_{2}$$

$$+ 144n_{1}n_{3} + 270n_{2}n_{3} - 135n_{3}^{2} + 801n_{4} - 738n_{1}n_{4}$$

$$- 108n_{3}n_{4} + 243n_{4}^{2} - 135n_{2}^{2} - 207n_{3} - 108n_{2}n_{4}$$
(80)

From the definition of $\tilde{\chi}^{(4)}$ in (2), and from (59,68), one gets:

$$\tilde{\chi}^{(4)} = 16 \, w^{16} \cdot \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \sum_{n_4=0}^{\infty} w^{n_1+n_2+n_3+n_4} \times$$

$$C(n_1, n_2, n_3, n_4) \cdot M(n_1, n_2, n_3, n_4)$$
(81)

The expansion method on the variables x_i introduces some summations free of any angular dependence. These summations appear in the coefficient $C(n_1, n_2, n_3, n_4)$ which is a kind of "geometrical factor" that appears for all the $\chi^{(n)}$'s with $n \geq 4$.

At this step the integrations have not yet been performed. They are contained in:

$$M(n_1, n_2, n_3, n_4) = \int_0^{2\pi} \frac{d\phi_1}{2\pi} \int_0^{2\pi} \frac{d\phi_2}{2\pi} \int_0^{2\pi} \frac{d\phi_3}{2\pi} \left(\prod_{i=1}^4 y_i \cdot x_i^{n_i + 3} \right) \cdot A^{(4)}$$
(82)

Due to its structure, it is obvious that $M(n_1, n_2, n_3, n_4)$ is completely symmetric in the indices n_i . We write $M(n_1, n_2, n_3, n_4)$ as an integral over the four angles $(\phi_1, \phi_2, \phi_3, \phi_4)$, by introducing Dirac delta function $\delta(\phi_1 + \phi_2 + \phi_3 + \phi_4)$ as a Fourier expansion that reads:

$$2\pi\delta(\phi_1 + \phi_2 + \phi_3 + \phi_4) = \sum_{k=-\infty}^{\infty} (Z_1 Z_2 Z_3 Z_4)^k$$
(83)

 $M(n_1, n_2, n_3, n_4)$ thus becomes

$$M(n_1, n_2, n_3, n_4) = \sum_{k=-\infty}^{\infty} M(n_1, n_2, n_3, n_4; k)$$
(84)

where:

$$M(n_1, n_2, n_3, n_4; k) = \int_0^{2\pi} \frac{d\phi_1}{2\pi} \int_0^{2\pi} \frac{d\phi_2}{2\pi} \int_0^{2\pi} \frac{d\phi_3}{2\pi} \int_0^{2\pi} \frac{d\phi_4}{2\pi} \left(\prod_{i=1}^4 y_i \cdot x_i^{n_i + 3} \right) A^{(4)} \cdot (Z_1 Z_2 Z_3 Z_4)^k.$$
(85)

The Fourier expansion (9) implies the following integration rule

$$\int_0^{2\pi} \frac{d\phi}{2\pi} (yx^m) \cdot Z^j = A(m, j). \tag{86}$$

The calculation of (85) thus becomes straightforward and does not induce any summation. The quantity $M(n_1, n_2, n_3, n_4; k)$ comes out as a huge expression with 201 terms, each of them being a product of four hypergeometric functions. Due to the definition (84) where k runs from $-\infty$ to ∞ , and to the symmetry of (84) under the permutation of the n_i 's, these 201 terms reduce to only 16 terms and $M(n_1, n_2, n_3, n_4; k)$ reads:

$$M(n_{1}, n_{2}, n_{3}, n_{4}; k) \equiv A(n_{1}, k)A(n_{2}, k)A(n_{3}, k)A(n_{4}, k)$$

$$-A(n_{1}, k)A(n_{2}, k)A(n_{3}, k)A(n_{4}, k+4)$$

$$A(n_{1}, k)A(n_{2}, k)A(n_{3}, k+2)A(n_{4}, k+2)$$

$$+2A(n_{1}, k)A(n_{2}, k)A(n_{3}, k+1)A(n_{4}, k+3)$$

$$-3A(n_{1}, k)A(n_{2}, k+1)A(n_{3}, k+1)A(n_{4}, k+2)$$

$$-A(n_{1}, k)A(n_{2}, k)A(n_{3}, k+3)A(n_{4}, k+5)$$

$$+A(n_{1}, k)A(n_{2}, k)A(n_{3}, k+4)A(n_{4}, k+4)$$

$$-2A(n_{1}, k)A(n_{2}, k+1)A(n_{3}, k+3)A(n_{4}, k+4)$$

$$+2A(n_{1}, k)A(n_{2}, k+1)A(n_{3}, k+2)A(n_{4}, k+5)$$

$$+2A(n_{1}, k)A(n_{2}, k+2)A(n_{3}, k+3)A(n_{4}, k+3)$$

$$-A(n_{1}, k)A(n_{2}, k+2)A(n_{3}, k+2)A(n_{4}, k+4)$$

$$+A(n_{1}, k)A(n_{2}, k+2)A(n_{3}, k+4)A(n_{4}, k+6)$$

$$-A(n_{1}, k)A(n_{2}, k+2)A(n_{3}, k+5)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+3)A(n_{3}, k+3)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+3)A(n_{3}, k+4)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+3)A(n_{3}, k+4)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+3)A(n_{3}, k+4)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+4)A(n_{3}, k+4)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+4)A(n_{3}, k+4)A(n_{4}, k+5)$$

$$-A(n_{1}, k)A(n_{2}, k+4)A(n_{3}, k+4)A(n_{4}, k+4)$$

$$(87)$$

which is equal to (85) for summation purposes.

Finally, collecting (81, 84), $\tilde{\chi}^{(4)}$ can be written as:

$$\tilde{\chi}^{(4)} = 16 \, w^{16} \cdot \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \sum_{n_3=0}^{\infty} \sum_{n_4=0}^{\infty} w^{n_1+n_2+n_3+n_4} \times C(n_1, n_2, n_3, n_4) \cdot M(n_1, n_2, n_3, n_4)$$
(88)

8. Appendix B

In this Appendix, we give the explicit derivation of the expression (14) given in Section 2.

Considering $H^{(4)}$ as in (13), we define the following relations

$$\prod_{i < j} \frac{\tilde{x}_i - \tilde{x}_j}{1 - \tilde{x}_i \tilde{x}_j} = P_{12-34}^{(x)} - (1 \leftrightarrow 3) - (2 \leftrightarrow 3)$$
(89)

$$\prod_{i < j} \frac{Z_i - Z_j}{1 - Z_i Z_j} = P_{12-34}^{(Z)} - (1 \leftrightarrow 3) - (2 \leftrightarrow 3)$$
(90)

where $P_{12-34}^{(x)}$ is given in (15). Using the constraint (6), $P_{12-34}^{(Z)}$ reads:

$$P_{12-34}^{(Z)} = \tilde{A}^{(4)} + (Z_2 - Z_1)(Z_3 - Z_4)(1 + Z_1 Z_2 + Z_3 Z_4)$$
(91)

with
$$\widehat{A}^{(4)} = (Z_2 - Z_1)(Z_3 - Z_4) \cdot \frac{Z_1 Z_2}{(1 - Z_1 Z_2)^2}$$
 (92)

Note that, while $H^{(4)}$ and $R^{(4)}$ are completely symmetric under the permutation of the angles ϕ_i , the right hand side of (89,90) are anti-symmetric under the same transformation.

From (13, 89, 90) and using the symmetry of the rest of the integrand (namely $R^{(4)}$ times the product over y_i variables), we can write $H^{(4)}$ as:

$$H^{(4)} \equiv \frac{1}{8} P_{12-34}^{(Z)} \cdot \left(P_{12-34}^{(x)} - (1 \leftrightarrow 3) - (2 \leftrightarrow 3) \right) \tag{93}$$

We obtain $H^{(4)}$, given in (14), from (93) with the first term at the right hand side of (91). Again using the property of complete symmetry of the integrand, that part of (93) with the second term at the right hand side of (91) can be written as:

$$-\frac{1}{8}P_{12-34}^{(x)} \cdot \left((Z_1 - Z_2)(Z_3 - Z_4)(1 + Z_1Z_2 + Z_3Z_4) - (1 \leftrightarrow 3) - (2 \leftrightarrow 3) \right)$$

$$(94)$$

and is identically null.

9. Appendix C

In this Appendix, we give the explicit integration rules corresponding to the form (16) given in the main text. For this purpose, let us consider

$$J(n_1, n_2, n_3, n_4) = \int_0^{2\pi} \frac{d\phi_1}{2\pi} \int_0^{2\pi} \frac{d\phi_2}{2\pi} \int_0^{2\pi} \frac{d\phi_3}{2\pi} \left(\prod_{i=1}^4 y_i \, x_i^{n_i} \right) \cdot \widehat{A}^{(4)}$$
(95)

where $\widehat{A}^{(4)}$ is given in (92). Expression $J(n_1, n_2, n_3, n_4)$ has the following properties:

$$J(n, n, n_3, n_4) = J(n_1, n_2, n, n) = 0$$

$$J(n_1, n_2, n_3, n_4) = -J(n_2, n_1, n_3, n_4) = J(n_3, n_4, n_1, n_2)$$
(96)

where the last identity comes from the fact that

$$\frac{Z_1 Z_2}{(1 - Z_1 Z_2)^2} = \frac{Z_3 Z_4}{(1 - Z_3 Z_4)^2}. (97)$$

Changing the integration variables from (ϕ_1, ϕ_2, ϕ_3) to $(\phi = \phi_1 + \phi_2, \phi_2, \phi_3)$, $\widehat{A}^{(4)}$, defined in (92), becomes:

$$\widehat{A}^{(4)} = \frac{1}{(1-Z)^2} \cdot \left[\left(\frac{1}{Z_2 Z_3} + Z_2 Z_3 \right) \cdot Z - \frac{Z_2}{Z_3} - \frac{Z_3}{Z_2} Z^2 \right]$$
(98)

with $Z = Z_1 Z_2$. Taking $\widehat{A}^{(4)}$ in the form (98), expanding $y_1 x_1^{n_1}$ and $y_4 x_4^{n_4}$, and using the Fourier expansion (9), the integration over ϕ_2 and ϕ_3 is straightforward \parallel . Some standard manipulations give:

$$J(n_1, n_2, n_3, n_4) = -\int_0^{2\pi} \frac{d\phi}{2\pi} \cdot \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} D(n_1, n_2; i) D(n_3, n_4; j) \cdot \frac{Z^{j-i+1}}{(1-Z)^2}$$
(99)

where:

$$D(n_1, n_2; k) = A(n_1, k+1) A(n_2, k) - A(n_1, k) A(n_2, k+1).$$
 (100)

From the definition (10) of A(n,k), one can easily show that

$$D(n_1, n_2; -k) = -D(n_1, n_2; k - 1)$$
(101)

With this property (101), and after some manipulations, $J(n_1, n_2, n_3, n_4)$ reads

$$J(n_1, n_2, n_3, n_4) = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} D(n_1, n_2; i) D(n_3, n_4; j) N(i, j)$$
 (102)

where

$$N(i,j) = \int_0^{2\pi} \frac{d\phi}{2\pi} Z^{-j-i} \frac{1 - Z^{2i+1}}{1 - Z} \frac{1 - Z^{2j+1}}{1 - Z}$$
(103)

which, by expansion in powers of Z and integration, simply reads:

$$N(i,j) = 2\min(i,j) + 1. \tag{104}$$

Finally, taking the form (102), together with (104), and using the definition (10) of A(n,k), we obtain:

$$J(n_1, n_2, n_3, n_4) = \frac{1}{2} \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} w^{2j+4k+2} (2k+1) (1+\theta (j-1)) \cdot \left(d(n_1, n_2; k) d(n_3, n_4; j+k) + d(n_1, n_2; j+k) d(n_3, n_4; k) \right)$$
(105)

where $d(n_1, n_2; k)$ is defined in (20).

|| see the integration rule (86).

10. Appendix D

In this Appendix, we give all the necessary details of the derivation of the result (17) given in the text. We consider $\tilde{\chi}^{(4)}$ in the form (2), with $H^{(4)}$ given in (14).

The first step is to consider the expansion of the product $R^{(4)} \cdot P_{12-34}^{(x)}$. Let us give the expansions, in the x_i 's variables, for the first two terms in the brackets in (15):

$$R^{(4)} \cdot \frac{\tilde{x}_1 \tilde{x}_2}{1 - \tilde{x}_1 \tilde{x}_2} \frac{\tilde{x}_3 \tilde{x}_4}{1 - \tilde{x}_3 \tilde{x}_4} = \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} (\tilde{x}_1 \tilde{x}_2)^m (\tilde{x}_3 \tilde{x}_4)^l \cdot \theta(m-1) \theta(l-1) (2 \min(m,l) - 1)$$
(106)

$$R^{(4)} \cdot \frac{(\tilde{x}_1 \tilde{x}_2)^2}{1 - \tilde{x}_1 \tilde{x}_2} = \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} (\tilde{x}_1 \tilde{x}_2)^m (\tilde{x}_3 \tilde{x}_4)^l \cdot \theta(m-l-2) \theta(m-2) (1 + \theta(l-1)).$$
(107)

 $R^{(4)} \cdot P_{12-34}^{(x)}$ can be expanded as

$$R^{(4)} \cdot P_{12-34}^{(x)} = (x_1 - x_2) \cdot (x_3 - x_4) \cdot \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} w^{2m+2l+2} \cdot (108)$$
$$(x_1 x_2)^m (x_3 x_4)^l \cdot c(m, l)$$

with:

$$c(m,l) = \theta(m-1) \theta(l-1) \cdot (2\min(m,l) - 1) + \theta(m-l-2) \theta(m-2) \cdot (1 + \theta(l-1)) + \theta(l-m-2) \theta(l-2) \cdot (1 + \theta(m-1)).$$
 (109)

Considering $\tilde{\chi}^{(4)}$ in the form (2), with $H^{(4)}$ given by (14), taking the expansion of $R^{(4)} \cdot P_{ij-kl}^{(x)}$ in the form (108), performing the angular integration using the definition (95), and collecting all terms, gives

$$\tilde{\chi}^{(4)} = 16 \, w^6 \cdot \sum_{m=0}^{\infty} \sum_{p=0}^{\infty} \frac{1}{2} w^{2m+2p} \, c(m,p) \, I(m,p)$$
 (110)

where:

$$I(m,p) = J(m,m+1,p,p+1) - J(m,p,m+1,p+1) - J(m,p+1,p,m+1).$$
(111)

Using the identities (96) for expression $J(n_1, n_2, n_3, n_4)$, one has:

$$I(p,m) = I(m,p)$$

 $I(m,m) = I(m,m+1) = 0.$ (112)

The symmetry of c(m, p), together with the relations (112), allow us, after some manipulations, to write $\tilde{\chi}^{(4)}$ as

$$\tilde{\chi}^{(4)} = 16 \, w^{10} \cdot \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} w^{4m+2n} \, (2m+1) \cdot I(m, n+m+2)$$
 (113)

where we have used the following identity c(m, n + m + 2) = 2m + 1.

From the previous equation, taking the definition (111) together with the expression (105), one obtains

$$\tilde{\chi}^{(4)} = 16w^{12} \cdot \sum_{m,k,n,j=0}^{\infty} w^{4m+4k+2n+2j} (2m+1) (2k+1) \times \frac{1}{2} \left(1 + \theta (j-1) \right) \cdot V(m,k,n,j-2)$$
(114)

where V(m, k, n, j) is defined in (19), with the constraints:

$$V(m,k,n,-2) = V(m,k,n,-1) = 0 (115)$$

Thus, $\tilde{\chi}^{(4)}$ reads :

$$\tilde{\chi}^{(4)} = 16 \, w^{16} \cdot \sum_{m,k,n,j=0}^{\infty} w^{4m+4k+2n+2j} \times (2m+1) \cdot (2k+1) \cdot V(m,k,n,j). \tag{116}$$

The last simplification comes from the following identity on V(m, k, n, j)

$$V(k, m, j, n) = V(m, k, n, j). (117)$$

which is obtained with the help of the symmetry of a(n, k), namely a(n, k) = a(k, n). Finally, using the general identity

$$\sum_{p_1=0}^{\infty} \sum_{p_2=0}^{\infty} s(p_1, p_2) = \sum_{p=0}^{\infty} \sum_{i=0}^{\infty} (1 + \theta(i-1)) \times \frac{1}{2} (s(p+i, p) + s(p, p+i))$$
(118)

and the identity (117), the expression (116) for $\tilde{\chi}^{(4)}$ can, after some manipulation, be written in the form (17) displayed in the text.

11. Appendix E

$$P_{10}(x) = 192598769664000 - 943722860380160 x \\ + 1154055263764480 x^2 + 223612469147648 x^3 \\ + 498965092419008 x^4 - 4709824359388336 x^5 \\ + 6098813329440179 x^6 - 2687506699337617 x^7 \\ - 752969324018818 x^8 + 1919011581320339 x^9 \\ - 1526656430056013 x^{10} + 660280621356073 x^{11} \\ - 134468923815612 x^{12} + 7980003107181 x^{13} \\ + 671991155376 x^{14} - 17122807680 x^{15} \\ - 5111390208 x^{16} + 43929600 x^{17}$$
 (119)

$$P_9(x) = 39290149011456000 - 297447362118287360 x$$

$$+ 785045342889902080 x^2 - 738039510467911680 x^3$$

$$+ 115430038068948224 x^4 - 1014554971757285568 x^5$$

$$+ 3530827584228091380 x^6 - 4191676869770939784 x^7$$

```
+2038314404276922993 x^{8} + 302242744021242129 x^{9}
-1228406406705470056 x^{10} + 1043856642401535101 x^{11}
-506114460611418219 x^{12} + 139123852417110463 x^{13}
-18756086615529738 x^{14} + 585977956065027 x^{15}
+80289868813296 x^{16} + 338076239232 x^{17}
-554882826240 x^{18} + 4524748800 x^{19} 
(120)
P_{8}(x) = -197221140135936000 + 2237384185085952000 x
-9353251554131968000 x^{2} + 17417661939469352960 x^{3}
```

 $-9353251554131968000 x^{2} + 17417661939469352960 x^{3} \\ -12764493479941255168 x^{4} + 5721656503410538176 x^{5} \\ -31513487740861948240 x^{6} + 74327124872294722259 x^{7} \\ -74793556916527926194 x^{8} + 31202576844037093699 x^{9} \\ +8393285642528898691 x^{10} - 21583664946799604164 x^{11} \\ +16831944005767359920 x^{12} - 7706219890490209273 x^{13} \\ +2036277808796847523 x^{14} - 267301496692115881 x^{15} \\ +8265260736796220 x^{16} + 1095242827496832 x^{17} \\ +4127794059264 x^{18} - 7320844222464 x^{19} \\ +58514227200 x^{20}$ (121)

 $P_{7}(x) = 764231918026752000 - 16971026878221516800 x \\ + 115553546628494786560 x^{2} - 357128015448284200960 x^{3} \\ + 527925844930277396480 x^{4} - 327328761204753779200 x^{5} \\ + 264654998297650108784 x^{6} - 1129192842855079731378 x^{7} \\ + 2154212025640162544967 x^{8} - 1923633718567453943246 x^{9} \\ + 708245350538757849333 x^{10} + 257335760183269925607 x^{11} \\ - 530877971609916541978 x^{12} + 387828430484350338476 x^{13} \\ - 169278351316199852707 x^{14} + 43213759197299807177 x^{15} \\ - 5540993509581072863 x^{16} + 171042063471920404 x^{17} \\ + 21596554335396864 x^{18} + 67975771296768 x^{19} \\ - 139524297965568 x^{20} + 1091386982400 x^{21}$ (122)

$$\begin{split} P_6(x) &= 2366653681631232000 + 15657511469396787200 \, x \\ &- 418375361108283228160 \, x^2 + 2413740972333717913600 \, x^3 \\ &- 6304389540169208659968 \, x^4 + 7885479774610742167552 \, x^5 \\ &- 4194762998349204963264 \, x^6 + 4787870532372577669360 \, x^7 \\ &- 18347979159424299125323 \, x^8 + 30953188609834801509491 \, x^9 \\ &- 25453291295484088365791 \, x^{10} + 8618062554065795648668 \, x^{11} \\ &+ 3501102053489313155231 \, x^{12} - 6497943932928673199052 \, x^{13} \end{split}$$

```
\begin{array}{l} +\ 4524991429929970477637\ x^{14} - 1899642804937798765560\ x^{15} \\ +\ 470903276066137118684\ x^{16} - 59187343449427131707\ x^{17} \\ +\ 1841781629480835082\ x^{18} + 217686233501214240\ x^{19} \\ +\ 505738155562752\ x^{20} - 1358525988175872\ x^{21} \\ +\ 10379070873600\ x^{22} \end{array} \tag{123}
```

 $P_5(x) = -4240254512922624000 + 62906863287743283200 \, x \\ - 176844924247909335040 \, x^2 - 926585736920055152640 \, x^3 \\ + 6926618061828909694976 \, x^4 - 17270267632607743877120 \, x^5 \\ + 19088006038191340885888 \, x^6 - 7801708756272263419344 \, x^7 \\ + 10929800840921163293912 \, x^8 - 45123693732451251946951 \, x^9 \\ + 72469347127122031920503 \, x^{10} - 57004940985415997264335 \, x^{11} \\ + 18768946563950160696028 \, x^{12} + 7071447265880540507399 \, x^{13} \\ - 12955437249853164400164 \, x^{14} + 8753288431922263351485 \, x^{15} \\ - 3567263867581256107712 \, x^{16} + 863585677517038259840 \, x^{17} \\ - 106841634606075117511 \, x^{18} + 3388641803935920834 \, x^{19} \\ + 366870695477680992 \, x^{20} + 460940531245824 \, x^{21} \\ - 2210964576190464 \, x^{22} + 16454358835200 \, x^{23} \tag{124}$

 $P_4(x) = 2366653681631232000 - 121147207913255731200 \, x \\ + 1204116891510820044800 \, x^2 - 4471702378853957632000 \, x^3 \\ + 3882247113210548060160 \, x^4 + 19050981728126452531200 \, x^5 \\ - 60173463653707794252800 \, x^6 + 61647224332324323946560 \, x^7 \\ - 8902468613107108100784 \, x^8 + 12826049043414461768619 \, x^9 \\ - 130168614999998943881548 x^{10} + 224001261613376678757254 x^{11} \\ - 177856326190823713976989 x^{12} + 61054390293590247070213 x^{13} \\ + 16702537875405165273835 \, x^{14} - 34504858377234313267137 \, x^{15} \\ + 23145136640956770024901 \, x^{16} - 9263503411249994637068 \, x^{17} \\ + 2206094527520924501173 \, x^{18} - 270104745702122623849 \, x^{19} \\ + 8842661536198397940 \, x^{20} + 853061197515225600 \, x^{21} \\ - 104681638038528 \, x^{22} - 4965596931735552 \, x^{23} \\ + 35863949721600 \, x^{24} \tag{125}$

 $P_3(x) = -11833268408156160000 + 296345375978029056000 x$ $-2144074027349363916800 x^2 + 6754783276113459937280 x^3$ $-8346802888116933918720 x^4 - 3858851854352328468480 x^5$ $+18632235498956809831680 x^6 - 5896832882414934369872 x^7$ $-20092581953702558202578 x^8 + 9039258514198720200789 x^9$

```
\begin{array}{l} +\,30407815233273202124059\,x^{10}-44070504829347073747333\,x^{11}\\ +\,24353838992694268159678\,x^{12}-2634376094447814881673\,x^{13}\\ -\,5786419996453628346078\,x^{14}+5219893433655665719185\,x^{15}\\ -\,2420429903132324343712\,x^{16}+639931684854726532004\,x^{17}\\ -\,85290635341359290409\,x^{18}+3288947218141266164\,x^{19}\\ +\,249618983188012416\,x^{20}-1896484503705600\,x^{21}\\ -\,1406458574438400\,x^{22}-9870981120000\,x^{23} \end{array} \tag{126}
```

```
P_{2}(x) = 36978963775488000000 - 561176784419998924800 \ x \\ + 3011344824723628359680 \ x^{2} - 7448695810925636485120 \ x^{3} \\ + 7522952469866279936000 \ x^{4} + 1571640457415428633600 \ x^{5} \\ - 6448286163781893470272 \ x^{6} - 5723224826762698171248 \ x^{7} \\ + 17523982043162827496339 \ x^{8} - 10871737924438446047496 \ x^{9} \\ - 1585720216853837551908 \ x^{10} + 4808127638071252147363 \ x^{11} \\ - 1996433111317397547653 \ x^{12} - 255305750980018902593 \ x^{13} \\ + 746533058733318361745 \ x^{14} - 451215245481551846087 \ x^{15} \\ + 139082951228390470494 \ x^{16} - 20614123445854835699 \ x^{17} \\ + 931298896466844519 \ x^{18} + 53593802936851536 \ x^{19} \\ - 845315534083200 \ x^{20} - 297344753664000 \ x^{21} \\ + 2006484480000 \ x^{22} \tag{127}
```

```
P_{1}(x) = -14878398029915750400 + 132065832050507120640 x - 489750894279960821760 x^{2} + 843938312965142528000 x^{3} - 436158709746751846400 x^{4} - 504050378774640931456 x^{5} + 362705915662042967696 x^{6} + 930077072222989626372 x^{7} - 1516628759565419342933 x^{8} + 919766159750634600991 x^{9} - 236614768621801865201 x^{10} - 4425339208940491244 x^{11} + 15751985529081115811 x^{12} + 356035582683101460 x^{13} - 3725421245175199951 x^{14} + 1704676309420947212 x^{15} - 304427228661108302 x^{16} + 16257997137256137 x^{17} + 635239438099728 x^{18} - 16870300732800 x^{19} - 3641326080000 x^{20} + 23063040000 x^{21}  (128)
```

12. Appendix F

Let us recall a few elementary facts on factorization of differential operators, without introducing any sophisticated \mathcal{D} -module theory. Two differential operators \mathcal{D}_1 and \mathcal{D}_2 of order N are said to be equivalent (see for instance equation (5) in [21]) if there exist two (intertwining) differential operators R and S of orders at most N-1, such

that:

$$\mathcal{D}_2 \cdot R = S \cdot \mathcal{D}_1. \tag{129}$$

The factorization of differential operators is not in general unique. There can be several factorizations, even an infinite number of factorizations, as can be seen in the simple exemple:

$$\frac{d^2}{dx^2} = \left(\frac{d}{dx} + \frac{a}{ax+b}\right) \cdot \left(\frac{d}{dx} - \frac{a}{ax+b}\right) \tag{130}$$

where the factorization (130) holds for every constant a and b. This can be seen as a consequence of the relation :

$$\left(\frac{d}{dx} - \frac{a}{ax+b}\right) \cdot \left(ax+b\right) = \left(ax+b\right) \cdot \frac{d}{dx} \tag{131}$$

which means that the multiplication by ax + b transforms the solutions of the differential operator d/dx into a solution of d/dx - a/(ax + b). Relation (131) is a simple exemple of equivalence (129) of two differential operators. The occurrence of this infinite set of factorizations can be seen to be a consequence of the fact that the various differential operators of (130), (131) have solutions the ratio of which can be rational functions. In general this is not the case and one will only have a finite number of factorizations, these various factorizations being basically the same up to some permutation of the differential operators, each operator in the factorization being changed ("by some transmutation" (129)) into an equivalent one during the permutation process.

This is typically the situation we encounter with the operator \mathcal{L}_{10} associated with the ordinary differential equation of order ten satisfied by $\tilde{\chi}^{(4)}$.

This order ten operator \mathcal{L}_{10} has 36 factorizations with two operators of order four $(M_1 \text{ and } M_2)$, ten operators of order two (N_0, N_1, \dots, N_9) and 26 operators of order one $(L_0, L_1, \dots L_{25})$. All these operators have been normalized in such a way that the coefficient of the highest derivative is normalized to +1. The operators L_0 , L_1 , L_2 , N_0 and N_8 below are such that $L_0(\mathcal{S}_0) = 0$, $L_1(\mathcal{S}_1) = 0$, $L_2(\mathcal{S}_2) = 0$, $N_0(\mathcal{S}_3) = 0$ and $N_8(\mathcal{S}_1^*) = 0$. These 36 factorizations* of \mathcal{L}_{10} read

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot L_{25} \cdot L_{12} \cdot L_3 \cdot L_0$$

$$= M_1 \cdot N_9 \cdot L_{25} \cdot L_{12} \cdot L_3 \cdot L_0$$

$$= M_1 \cdot L_{24} \cdot N_4 \cdot L_{12} \cdot L_3 \cdot L_0$$

$$= M_1 \cdot L_{24} \cdot L_{13} \cdot N_6 \cdot L_3 \cdot L_0$$

$$= M_1 \cdot L_{24} \cdot L_{13} \cdot L_{17} \cdot N_3 \cdot L_0$$

$$= M_1 \cdot L_{24} \cdot L_{13} \cdot L_{17} \cdot L_{11} \cdot N_0$$
(132)

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot L_{25} \cdot L_{14} \cdot L_4 \cdot L_0
= M_1 \cdot N_9 \cdot L_{25} \cdot L_{14} \cdot L_4 \cdot L_0
= M_1 \cdot L_{24} \cdot N_4 \cdot L_{14} \cdot L_4 \cdot L_0
= M_1 \cdot L_{24} \cdot L_{15} \cdot N_7 \cdot L_4 \cdot L_0
= M_1 \cdot L_{24} \cdot L_{15} \cdot L_{16} \cdot N_3 \cdot L_0
= M_1 \cdot L_{24} \cdot L_{15} \cdot L_{16} \cdot L_{11} \cdot N_0$$
(133)

^{*} The DFactor command [27] of DEtools can give the second factorization in (132), namely $\mathcal{L}_{10} = M_1 \cdot N_9 \cdot L_{25} \cdot L_{12} \cdot L_3 \cdot L_0$.

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot L_{25} \cdot L_{18} \cdot L_5 \cdot L_1$$

$$= M_1 \cdot N_9 \cdot L_{25} \cdot L_{18} \cdot L_5 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot N_4 \cdot L_{18} \cdot L_5 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot L_{19} \cdot N_5 \cdot L_5 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot L_{19} \cdot L_{23} \cdot N_1 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot L_{19} \cdot L_{23} \cdot L_{10} \cdot N_0$$
(134)

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot L_{25} \cdot L_{14} \cdot L_6 \cdot L_1$$

$$= M_1 \cdot N_9 \cdot L_{25} \cdot L_{14} \cdot L_6 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot N_4 \cdot L_{14} \cdot L_6 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot L_{15} \cdot N_7 \cdot L_6 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot L_{15} \cdot L_{20} \cdot N_1 \cdot L_1$$

$$= M_1 \cdot L_{24} \cdot L_{15} \cdot L_{20} \cdot L_{10} \cdot N_0$$
(135)

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot L_{25} \cdot L_{18} \cdot L_7 \cdot L_2$$

$$= M_1 \cdot N_9 \cdot L_{25} \cdot L_{18} \cdot L_7 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot N_4 \cdot L_{18} \cdot L_7 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot L_{19} \cdot N_5 \cdot L_7 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot L_{19} \cdot L_{21} \cdot N_2 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot L_{19} \cdot L_{21} \cdot L_9 \cdot N_0$$
(136)

$$\mathcal{L}_{10} = N_8 \cdot M_2 \cdot L_{25} \cdot L_{12} \cdot L_8 \cdot L_2$$

$$= M_1 \cdot N_9 \cdot L_{25} \cdot L_{12} \cdot L_8 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot N_4 \cdot L_{12} \cdot L_8 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot L_{13} \cdot N_6 \cdot L_8 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot L_{13} \cdot L_{22} \cdot N_2 \cdot L_2$$

$$= M_1 \cdot L_{24} \cdot L_{13} \cdot L_{22} \cdot L_9 \cdot N_0$$
(137)

All these 36 factorizations can be seen as the consequence of "transmutations" like (129), where one differential operator becomes an equivalent one, thus yielding some elementary permutation of the operator. For instance, the two differential operators of order four, namely M_1 and M_2 , are equivalent, their "intertwiners" being two order two differential operators N_8 and N_9 :

$$N_8 \cdot M_2 = M_1 \cdot N_9 \tag{138}$$

Actually, one sees clearly that the two first factorizations in (132) are just deduced from one another by (138).

Nine order-two differential operators N_0, \dots, N_8 can be seen to be related by various equivalence relations requiring the introduction of 26 differential operators L_i 's of order one (the intertwiners):

$$N_{1} \cdot L_{1} = L_{10} \cdot N_{0}, \qquad N_{2} \cdot L_{2} = L_{9} \cdot N_{0}$$

$$N_{3} \cdot L_{0} = L_{11} \cdot N_{0}, \qquad N_{4} \cdot L_{12} = L_{13} \cdot N_{6}$$

$$N_{4} \cdot L_{14} = L_{15} \cdot N_{7}, \qquad N_{4} \cdot L_{18} = L_{19} \cdot N_{5}$$

$$N_{5} \cdot L_{5} = L_{23} \cdot N_{1}, \qquad N_{5} \cdot L_{7} = L_{21} \cdot N_{2}$$

$$N_{6} \cdot L_{3} = L_{17} \cdot N_{3}, \qquad N_{6} \cdot L_{8} = L_{22} \cdot N_{2}$$

$$N_{7} \cdot L_{4} = L_{16} \cdot N_{3}, \qquad N_{7} \cdot L_{6} = L_{20} \cdot N_{1}$$

$$N_{9} \cdot L_{25} = L_{24} \cdot N_{4}.$$

$$(139)$$

The 26 order-one differential operators L_i 's can also be seen to be related by various equivalence relations. The L_i 's being order one differential operators, their intertwiners (R and S in (129)) should be order zero differential operators, that is some function $f_{i,j}$. As a consequence of the normalization to +1 of the highest order derivative (here d/dx), the two intertwiners R and S in (129) are identical and one thus gets relations like:

$$L_i \cdot f_{i,j} = f_{i,j} \cdot L_j \tag{140}$$

Furthermore one also verifies the following relations between the order one operators L_i 's :

$$L_{3} \cdot L_{0} = L_{8} \cdot L_{2}, \qquad L_{4} \cdot L_{0} = L_{6} \cdot L_{1}$$

$$L_{5} \cdot L_{1} = L_{7} \cdot L_{2}$$

$$L_{12} \cdot L_{3} = L_{14} \cdot L_{4}, \qquad L_{13} \cdot L_{17} = L_{15} \cdot L_{16}$$

$$L_{18} \cdot L_{5} = L_{14} \cdot L_{6}, \qquad L_{19} \cdot L_{23} = L_{15} \cdot L_{20}$$

$$L_{18} \cdot L_{7} = L_{12} \cdot L_{8}, \qquad L_{19} \cdot L_{21} = L_{13} \cdot L_{22}$$

$$L_{17} \cdot L_{11} = L_{22} \cdot L_{9}, \qquad L_{16} \cdot L_{11} = L_{20} \cdot L_{10}$$

$$L_{23} \cdot L_{10} = L_{21} \cdot L_{9}$$

$$(141)$$

It is straightforward to see that a relation like $L_i \cdot L_j = L_k \cdot L_m$ yields relations like:

$$L_{i} \cdot (L_{j} - L_{m}) = (L_{k} - L_{i}) \cdot L_{m},$$

$$(L_{i} - L_{k}) \cdot L_{j} = L_{k} \cdot (L_{m} - L_{j}),$$

$$f_{i,m} = L_{j} - L_{m} = L_{k} - L_{i}, \qquad f_{k,j} = L_{m} - L_{j} = L_{i} - L_{k}$$

The number of intertwiners $f_{i,j}$ between the operators of order one in (140), is at first sight 325, but due to the above identities only 15 are independent.

- [1] T.T. Wu, B.M. McCoy, C.A. Tracy and E. Barouch, 1976 Phys. Rev. B 13, 316-374
- [2] C. R. Nappi, 1978 Nuovo Cim. A 44 392
- [3] J. Palmer, C. Tracy, 1981 Adv. Appl. Math. 2 329
- [4] K. Yamada, 1984 Prog. Theor. Phys. 71 1416
- [5] K. Yamada, 1985 Phys. Lett. A 112 456-458
- [6] B. Nickel, 1999 J. Phys. A: Math. Gen. 32 3889-3906
- [7] B. Nickel, 2000 J. Phys. A: Math. Gen. **33** 1693-1711
- [8] W.P. Orrick, B.G. Nickel, A.J. Guttmann, J.H.H. Perk, 2001 J. Stat. Phys. 102 795-841
- [9] W.P. Orrick, B.G. Nickel, A.J. Guttmann, J.H.H. Perk, 2001 Phys. Rev. Lett. 86 4120-4123
- [10] B.M. McCoy, T.T. Wu, 1980 Phys. Rev. Lett. 45 675-678
- [11] J.H.H. Perk, 1980 Phys. Lett. A79 3-5
- [12] M. Jimbo, T. Miwa, 1980 Proc. Japan Acad. A **56** 405; 1981 Proc. Japan Acad. A **57** 347

- [13] B.M. McCoy and T.T. Wu, Nucl. Phys. B180 (1981) 89
- [14] D. Hansel and J-M. Maillard, 1988 J. Phys. A: Math. Gen. 21 213-225
- [15] D. Hansel, J.M. Maillard, J. Oitmaa, M.J. Vergakis, J. Stat. Phys. 1987 48 69-80
- [16] A.J. Guttman, I.G. Enting, 1996 Phys. Rev. Lett. 76 344-347
- [17] N. Zenine, S. Boukraa, S. Hassani, J.M. Maillard, 2004 J. Phys. A: Math. Gen. 37 9651-9668 and arXiv:math-ph/0407060
- [18] N. Zenine, S. Boukraa, S. Hassani, J.M. Maillard, Square lattice Ising model susceptibility: Series expansion method and differential equation for $\chi^{(3)}$, 2005 to be published in J. Phys. A: Math. Gen. and arXiv:hep-ph/0411051
- [19] J. H. Davenport, The Difficultie od Definite Integration, http://www-calfor.lip6.fr/rr/Calculemus03/davenport.pdf
- [20] M. van der Put and M. F. Singer, Galois theory of difference equations, Springer-Verlag, Berlin, (1997), Lecture Notes in Mathematics. http://www4.ncsu.edu/singer/papers/dbook.ps
- [21] P. H. Berman and M. F. Singer, Calculating the Galois Group of $L_1(L_2(y)) = 0$, L_1 , L_2 Completly Reducible Operators., (1998) http://www4.ncsu.edu/singer/papers/12Inhom.ps
- [22] M. Van Hoeij, Formal solutions and factorization of differential operators with power series coefficients., Journal of Symbolic Computation, vol. 24, no1, July (1997), 1-30.
- [23] J-A. Weil, Absolute Factorization of Differential Operators, http://pauillac.inria.fr/algo/seminars/sem96-97/weil.html
- [24] S.A. Abramov, J.J. Carette, K.O. Geddes, and H.Q. Le, Telescoping in the context of symbolic summation in Maple. Journal of Symbolic Computation, 38 (4), (2004) 1303-1326.
- [25] M. Giesbrecht, Y. Zhang, Factoring and Decomposing Ore Polynomials over Fq(t). ACM International Symposium on Symbolic and Algebraic Computation (ISSAC), (2003) 127-134.
- [26] M. van Hoeij, help page for diffop, $http://www.math.fsu.edu/hoeij/daisy/help_pages$
- [27] DEtools library for differential equations http://www.math.upenn.edu/ugrad/calc/manual/deplot.pdf
- [28] H.K. Ince, Ordinary differential equations, (Longmans, London, 1927; also Dover Pubs., NY,
- [29] A. R. Forsyth, Theory of Differential Equations, 6 vols. (New York: Dover, 1959)
- [30] Mgfun's project : see http://algo.inria.fr/chyzak/mgfun.html ; gfun generating functions package see gfun in : http://algo.inria.fr/libraries
- [31] Michael F. Singer, Testing reducibility of linear differential operators: A group theoretic perspective., Applicable Algebra in Engineering, Communication and Computing, vol. 7, no2, (1996), 77-104.