

Vanishing Neel Ordering of $SU(n)$ Heisenberg Model in Three Dimensions

Noboru Fukushima

Department of Physics, University of the Saarland, 66041 Saarbrücken, Germany

The $SU(n)$ Heisenberg model represented by exchange operators is studied by means of high-temperature series expansion in three dimensions, where n is an arbitrary positive integer. The spin-spin correlation function and its Fourier transform $S(q)$ is derived up to $O[(J)^{10}]$ with J being the nearest-neighbor antiferromagnetic exchange in units of temperature. The temperature dependence of $S(q)$ and next-nearest-neighbor spin-spin correlation in the large n cases show that dominant correlation deviates from $q = (\pi, \pi, \pi)$ at low temperature, which is qualitatively similar to that of this model in one dimension. The Neel temperature of $SU(2)$ case is precisely estimated by analyzing the divergence of $S(\pi, \pi, \pi)$. Then, we generalize n of $SU(n)$ to a continuous variable and gradually increases from $n = 2$. We conclude that the Neel ordering disappears in the range $2 < n < 3$.

I. INTRODUCTION

It is known that properties of quantum spin systems tend to approach those of their corresponding classical spin systems as the spin magnitude increases. However, this is not necessarily the case for a sequence of models in which the number of multipolar-interaction terms increases as the spin magnitude increases. Consequently the higher-spin system may have stronger quantum effects in this case. In other words, such additional terms can break a classical correspondence down even in high dimensions. These "large-spin-magnitude" systems mentioned above include systems in which one unit has more than two degrees of freedom, such as orbitally degenerate systems^{1,2,3}. In such systems, many coupling constants appear in general. However, experimental information about multipolar couplings is limited. As a starting point to explore such systems, understanding of one of the extreme limits must be useful. Therefore in this paper, we investigate an $SU(n)$ symmetric case^{4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22} in three dimensions, where n is the total number of internal degrees of freedom, by means of high-temperature series expansion (HTSE).

We consider a simple cubic lattice. Let each site take one of the n colors denoted by j with $j = 1, 2, \dots, n$. Using the Hubbard operator $X_{ij} = |j\rangle\langle i|$, the exchange operator is expressed as

$$P_{ij} = \sum_{j=1}^n \sum_{i=1}^n X_i X_j : \quad (1)$$

Colors of sites i and j are exchanged when P_{ij} is applied. Then, an $SU(n)$ symmetric Hamiltonian reads

$$H = J \sum_{\langle ij \rangle} P_{ij} \quad (2)$$

where the summation is taken over all the nearest neighbors. We consider the antiferromagnetic case, $J > 0$. Let us show the relations with spin operators explicitly for some of the special cases below.

(i) When $n = 2$, this model is reduced to the ordinary Heisenberg model with $s = 1/2$ by relation $2P_{ij} = 4s_i \cdot s_j$.

(ii) The $SU(3)$ case corresponds to $s = 1$, competing quadratic and biquadratic exchange interaction,²³

$$1 + P_{ij} = s_i \cdot s_j + (s_i \cdot s_j)^2 : \quad (3)$$

(iii) The $SU(4)$ Heisenberg model is related to spin $3/2$ systems but more often discussed in the context of orbital- and spin-degenerate systems^{10,11,12,13,14,15,16,17,18,19,20,21}. In particular, the model in three dimensions has been used as an effective model of CeB_6 to explain magnetic field dependence of the transition temperature of an antiferro-orbital ordering.^{20,21} The four local states can be represented by $|\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle, |\downarrow\uparrow\rangle, |\downarrow\downarrow\rangle$, where $|\uparrow\rangle$ and $|\downarrow\rangle$ represent an orbital state and a spin state, respectively. The pseudo-spin operators are defined by $t^z_j = \frac{1}{2}j_z$, $t^+_j = |\downarrow\uparrow\rangle\langle\uparrow\uparrow| + |\uparrow\downarrow\rangle\langle\downarrow\uparrow|$, $t^-_j = |\uparrow\uparrow\rangle\langle\downarrow\uparrow| + |\downarrow\downarrow\rangle\langle\uparrow\downarrow|$, and the exchange operator for $n = 4$ is rewritten as

$$P_{ij} = t_i \cdot t_j + s_i \cdot s_j + 4(t_i \cdot t_j)(s_i \cdot s_j) + \frac{1}{4} : \quad (4)$$

The Pauli matrices $\sigma = 2t$ and $\tau = 2s$ may simplify this expression, i.e.,

$$4P_{ij} = \sigma_i \cdot \sigma_j + \tau_i \cdot \tau_j + (\sigma_i \cdot \sigma_j)(\tau_i \cdot \tau_j) + 1 : \quad (5)$$

Note that there is a different representation of $SU(n)$ Heisenberg model studied in detail using Quantum Monte Carlo method (QMC)²⁴. However, the QMC for the Hamiltonian (2) suffers from minus sign problems in more than one dimensions^{11,24}.

If the number of competing order parameters is large and frustration exists, the transition temperature can greatly be reduced from the mean-field value even in three dimensions. That will be the case with the Hamiltonian (2). First, it is isotropic with respect to $n^2 - 1$ independent interacting components. Furthermore, it contains frustration as most clearly seen in Eq. (5) of the $SU(4)$ case. Each of the 15 components, $\sigma_x, \sigma_y, \sigma_z, \tau_x, \tau_y, \tau_z, \sigma_x\tau_x, \sigma_y\tau_y, \sigma_z\tau_z, \sigma_x\tau_y, \sigma_y\tau_x, \sigma_x\tau_z, \sigma_z\tau_x, \sigma_z\tau_y$, attempts an antiparallel alignment. However, it cannot




FIG. 2: The Fourier transform of the correlation function as a difference from its high-temperature limit, $S(q; q; q)$, for the SU(16) model.

bad convergence for several q . This is probably because information at position x is lacking when $\cos(x - q) \neq 0$. In order to avoid it, we extrapolate the series of a complex function $\sum_{j=0}^{\infty} hX_j X_{j+x} i e^{ix} \Psi(x)$, and after that take the real part of the extrapolated function. Here, the summation is taken for $x=0, y=0, z=0$, and $w(x)$ is two to the power of the number of nonzeros in $x; y; z$. This method also makes use of its imaginary part, and the convergence becomes better than the naive extrapolation. As we have seen in the next-nearest-neighbor correlation function, the analysis becomes easier as n increases. Therefore, we show $S(q)$ of the SU(16) case in Fig. 2 for a diagonal direction in the q -space, $S(q; q; q)$. Here, its high-temperature limit, which does not depend on q , is subtracted. As temperature decreases, antiferromagnetic correlation develops and $S(q; q; q)$ increases as the curve at $T = 2J$ shows. However, with further decrease of temperature, $S(q; q; q)$ starts decreasing and $S(q)$ with other q increases. It is not very clear from Fig. 2 if the maximum of $S(q)$ starts moving from $(\pi; \pi; \pi)$. Hence, let us show another quantity. Note that if the second derivative of $S(q)$ at $q = (\pi; \pi; \pi)$ changes sign, the position of the maximum clearly moves. Figure 3 shows its temperature dependence for the SU(16) model. It shows that it changes the sign around $T = J$. In fact, this temperature of changing sign seems to hardly depend on n of SU(n). The result above suggests that the Neel order disappear at least in the SU(n) model with large n , i.e., order with wave number $q \notin (\pi; \pi; \pi)$, or disorder, should appear. Then, the next question is, at which n the Neel order disappears.

III. NEEL TEMPERATURE

It is known that the SU(2) Heisenberg model has the Neel order at low temperature. Therefore in this sec-




FIG. 3: Temperature dependence of $d^2S(q; q; q)/dq^2$ at $q = (\pi, \pi, \pi)$ for the SU(16) model by two different PAs.

tion, we gradually increase n of SU(n) from $n = 2$. As a preparation for that, we find a reliable way to estimate a transition temperature first, using the SU(2) model. The Neel temperature T_N can be characterized by divergence of $S(q; q; q)$, namely, by a singularity of $S(q; q; q)$ as a function of J . In order to analyze such a singularity, we use so-called D-log-Pade approximation (DLPA), i.e., the PA for the logarithmic derivative. In using the DLPA, transformation of the expansion variable may improve the convergence of extrapolation. We choose a transformation^{27,28}

$$J = \frac{x}{1 - a^2 x^2}; \quad (8)$$

where a is an adjustable parameter that improves convergence. In fact, the singularity closest to the origin in the original series is near the imaginary axis, while T_N corresponds to a singularity on the real axis. With this transformation, singularities on the real axis approach to the origin and those near the imaginary axis go away. Since the DLPA can estimate a position of the nearest singularity most accurately, errors of the DLPA can become smaller by this transformation.

In order to find an optimal a , we calculate T_N as a function of a for a couple of choices of the DLPA, and we adopt a at which difference among different DLPA is the smallest. Figure 4 shows T_N as a function of a for three different choices of the DLPA. Here, $[m/n]$ denotes the DLPA with a polynomial of order m over a polynomial of order n . Since $[4/4]$ is from the series one-order lower than the others, we choose a at which $[5/4]$ and $[4/5]$ are the closest, namely, $a = 1.04$. Then, $T_N = J$ obtained here is 1.898, which is close to those in the literature, 1.892 by the QMC²⁹ and 1.888 by the HTSE³⁰. In addition, we have obtained a critical exponent simultaneously. We assume that the critical exponent does not depend on the spin magnitude, and compare it with that of the classical antiferromagnetic Heisenberg model, in which $S(q; q; q)$ is identical to the staggered susceptibility equivalent to




FIG. 4: The Neel Temperature T_N of the SU (2) Heisenberg model obtained by three different DLPAs as a function of the parameter a .

the uniform susceptibility of the ferromagnetic model. Hence we can compare the critical exponent. The estimation from our calculation above is $\beta = 1.399$, which is close to 1.396 by a Monte Carlo method³¹, 1.406 by HTSE³², and 1.388 by a field theory³³ of the classical O (3) model. Therefore, we trust this way of analyzing T_N , and use it also for the SU (n) model with $n > 2$.

Originally n is an integer because it is the number of internal degrees of freedom. However, after obtaining the series as an analytic function of n , we can regard n as a continuous variable that has a physical meaning only when it happens to take integer values. In fact, as n changes continuously, the properties of the series change also continuously. Therefore, we gradually increase n from $n = 2$ and see n -dependence of T_N . Figure 5 shows the results. Here, we fix $a = 1.04$, and we plot [5/4] and [4/5] together. The optimal a may depend on n . However, since the difference between the two curves is very small, we regard it as an optimal a . As n increases, T_N decreases almost linearly, and at $n = 2.45$ the singularity corresponding to T_N runs away from the real axis to the complex value with a finite imaginary part. For larger n , we do not find any singularity on the antiferromagnetic side of the real axis. Therefore, we conclude that the Neel order disappears in the range $2 < n < 3$.

In addition, we have also analyzed $S(q)$ with $q \in (\pi, 2\pi)$. However, we have not found any symptom of ordering with $q \in (\pi, 2\pi)$ in the temperature region that we can reach by the present order of the series.

IV. SUMMARY

In summary, we have performed high-temperature series expansions for the SU (n) Heisenberg model in three

dimensions with arbitrary n . First of all, we have calculated a next-nearest-neighbor correlation function. At least at large n , it changes the sign at low temperature, which suggests that the correlation should not be like




FIG. 5: The Neel Temperature T_N of the SU (n) Heisenberg model as a function of n with $a = 1.04$. Here, [5/4] and [4/5] are plotted together.

Neel order, but have a longer period. Analysis of the Fourier transform of the correlation function also supports that the ground state of the large- n SU (n) Heisenberg model does not have the Neel-order correlation. Then, we have turned to an approach from $n = 2$. Since the SU (2) Heisenberg model has the Neel order, it should disappear at a certain n . We have first found a reliable way to estimate a transition temperature by analyzing the divergence of the (π, π) component of the correlation function. Next, we generalize n to a continuous variable and increase n gradually from $n = 2$. We have concluded that the Neel order disappears in the range $2 < n < 3$.

Acknowledgments

The author would like to thank Y. Kuramoto for directing him to this topic and for stimulating discussions. He also appreciates useful comments on the manuscript from A. Honecker. This work was supported in part by the Deutsche Forschungsgemeinschaft (DFG). Early stages of this work were supported by the Japan Society for the Promotion of Science and by the Max-Planck Institute for the Physics of Complex Systems. Different stages of this work were supported by the DFG and the Technical University Braunschweig. Parts of the numerical calculations were performed on the cfgauss at the computing centers of the TU Braunschweig.

- (1982)
- ² N. Fukushima and Y. Kuramoto, J. Phys. Soc. Jpn. 67, 2460 (1998).
 - ³ Y. Kuramoto and N. Fukushima, J. Phys. Soc. Jpn. 67, 583 (1998).
 - ⁴ G. V. Uimin, JETP Lett. 12, 225 (1970).
 - ⁵ B. Sutherland, Phys. Rev. B 12, 3795 (1975).
 - ⁶ A. Fujii, A. Klumper, Nucl. Phys. B 546, 751 (1999).
 - ⁷ I. A. A. Eck, Nucl. Phys. B 265, 409 (1986).
 - ⁸ N. Kawakami, Phys. Rev. B 46, 3191 (1992).
 - ⁹ M. T. Batchelor, X.-W. Guan, N. Oelkers, K. Sakai, Z. Tsuboi, A. Foerster, Phys. Rev. Lett. 91, 217202 (2003).
 - ¹⁰ Y. Yamashita, N. Shibata and K. Ueda, Phys. Rev. B 58, 9114 (1998).
 - ¹¹ B. Frischmuth, F. M. Ila and M. Troyer, Phys. Rev. Lett. 82, 835 (1999).
 - ¹² N. Fukushima and Y. Kuramoto, J. Phys. Soc. Jpn. 71, 1238 (2002).
 - ¹³ N. Fukushima, J. Stat. Phys. 111, 1049 (2003).
 - ¹⁴ Y. Q. Li, M. Ma, D. N. Shi, and F. C. Zhang, Phys. Rev. Lett. 81, 3527 (1998).
 - ¹⁵ M. van den Bossche, P. Azaria, P. Lecheminant, F. M. Ila, Phys. Rev. Lett. 86, 4124 (2001).
 - ¹⁶ M. van den Bossche, F. C. Zhang, and F. M. Ila, Eur. Phys. J. B 17, 367-370 (2000).
 - ¹⁷ A. Mishra, M. Ma, and F. C. Zhang, Phys. Rev. B 65, 214411 (2002).
 - ¹⁸ S. Q. Shen, Phys. Rev. B 66, 214516 (2002).
 - ¹⁹ K. Penc, M. Mambrini, P. Fazekas, F. M. Ila, Phys. Rev. B 68, 12408 (2003).
 - ²⁰ F. J. Ohkawa: J. Phys. Soc. Jpn. 54, (1985). 3909.
 - ²¹ R. Shiina, H. Shiba and P. Thalmeier, J. Phys. Soc. Jpn. 66, 1741 (1997).
 - ²² H. H. Chen and R. K. Joseph, J. Math. Phys. 13, 725 (1972).
 - ²³ N. Papanicolaou, Nucl. Phys. B 305, 367 (1988).
 - ²⁴ K. Harada and N. Kawashima, Phys. Rev. B 65, 52403 (2002).
 - ²⁵ G. S. Rushbrooke, G. A. Baker, Jr., and P. J. Wood, in Phase Transitions and Critical Phenomena, edited by C. Domb and M. S. Green (Academic, London, 1974), Vol. 3, p. 245.
 - ²⁶ D. C. Handscomb, Proc. Camb. Phil. Soc. 60, 115 (1964).
 - ²⁷ A. J. Guttmann, in Phase Transitions and Critical Phenomena, edited by C. Domb and J. L. Lebowitz (Academic, San Diego, 1989), vol. 13, p. 1.
 - ²⁸ K. K. Pan, Phys. Rev. B 59, 1168 (1999).
 - ²⁹ A. W. Sandvik, Phys. Rev. Lett. 80, 5196 (1998).
 - ³⁰ J. Oitmaa and W. Zheng, preprint (cond-mat/0409041).
 - ³¹ M. Campostrini, M. Hasenbusch, A. Pelissetto, P. Rossi, E. Vicari, Phys. Rev. B 65, 144520 (2002).
 - ³² P. Butera and M. Comi, Phys. Rev. B 56, 8212 (1997).
 - ³³ F. Jasch and H. Kleinert, J. Math. Phys. 42, 52 (2001).