

# Computation of the Kolmogorov-Sinai entropy using statistical mechanics: Application of an exchange Monte Carlo method

SHIN-ICHI SASA<sup>1</sup> (\*) and KUMIKO HAYASHI<sup>1</sup> (\*\*)

<sup>1</sup> *Department of Pure and Applied Sciences, University of Tokyo, Komaba, Tokyo 153-8902, Japan*

PACS. 75.10.Nr – First pacs description.

PACS. 05.20.-y – Second pacs description.

PACS. 05.45.Ac – Third pacs description.

**Abstract.** – We propose a method for computing the Kolmogorov-Sinai (KS) entropy of chaotic systems. In this method, the KS entropy is expressed as a statistical average over the canonical ensemble for a Hamiltonian with many ground states. This Hamiltonian is constructed directly from an evolution equation that exhibits chaotic dynamics. As an example, we compute the KS entropy for a chaotic repeller by evaluating the thermodynamic entropy of a system with many ground states.

Chaotic systems are characterized by properties of  $O(e^N)$  distinguishable trajectories over a time interval  $N$  in a phase space in which the limit of observational accuracy is given by some finite value  $\epsilon$  [1–3]. In particular, if these trajectories are coded as sequences of symbols, the optimal compression ratio of such sequences is an important quantity from the point of view of information theory [4, 5]. This quantity corresponds to the Kolmogorov-Sinai (KS) entropy,  $h_{\text{ks}}$ .

Although the KS entropy provides a simple characterization of chaotic systems, its direct numerical evaluation is not simple. One reason for this is that a precise mathematical definition of distinguishable trajectories is difficult to investigate numerically [1]. However, this difficulty can be overcome when a set of periodic orbits alone can provide information that is sufficient to obtain a useful characterization of the set of all distinguishable trajectories [6–8]. However, even in such cases, the difficulty involved in numerical computations remains, because the number of periodic orbits of period  $N$  increases exponentially as a function of  $N$ .

Here, let us recall that in the statistical mechanics of glassy systems [9] and decision problems [10], the number of explored states increases exponentially as a function of the system size. Recently, efficient methods for computing thermodynamic quantities of glassy systems have been developed [11, 12] and applied to several problems [13–15]. Inspired by this success, in this paper, we propose a method for computing the KS entropy employing a quantity that can be calculated in a system with many ground states.

---

(\*) E-mail:sasa@jiro.c.u-tokyo.ac.jp

(\*\*) E-mail:hayashi@jiro.c.u-tokyo.ac.jp

Specifically, we consider a map  $\Gamma_{n+1} = F(\Gamma_n)$ , with  $\{\Gamma_n\} \in \Omega \subset \mathbf{R}^d$ , and assume that its trajectories can be coded with sequences of  $D$  symbols. The frequency distribution of an  $N$  symbol sequence  $[s^k] = (s_1^k, s_2^k, \dots, s_N^k)$ , with the natural measure for the initial conditions, is denoted by  $\mu([s^k])$ , where  $s_n^k \in \{1, \dots, D\}$ , and  $k$  is the index identifying the sequence. Letting  $M(N)$  be the total number of possible symbol sequences, the KS entropy,  $h_{\text{ks}}$ , is defined in terms of the Shannon entropy rate [4] as

$$h_{\text{ks}} \equiv - \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^{M(N)} \mu([s^k]) \log \mu([s^k]). \quad (1)$$

Note that  $h_{\text{ks}}/\log(D)$  represents the optimal compression ratio of symbol sequences [5].

In order to study simple cases, we further assume that for each  $[s^k]$  there exists an  $N$ -periodic orbit  $\Gamma_1^k, \Gamma_2^k, \dots, \Gamma_N^k, \Gamma_1^k$  such that the symbol corresponding to  $\Gamma_i^k$  is  $s_i^k$  for all  $i \leq N$ . Then there is a one-to-one correspondence between the set of all symbol sequences  $\{[s^k]\}$  and the set of all  $N$ -periodic orbits  $\{[\Gamma^k]\}$ . Furthermore, when  $N$  is sufficiently large,  $\mu([s^k])$  is accurately approximated by the probability

$$p_k \equiv \frac{1}{Z_d(N)} e^{-N\lambda([\Gamma^k])}, \quad (2)$$

where  $\lambda([\Gamma^k])$  is the volume expansion ratio in the unstable directions around the periodic orbit  $[\Gamma^k]$ . The normalization constant  $Z_d$  is determined by the condition  $\sum_{k=1}^{M(N)} p_k = 1$ . In terms of  $p_k$ , the KS entropy is expressed as

$$h_{\text{ks}} = - \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^{M(N)} p_k \log p_k. \quad (3)$$

Note that the relation  $\lim_{N \rightarrow \infty} \log Z_d(N)/N = 0$  holds for many physically interesting systems. In this case, the expression (3) leads to the useful relation (Pesin's formula) that  $h_{\text{ks}}$  is equal to the sum of the positive Lyapunov exponents [1, 3]. However, there are systems to which this formula cannot be applied. A typical example is a chaotic repeller. Because the KS entropy of a repeller is related to transport properties of nonequilibrium phenomena described by deterministic models [16], it is important to study a computational method for  $h_{\text{ks}}$  on the basis of (3) that does not employ Pesin's formula.

*Basic formula.* – Here and hereafter, we use  $[\Gamma] = (\Gamma_1, \dots, \Gamma_N)$  to represent an array of  $N$  phase space points, not a trajectory given by the map  $\Gamma_{n+1} = F(\Gamma_n)$ . We also introduce  $\Gamma_0$  as  $\Gamma_0 = \Gamma_N$  for later convenience. Let  $\Lambda_m([\Gamma])$ , with  $1 \leq m \leq d$ , be the eigenvalues of the matrix  $F'(\Gamma_N)F'(\Gamma_{N-1}) \cdots F'(\Gamma_1)$ , where  $F'(\Gamma_n)$  is the  $d \times d$  Jacobi matrix determined from the map  $\Gamma_{n+1} = F(\Gamma_n)$ . The volume expansion ratio  $\lambda([\Gamma^k])$  in the unstable directions around the periodic orbit  $[\Gamma^k]$  is defined as

$$\lambda([\Gamma^k]) \equiv \frac{1}{N} \sum_{m; |\Lambda_m([\Gamma^k])| > 1} \log |\Lambda_m([\Gamma^k])|. \quad (4)$$

Here, we assume that all periodic orbits are hyperbolic and hence satisfy  $|\Lambda_m| \neq 1$  for  $1 \leq m \leq d$ . The volume expansion ratio  $\lambda([\Gamma])$  for an arbitrary array  $[\Gamma]$  can be defined by replacing  $\Gamma^k$  in (4) with  $\Gamma$ .

For an arbitrary function  $A([\Gamma])$ , we define the quantity

$$I(A) \equiv \sum_{k=1}^{M(N)} A([\Gamma^k]) e^{-N\lambda([\Gamma^k])}. \quad (5)$$

Substituting (2) into (3) and using (5), we rewrite  $h_{\text{ks}}$  as

$$h_{\text{ks}} = \lim_{N \rightarrow \infty} \left[ \frac{I(\lambda)}{I(1)} + \frac{1}{N} \log I(1) \right]. \quad (6)$$

We now seek to express  $I(A)$  in a form that can be computed easily. We first rewrite (5) as

$$I(A) = \sum_{k=1}^{M(N)} \left[ \prod_{n=1}^N \int d\Gamma_n \delta(\Gamma_n - \Gamma_n^k) \right] A([\Gamma]) e^{-N\lambda([\Gamma])}. \quad (7)$$

Then, defining  $\phi_n([\Gamma_n]) \equiv \Gamma_n - F(\Gamma_{n-1})$ ,  $n = 1, 2, \dots, N$ , we obtain the formula

$$\prod_{n=1}^N \delta(\phi_n) = \sum_{k=1}^{M(N)} \frac{\prod_{n=1}^N \delta(\Gamma_n - \Gamma_n^k)}{|J([\Gamma])|}, \quad (8)$$

where

$$J([\Gamma]) = \det \frac{\partial(\phi_1, \dots, \phi_N)}{\partial(\Gamma_1, \dots, \Gamma_N)}. \quad (9)$$

Here, we note that  $J([\Gamma^k]) \neq 0$  for all hyperbolic periodic orbits, because in this case it can be proved that

$$J([\Gamma]) = \prod_{m=1}^d (1 - \Lambda_m([\Gamma])). \quad (10)$$

Also, from the expression (10), we can derive

$$|J([\Gamma^k])| = e^{N\lambda([\Gamma^k]) + o(N)}. \quad (11)$$

Then, expressing the delta function as

$$\delta(x) = \lim_{\beta \rightarrow \infty} \sqrt{\frac{\beta}{\pi}} e^{-\beta x^2}, \quad (12)$$

and ignoring the sub-exponential contribution, we can rewrite  $I(A)$  as

$$I(A) = \lim_{\beta \rightarrow \infty} \left( \frac{\beta}{\pi} \right)^{Nd/2} \int d\Gamma_1 \dots d\Gamma_N e^{-\beta H([\Gamma])} A([\Gamma]), \quad (13)$$

with

$$H([\Gamma]) = \sum_{n=1}^N |\Gamma_n - F(\Gamma_{n-1})|^2. \quad (14)$$

The expression (13) leads us to introduce a canonical ensemble in the form

$$p_c([\Gamma]) = \frac{1}{Z(\beta)} e^{-\beta H([\Gamma])}, \quad (15)$$

with the partition function

$$Z(\beta) = \int d\Gamma_1 \cdots d\Gamma_N e^{-\beta H(\Gamma)}. \quad (16)$$

Then, (13) becomes

$$I(A) = \lim_{\beta \rightarrow \infty} \left( \frac{\beta}{\pi} \right)^{Nd/2} \langle A \rangle_{\beta} Z(\beta), \quad (17)$$

where  $\langle \cdot \rangle_{\beta}$  represents the average taken with respect to (15). Substituting this result into (6), we obtain the formula

$$h_{\text{ks}} = \lim_{N \rightarrow \infty} \lim_{\beta \rightarrow \infty} \left[ \langle \lambda \rangle_{\beta} + \frac{d}{2} \log \left( \frac{\beta}{\pi} \right) + \frac{1}{N} \log Z(\beta) \right]. \quad (18)$$

Recall that the Hamiltonian (14) is non-negative and is equal to zero when  $[\Gamma]$  coincides with a periodic orbit  $[\Gamma^k]$ . Also, the number of periodic orbits  $M(N)$  is of order  $e^N$  for a chaotic map  $F$ . These facts imply that the KS entropy,  $h_{\text{ks}}$ , can be obtained using the formalism of statistical mechanics with the Hamiltonian (14), which possesses  $O(e^N)$  ground states.

*Computational method:* – Let us define the entropy function  $S(E)$  through the relation

$$e^{S(E)} dE = \int d\Gamma_1 \cdots d\Gamma_N E \leq H([\Gamma]) \leq E + dE. \quad (19)$$

We then evaluate the partition function  $Z(\beta)$  as

$$Z(\beta) = e^{S(E_*(\beta)) - \beta E_*(\beta) + O(\log(N))}, \quad (20)$$

where  $E_*(\beta)$  is defined as

$$\frac{\partial S(E_*)}{\partial E_*} = \beta. \quad (21)$$

Then, from a standard argument employed in statistical mechanics, the relation  $E_*(\beta) = \langle H \rangle_{\beta}$  is obtained, and the quantity  $\tilde{S}(\beta) \equiv S(E_*(\beta))$  is calculated from the identity

$$-\beta \frac{\partial \tilde{S}(\beta)}{\partial \beta} = C(\beta), \quad (22)$$

where  $C(\beta)$  represents the heat capacity defined by

$$C(\beta) \equiv \beta^2 \langle (\delta E)^2 \rangle_{\beta}. \quad (23)$$

Here, because  $C_{\infty} \equiv \lim_{\beta \rightarrow \infty} C(\beta)$  takes a nonzero value,  $\tilde{S}(\beta)$  diverges as  $\sim \log \beta$  in the limit  $\beta \rightarrow \infty$ . Therefore, in order to avoid a singularity, we define  $\tilde{S}_0(\beta)$  according to

$$\tilde{S}(\beta) = -C_{\infty} \theta(\beta - 1) \log \beta + \tilde{S}_0(\beta), \quad (24)$$

where  $\theta(\cdot)$  is the Heaviside function. Then, from (22) and (24), we can express  $\tilde{S}_0(\beta)$  in terms of  $C(\beta)$ , using the condition

$$e^{\tilde{S}_0(0) + o(N)} = |\Omega|^N. \quad (25)$$

Combining this result with (20) and (24), we rewrite the second and third terms of (18) as

$$-f(N) \equiv \log \frac{|\Omega|}{(\pi e)^{d/2}} - \frac{1}{N} \left[ \int_0^\infty d\beta \frac{C(\beta) - C_\infty \theta(\beta - 1)}{\beta} \right], \quad (26)$$

where we have used the relation  $\lim_{\beta \rightarrow \infty} \beta E_*(\beta) = C_\infty = Nd/2$ . Using this quantity, the KS entropy given by (18) can be expressed as

$$h_{\text{ks}} = \lim_{N \rightarrow \infty} \left[ \lim_{\beta \rightarrow \infty} \langle \lambda \rangle_\beta - f(N) \right]. \quad (27)$$

Therefore, the KS entropy is obtained from  $\langle \lambda \rangle_\beta$  and  $C(\beta)$ , which can be computed using the canonical ensemble (15) with the Hamiltonian (14).

As one method to obtain the canonical ensemble, we consider the Langevin equation

$$\dot{\Gamma}_n(t) = -\frac{\partial H([\Gamma])}{\partial \Gamma_n} + \xi_n, \quad (28)$$

where  $\xi_n$  represents Gaussian white noise satisfying

$$\langle \xi_n(t) \xi_m(t') \rangle = 2\beta^{-1} \delta(t - t') \delta_{nm}. \quad (29)$$

The steady distribution of arrays  $[\Gamma]$  in this system is given by the canonical ensemble (15) with the Hamiltonian (14). However, because there are  $O(e^N)$  ground states of  $H$ , it is difficult to obtain accurately the canonical ensemble (15) by direct numerical integration of (28) when  $\beta$  is large. For this reason, we employ an exchange Monte Carlo method [12] in order to improve the accuracy of the numerical computations. Below we describe this method.

We first prepare  $L$  copies of the Langevin system. Each system is subject to noise with a time-dependent inverse temperature  $\beta^{(i)}(t)$  that takes values in the set  $\{\beta_0^{(1)}, \dots, \beta_0^{(L)}\}$ . We have the initial values  $\beta^{(i)}(0) = \beta_0^{(i)}$ , and the time evolution of each  $\beta^{(i)}(t)$  is described by the following rule: At time  $k\tau$  with  $k = 1, 2, \dots$ , two inverse temperatures  $\beta^{(j)}$  and  $\beta^{(j')}$ , given by  $\beta^{(j)} = \beta_0^{(i)}$  and  $\beta^{(j')} = \beta_0^{(i+1)}$  for some  $i$  satisfying  $i = (1 - (-1)^k)/2 + 2i' - 1$ , with  $i' = 1, 2, \dots$ , are exchanged with probability

$$\frac{1}{1 + e^{(\beta_0^{(i+1)} - \beta_0^{(i)}) (H([\Gamma]^{(i)}) - H([\Gamma]^{(i+1)}))}}, \quad (30)$$

where  $[\Gamma]^{(i)}$  is the configuration of the system subject to the noise with the inverse temperature  $\beta_0^{(i)}$ . It can be shown that the steady distribution of  $([\Gamma]^{(1)}, \dots, [\Gamma]^{(L)})$  satisfies

$$P_c([\Gamma]^{(1)}, \dots, [\Gamma]^{(L)}) = \frac{1}{\tilde{Z}} e^{-\sum_{i=1}^L \beta_0^{(i)} H([\Gamma]^{(i)})}, \quad (31)$$

where  $\tilde{Z}$  is a normalization constant. It has been demonstrated that the exchange Monte Carlo method is useful for calculating thermodynamic quantities in systems with many metastable states [9], and therefore we expect that this method will be effective in application to the present problem.

*Example.* – As a simple demonstration of the computational method described above, we study a map  $F$  given by

$$\begin{aligned} x_{n+1} &= x_n + z_n, \\ z_{n+1} &= z_n + \frac{4}{g} \cos(x_n + z_n), \end{aligned} \quad (32)$$




Fig. 1




Fig. 2

Fig. 1 – Heat capacity as a function of  $\log \beta$ . The circles, squares, triangles, stars and pluses represent the results for the cases  $N = 4, 8, 12, 16$  and  $20$ .

Fig. 2 –  $\langle \lambda \rangle_\beta$  as a function of  $\log \beta$ . Here,  $N = 20$ .

which is defined in the region  $\Omega = \{(x, z) | 0 \leq x \leq 2\pi, 0 \leq x + z \leq 2\pi\} \subset \mathbf{R}^2$ . This map is obtained as an approximation of a two-dimensional map that describes a bouncing ball on a vibrating table [2]. It also becomes the so-called standard map under a variable transformation. We wish to compute the KS entropy of the repeller of  $\Omega$ , which is defined as the set  $\{\Gamma | F^n(\Gamma) \in \Omega \text{ for all } n \in \mathbf{Z}\}$ .

Specifically, we investigate the case  $g = 0.3$ . In the numerical simulations, we set  $\tau = 10^{-3}$  and  $\beta_0^{(i)} = 1.4^{-(i-8)}$ , where  $1 \leq i \leq L = 30$ , with a time step  $\Delta t = 10^{-5}$  for the integration of (28). The heat capacity,  $C(\beta)/N$ , is plotted for  $N = 4, 8, 12, 16$  and  $20$  in Fig. 1. From the value of the heat capacity thus obtained numerically, we derive the estimation  $\lim_{N \rightarrow \infty} f(N) = 1.8(3)$ . Furthermore,  $\langle \lambda \rangle_\beta$  as a function of  $\beta$  is displayed in Fig. 2. Using this result, we derive  $\langle \lambda \rangle_\infty = 2.5(0)$ . Thus, from (26), we obtain  $h_{\text{KS}} = 0.6(7)$ .

Let us check the validity of our computational result. First, because the escape rate formula holds for hyperbolic maps [17], the value  $\lim_{N \rightarrow \infty} f(N)$  should be equal to the escape rate,  $\gamma$ . Directly computing the escape rate, we obtain  $\gamma = 1.8(3)$ . Thus, our computational result is consistent with the escape rate formula. Second, in this model, we can directly evaluate the right-hand side of (1) by using the binary symbols 0 and 1 for  $x < \pi$  and  $x \geq \pi$ , respectively. Collecting many segments of trajectories on the repeller over the time intervals  $-\tilde{N} \leq n \leq \tilde{N}$ , with for  $\tilde{N} = 1, 2$  and  $3$ , we obtain  $h_{\text{KS}} = 0.6(8)$ . This value is also consistent with that found using our method.

*Discussion.* – Let us finally comment on topics related to the basic formula (18). First, the statistical mechanical formulation used in the present paper is not related to the so-called thermodynamic formalism of chaos [4], in which, instead of (2), the probability

$$p_k = \frac{1}{Z_d(\tilde{\beta}, N)} e^{-N\tilde{\beta}\lambda(\Gamma^k)} \quad (33)$$

is assumed for each periodic orbit. In a manner similar to that used here,  $Z_d(\tilde{\beta}, N)$  can be obtained with a statistical mechanical formulation, but the Hamiltonian for cases in which  $\tilde{\beta} \neq 1$  includes the Jacobian  $J([\Gamma])$ , and this makes it difficult to carry out the computation.

Second, it is well known that the chaotic trajectories of the standard map correspond to the stationary configurations of the Frenkel-Kontorowa model [18]. However, such a corre-

spondence can be found only for a certain class of maps. Also, even for the standard map, it seems difficult to develop a method for computing the KS entropy by using this correspondence, because periodic orbits correspond to local minima and maxima, not ground states of the Frenkel-Kontorowa Hamiltonian. Nevertheless, it is worth noting that this correspondence has led to a mathematical study to evaluate the lower bound of the topological entropy [19].

Finally, let us recall that the expression (18) is valid in the case that none of the eigenvalues  $\Lambda_m$  ( $1 \leq m \leq d$ ) of the periodic orbits is equal to 1. However, even if there is a non-hyperbolic periodic orbit whose eigenvalue  $\Lambda_m$  is unity, the right-hand side of the expression (18) can be computed. It might be an interesting problem to investigate the expression (18) for non-hyperbolic systems.

In conclusion, we derived the formulae (18) and (27) for the KS entropy. We then applied these formulae to compute the KS entropy for the case of the repeller of a simple map by using the corresponding Langevin equation together with the exchange Monte Carlo method. We hope that these formulae can be similarly applied to many problems.

\* \* \*

The authors acknowledge K. Hukushima for teaching them some techniques of the exchange Monte Carlo method. They also thank Y. Ishii for informing them of Ref. [19]. This work was supported by a grant from the Ministry of Education, Science, Sports and Culture of Japan (No. 16540337) and a Research Fellowship for Young Scientists from the Japan Society for the Promotion of Science (No. 1711222).

## REFERENCES

- [1] KATOK A. and HASSELBLATT B., *Introduction to the Modern Theory of Dynamical Systems* (Cambridge University Press, Cambridge) 1995.
- [2] GUCKENHEIMER J. and HOLMES P., *Nonlinear Oscillations, dynamical Systems and Bifurcations of Vector Fields* (Springer, Berlin) 1983.
- [3] DORFMAN J. R., *An introduction to Chaos in Nonequilibrium Statistical Mechanics* (Cambridge University Press, Cambridge) 1999.
- [4] BECK C. and SCHLÖGEL F., *Thermodynamic formalism* (Cambridge University Press, Cambridge) 1993.
- [5] COVER T. M. and THOMAS J. A., *Elements of Information Theory* (Wiley) 1991.
- [6] KAI T. and TOMITA K., *Prog. Theor. Phys.*, **64** (1980) 1532.
- [7] GUNARATNE G. and PROCACCIA I., *Phys. Rev. Lett.*, **59** (1987) 1377.
- [8] CVITANOVIĆ P., *Phys. Rev. Lett.*, **61** (1988) 2729.
- [9] BRAY A. J. and MOORE M. A., *J. Phys. C: Solid St. Phys.*, **13** (1980) L469.
- [10] MONASSON R. AND ZECCHINA R., *Phys. Rev. Lett.*, **76** (1996) 3881.
- [11] BERG B. A. AND NEUHAUS T., *Phys. Lett. B*, **267** (1991) 249.
- [12] HUKUSHIMA K. and NEMOTO K., *J. Phys. Soc. Jpn.*, **65** (1996) 1604.
- [13] PINN K. and WIECZERKOWSKI C., *Int. J. Mod. Phys. ,* **C9** (1998) 541.
- [14] LIMA A. P. and ARGOLL DE MENEZES M., *Phys. Rev. E*, **63** (2001) 020106.
- [15] HUKUSHIMA K., *Computer Physics Communications*, **147** (2002) 77.
- [16] GASPARD P. and NICOLIS G., *Phys. Rev. Lett.*, **65** (1990) 1693.
- [17] KADANOFF L. P. and TANG C., *Proc. Natl. Acad. Sci.*, **81** (1984) 1276.
- [18] AUBRY S. and ABRAMOVICI G., *Physica D*, **43** (1990) 199.
- [19] KNILL O., *Trans. Amer. Math. Soc.*, **348** (1996) 2999.