

Fluctuation-dissipation theorem in an aging colloidal glass

Sara Jabbari-Farouji¹, Daisuke Mizuno², Maryam Atakhorrami², Fred C. MacKintosh²,
Christoph F. Schmidt², Erika Eiser³, Gerard H. Wegdam¹ and Daniel Bonn^{1,4}

¹Van der Waals-Zeeman Institute, University of Amsterdam, 1018XE Amsterdam, the Netherlands

²Department of Physics and Astronomy, Vrije Universiteit Amsterdam, 1081HV Amsterdam, The Netherlands

³van't Hoff Institute for Molecular Sciences, University of Amsterdam, 1018WV Amsterdam, The Netherlands and

⁴Laboratoire de Physique Statistique, UMR 8550 CNRS associated with the University of Paris 6 and Paris 7,
Ecole Normale Supérieure, 75231 Paris Cedex 05, France

(Dated: February 25, 2019)

We provide a direct experimental test of the Stokes-Einstein relation as a special case of the fluctuation-dissipation theorem (FDT) in an aging colloidal glass. The use of combined active and passive microrheology allows us to independently measure both the correlation and response functions in this non-equilibrium situation. Contrary to previous reports, we find no deviations from the FDT over several decades in frequency (1 Hz-10 kHz) and for all aging times. In addition, we find two distinct viscoelastic contributions in the aging glass, including a nearly elastic response at low frequencies that grows during aging. This is the clearest change in material properties of the system with aging.

PACS numbers:

Developing a statistical mechanical description of non-equilibrium systems such as glasses still remains an important challenge in physics. One of the most interesting recent developments along these lines is the proposal to generalize the fluctuation dissipation theorem (FDT) to non-equilibrium situations [1]. The FDT relates the response of a system to a weak external perturbation to the relaxation of the spontaneous fluctuations in equilibrium. More precisely, it states that the response function is proportional to the power spectral density of thermal fluctuations, with a prefactor given by the temperature of the system. This suggests a generalization for systems out of equilibrium, in which the (non-equilibrium) fluctuations are related to the response via a time-scale-dependent *effective temperature*. While this topic has been the subject of many theoretical studies and simulations [2], the experimental support for a meaningful effective temperature is unclear. There have been few experiments [3, 4, 5] and the usefulness of the extension of the FDT to non-equilibrium situation is still a matter of controversy.

Here, we introduce a combination of both *active* and *passive* (fluctuation-based) microrheology techniques [6, 7, 8] that provide a way to *directly* test the applicability of the FDT. We examine the validity of FDT in the form of the Stokes-Einstein relation (SER) in a colloidal glass, the synthetic clay Laponite [9, 10, 11]. For this system conflicting results have been reported previously [4, 5], that may in part be due to the use of a limited experimental window in both frequency and aging time. Here, we perform measurements over a wide range of frequencies and aging times. Contrary to previous reports, we find no violation of the FDT and thus no support for an effective temperature different from the bath temperature.

In addition, these measurements provide a new insight

into the physics of the aging process. The microrheology done during the aging process allows us to explore the evolution of viscoelastic properties of the glass over a wider frequency range than hitherto explored, spanning nearly 6 decades in frequency. The measurements reveal the existence of two distinct contributions to the viscoelasticity of the system: (i) a high-frequency viscoelastic response in which the shear modulus increases rapidly with frequency; and (ii) a predominantly elastic (weakly frequency-dependent) response at lower frequencies, which becomes increasingly important as the system ages.

The Stokes-Einstein relation (SER) relates the diffusion of the particle (*i.e.*, position fluctuations) to its mobility (response). The generalization for a viscoelastic system in Fourier space relates the power spectral density (PSD) of position fluctuations to the imaginary part of the complex response function $\alpha''(\omega)$ [7]:

$$\langle |x(\omega)|^2 \rangle = \frac{2k_B T}{\omega} \alpha''(\omega) \quad (1)$$

Here, $\langle |x(\omega)|^2 \rangle$ denotes the Fourier transform of $\langle x(t)x(0) \rangle$. In a non-equilibrium system the application of FDT suggests the introduction of an effective temperature in which T is replaced by $T_{\text{eff}}(\omega)$ in Eq. (1).

To investigate the aging of colloidal glass, we use microrheology, in which the motion of a probe particle within a viscoelastic sample is studied using optical tweezers. A custom-built inverted microscope [7], equipped with two overlapping optical tweezers formed by two independent lasers (wavelengths $\lambda = 830$ nm and 1064 nm focused to diffraction-limited spots. The latter drives the oscillations of the trapped particle with an Acousto-Optical Deflector, allowing us to measure the (active) response to a driving force. The (x, y) position of the particle is determined by a quadrant photo diode [12]




FIG. 1: The displacements power spectral densities (PSD) as a function of frequency of $1.16 \mu\text{m}$ silica probe particles in a 2.8% Laponite solution in water with increasing age after preparing the sample. Fluctuations were recorded for 45 seconds with the 830 nm laser focus and results averaged in x-y directions for 1 bead several times. Aging times are given in the legend. The filled squares show the PSD of a bead in pure water for comparison. The filled squares show the PSD of a bead in pure water for comparison. An acoustic noise signal around $f \approx 200\text{Hz}$ is cut out from the curve at the latest stage of aging where the displacement signal was the lowest. All experiments were done at 21°C .

with a spatial resolution of $\sim 0.1 \text{ nm}$. The output signal from the photodiode is fed into a lock-in amplifier that measures the amplitude and phase of the response function. For the passive measurements the shutter in front of the driving laser is closed and the spontaneous fluctuations of the particle position are recorded for a minimum time of 45 s. From the displacement time series, we calculate the displacement power spectral density by Fast Fourier Transform [7]. Comparing the response function from the active microrheology with the fluctuation spectra, we can directly check the validity of the FDT during the aging of the system, as well as resolve the frequency-dependent viscoelastic properties during the aging of the glass.

The colloidal glass under study is a suspension of Laponite XLG in ultra pure water. After mixing the colloidal particles with the water, the system spontaneously evolves from an initially liquid and ergodic state to a non-ergodic glassy state that exhibits elastic behavior [10]. For a particle concentration of 2.8 wt%, the rate of aging is slow enough, on the one hand, that no significant structural and dynamic changes occur during each individual active and passive microrheology measurement lasting at most 2 min. On the other hand, the system evolves fast enough to allow us to follow the evolution from ‘liquid’ to ‘solid’ (the glass no longer flows when the sample cell

is tilted) within about 8 hours. The dispersions are filtered (Millipore Millex AA $0.8 \mu\text{m}$ filter) to obtain a reproducible initial state [9]. This defines the zero of aging time $t_a = 0$. Immediately after filtration, a small fraction ($< 10^{-4} \text{ vol\%}$) of silica probe beads (diameter $1.16 \mu\text{m} \pm 5\%$) are mixed with the Laponite dispersion. The solution is then introduced into a sample chamber of about $50 \mu\text{l}$ volume, consisting of a coverslip and a microscope slide separated by a spacer of thickness $70 \mu\text{m}$. This is sealed with vacuum grease to avoid evaporation of sample. We then trap a silica bead and perform the experiments.

Figure 1 shows the (passive) displacement PSD for different aging times. It is evident that the particle motion progressively slows down with increasing aging time, reflecting the increase in viscosity of the system. Qualitatively two regimes of aging are seen: for $t_a < 200 \text{ min}$ the PSD can be described by a single power law. At longer aging times two distinct slopes appear in the log-log plot (Fig. 1).

To check the validity of the Stokes-Einstein relation in our aging system, we also measure the (active) response, again as a function of aging time and for oscillation frequencies of $f = 1.2, 10.8, 116, 1035$ and 12000 Hz . To directly compare the (passive) fluctuations with the (active) response, we convert fluctuation PSDs to response functions assuming that the FDT is valid and compare both the real and imaginary parts of the response functions. We obtain α'' directly from the measured PSD (Eq. (1)). Assuming that FDT holds, the real part can be calculated using a Kramers-Kronig (principal-value) integral $\alpha'(\omega) = \frac{2}{\pi} P \int_0^\infty \frac{\xi \alpha''(\xi)}{\xi^2 + \omega^2} d\xi$. The cutoff error due to a finite range of frequencies sets an upper limit to $\alpha'(\omega)$ about a decade lower than that of $\alpha''(\omega)$. Fig. 2 depicts the real and imaginary parts obtained from the active and passive methods at an early ($t_a = 100 \text{ min}$) and a late stage of aging ($t_a = 300 \text{ min}$). We see that the results are identical to within the experimental accuracy, showing that there are no deviations from the FDT in this system over the range of frequencies and aging times probed in our experiments. Note that the small deviations between the respective $\alpha'(\omega)$ values at high frequencies are likely due to cut-off errors in the Kramers-Kronig integrals because the drawn lines, obtained by extrapolating the measured power law of α'' to infinity, show significantly better agreement. Since we have directly compared both the real and imaginary parts of the response functions, this represents a stronger test of the FDT than previous measurements [5] and demonstrates that the FDT holds in this non-equilibrium system.

In Fig. 3, we plot the resulting $T_{\text{eff}}/T_{\text{bath}} = \alpha''_{\text{passive}}/\alpha''_{\text{active}}$ as a function of aging time for several different frequencies. This figure confirms again that to within the experimental uncertainty the FDT holds: the measured effective temperature does not differ from bath




FIG. 2: Comparison of active and passive results: Real $\alpha'(\omega)$ and imaginary $\alpha''(\omega)$ parts of the frequency-dependent response functions at $t_a = 100$ and 300 min obtained from active (solid symbols) and passive (open symbols) microrheology using identical $1.16\ \mu\text{m}$ diameter silica beads in one and the same aging sample. Imaginary parts of the response functions are obtained directly and real parts are calculated with a Kramers-Kronig integral from the PSDs for the passive experiments. Both parts were measured directly in the active experiments. At early stages of aging the data can be described with one power law., while at later stages, superposition of two power laws are needed to describe the whole frequency range. The amplitude of oscillation for the active experiments was $77\ \text{nm}$.

temperature. The errors for the lowest frequency ($1.2\ \text{Hz}$) are relatively large and there appears to be a trend towards an increased ratio $T_{\text{eff}}/T_{\text{bath}}$ with increasing aging time. This effect, however, is most certainly due to $1/f$ noise from laser pointing fluctuations. This is directly visible in the passive data at low frequencies: when the fluctuations become small the error becomes large. In addition, for the active measurements the errors at low frequencies become large when the elastic modulus of the material becomes large. We conclude that the measurements show that the FDT in the form of the SER is valid for all frequencies probed here, and can be used for all the stages of aging in this system.

These measurements also demonstrate, for the first time, the applicability of fluctuation-based (passive) microrheology in a non-equilibrium glassy system. The method allows to obtain the viscoelastic properties over a very wide frequency range; classical (macroscopic) rheology is limited to frequencies up to about $10\ \text{Hz}$ [10]. We therefore now turn to an analysis of the rheological properties of the system.

In Fig. 1 we observe a gradual decrease of the PSD for higher frequencies and a more rapid change at lower fre-




FIG. 3: The ratio of (apparent) response derived from fluctuations via FDT and actively measured response $T_{\text{eff}}/T_{\text{bath}}$ as a function of aging time for different frequencies ($f = 1.2, 10.8, 116, 1035$ and $12000\ \text{Hz}$) from bottom to the top in Laponite colloidal glass of $2.8\ \text{wt}\%$ concentration. This ratio can also be seen as a measure of effective temperature.

quencies. The response function is directly proportional to the PSD which in turn is related to the inverse of complex shear modulus $G^* = G' + iG''$. With increasing aging time, as the sample becomes more viscous and solid-like, one would expect that both the elastic modulus G' and the viscous modulus G'' increase. This is consistent with the decrease of the PSD and consequently of the response function. As was mentioned above, at late stages of aging two distinct slopes appear in the PSD (Fig. 1). This suggests the existence of two distinct contributions to the viscoelasticity during aging. We obtain excellent fits to the data assuming a simple addition of two power law contributions to the complex shear modulus (only a single power-law contribution at the early stages of aging). Thus, the complex compliance is fitted with:

$$\alpha^*(\omega) = \frac{1}{6\pi R G^*(\omega)} = \frac{1}{C_1(i\omega)^a + C_2(i\omega)^b} \quad (2)$$

The fit of the imaginary part of the response function from the passive measurements with the imaginary part of the above functional form is shown by the drawn lines in Fig. 2. To demonstrate the quality of this model for describing the data, we also plot the real part with the fitting parameters obtained from the imaginary part. The agreement is remarkable, especially with the active data at high frequencies. Other functional forms such as notably the much-used Havriliak-Negami model [13] cannot describe the data over the whole frequency range.




FIG. 4: For $t_a < 200$ min, the complex response function can be described by single power law of the form $\frac{1}{(i\omega)^\alpha}$. After about 200 min, a slowly relaxing contribution emerges leading to a response function of the form in Eq. (2). Note that data around 200 min could not be fitted with either a single or with two power laws and are consequently not shown.

Fig. 4 depicts the evolution of the fitting parameters, *i.e.*, the exponents of power laws and the weight factors for the contribution of the two viscoelastic contributions as a function of aging time. The exponent and amplitude of the one of the components do not change with aging time while the amplitude of the other one grows appreciably when the aging time becomes larger than about 250 min.

These results demonstrate the existence of two distinct contributions in the viscoelasticity of the system. In addition to a strongly frequency-dependent viscoelastic response at high frequencies, we also observe the slow development of a more elastic (weakly frequency-dependent) response during the aging. In fact, this appears to be the main characteristic of the aging in this system. Similar behavior has been reported previously for gel formation in polymeric systems, and can be attributed to the growth of a tenuous network-like structure, in addition to the more fluid-like response that is always present. We note, however, that although our system also becomes non-ergodic there is no evidence for spatial inhomogeneity, neither in our system, nor in glassy systems in general. Recently, it has been suggested that glassy systems may be dynamically heterogeneous [14], and that one should look into the correlation of spatial and temporal dynamics in order to detect the heterogeneity. In view of the existence of the two distinct viscoelastic contributions in our system, that show some similarity to the modes observed for very heterogeneous polymeric systems, the homogeneity of the spatial and temporal dynamics merit

further study, which is planned.

In summary, our measurements of both active and passive microrheology, show good quantitative agreement between the response function and the spontaneous thermal fluctuations, implying that we observe no violation of the FDT in this non-equilibrium system. Equivalently, we find an effective temperature that does not differ from the system temperature. It is important to note that these measurements provide a much more direct test of the FDT than prior experiments, since we directly measure the response and the *corresponding* fluctuations over the same and very large range of frequencies.

Acknowledgments

The research has been supported by the Foundation for Fundamental Research on Matter (FOM), which is financially supported by Netherlands Organization for Scientific Research (NWO). LPS de l'ENS is UMR8550 of the CNRS, associated with the universities Paris 6 and 7. We would like to thank N. Israeloff and H. Tanaka for helpful discussions.

-
- [1] L.F. Cugliandolo, J. Kurchan, and L. Peliti Phys. Rev. E **55**, 3898 (1997).
 - [2] W. Kob, J.L. Barrat, Phys. Rev. Lett. **78**, 4581 (1997); G. Parisi, Phys. Rev. Lett. **79**, 3660 (1997).
 - [3] T. Grigera, N. Israeloff, Phys. Rev. Lett. **83**, 5083 (1999); D. Hérisson and M. Ocio, Phys. Rev. Lett. **88**, 257202 (2002); D. Bonn, W.K. Kegel, J. Chem. Phys. **118**, 2005 (2003); C.M. Song, P. Wang, H. A. Makse, Proc. Nat. Acad. Sci. **102**, 2299 (2005).
 - [4] L. Bellon, S. Ciliberto, C. Laroche, Europhys. Lett. **53**, 511 (2001); L. Bellon, S. Ciliberto, Physica D **168-169**, 325 (2002); L. Buisson, L. Bellon, and S. Ciliberto, J. Phys. Cond. Matt. **15**, S1163 (2003).
 - [5] B. Abou and F. Gallet, Phys. Rev. Lett. **93**, 160603 (2004).
 - [6] T.G. Mason, and D.A. Weitz, Phys. Rev. Lett. **74**, 1250 (1995).
 - [7] F. Gittes, B. Schurr, B. and Olmsted, P.D. F.C. MacKintosh, and C.F. Schmidt, Phys. Rev. Lett. **79**, 3286 (1997); B. Schnurr, F. Gittes, F.C. MacKintosh, C.F. Schmidt, Macromolecules **30**, 7781 (1997).
 - [8] L.A. Hough and H.D. Ou-Yang, Phys. Rev. **E65**, 021906 (2002); A.W.C. Lau, B.D. Hoffman, A. Davies, J.C. Crocker, and T.C. Lubensky, Phys. Rev. Lett. **91**, 198101 (2003).
 - [9] D. Bonn, H. Kellay, H. Tanaka, G.H. Wegdam and J. Meunier, Langmuir **15**, 7534-7536 (1999), D. Bonn, H. Tanaka, H. Kellay, G.H. Wegdam and J. Meunier, Europhys. Lett. **45**, 52 (1998).
 - [10] D. Bonn, P. Coussot, H.T. Huynh, F. Bertrand and G. Debregeas, Europhys. Lett. **59**, 786 (2002).
 - [11] M. Bellour, A. Knaebel, J.L. Harden, F. Lequeux and J.-P. Munch, Phys. Rev. **E67**, 031405 (2003), S. Kaloun, R. Skouri, M. Skouri, J. P. Munch and F. Schosseler Phys. Rev. **E72**, 011403 (2005).
 - [12] F. Gittes and C. F. Schmidt, Optics Lett. **23**, 7 (1998).
 - [13] S. Havriliak and S. Negami, Polymer **8** (4), 161 (1967)

- [14] M. Merolle, J. P. Garrahan and D. Chandler, Proc. Nat. Acad. Sci. **102** (31), 10837 (2005); P. Mayer, H. Bissig, L. Berthier, L. Cipelletti, J.P. Garrahan, P. Sollich, and V. Trappe, Phy. Rev. Lett. **93**, 115701 (2004)