

Wavefunctions and counting formulas for quasiholes of clustered quantum Hall states on a sphere

N. Read

Department of Physics, Yale University, P.O. Box 208120, New Haven, CT 06520-8120

(May 5, 2006)

The quasiholes of the Read-Rezayi clustered quantum Hall states are considered, for any number of particles and quasiholes on a sphere, and for any degree k of clustering. A set of trial wavefunctions, that are zero-energy eigenstates of a $k+1$ -body interaction, and so are symmetric polynomials that vanish when any $k+1$ particle coordinates are equal, is obtained explicitly and proved to be both complete and linearly independent. Formulas for the number of states are obtained, without the use of (but in agreement with) conformal field theory, and extended to give the number of states for each angular momentum. An interesting recursive structure emerges in the states that relates those for k to those for $k-1$. It is pointed out that the same numbers of zero-energy states can be proved to occur in certain one-dimensional models that have recently been obtained as limits of the two-dimensional $k+1$ -body interaction Hamiltonians, using results from the combinatorial literature.

Non-Abelian quantum Hall states have been a subject of great interest for some time [1]. Moore and the author showed that non-Abelian statistics, in which adiabatic exchange of well-separated quasiparticles produces a matrix operation on a space of degenerate states, was a possibility in a condensed matter system, and gave an example, now often termed the Moore-Read (MR) state. They related the deep structure underlying the statistics and other aspects of the quasiparticles of these phases to conformal field theory, and to what are now called modular tensor categories. Many trial wavefunctions of quantum Hall states can be related to conformal blocks (i.e. chiral parts of correlation functions) in the corresponding conformal field theory. A further sequence of examples was constructed by Rezayi and the author (RR) [2]. These are parametrized by two integers, $k \geq 1$ and $M \geq 0$; the filling factor for each is $\nu = k/(Mk+2)$, and they describe bosons for M even, fermions for M odd (the case $k=1$ is the Laughlin state, while $k=2$ is the MR state). For $M=0$ or 1, trial ground state and quasihole states can be obtained as zero-energy eigenstates of a Hamiltonian that consists of a purely $k+1$ -body interaction, acting within the lowest Landau level. (For $M > 0$, the trial wavefunctions are those for $M=0$ multiplied by the factor $\prod_{i < j} (z_i - z_j)^M$, where $z_i = x_i + iy_i$ is the complex coordinate representing position (x_i, y_i) of particle i in two dimensions. These trial wavefunctions are zero-energy eigenstates for a Hamiltonian that contains two-body as well as $k+1$ body interactions [2].) For $M=0$, the wavefunctions are closely related to conformal blocks of $SU(2)$ level k current algebra conformal field theories [2]. While the explicit trial ground state wavefunctions were obtained, for $k \geq 3$ the trial wavefunctions for quasiholes were not, except for some of those for small numbers of quasiholes [2,3]. The number of degenerate zero-energy states for fixed positions of the quasiholes is basic to the theory of non-Abelian statistics, and for the MR and RR states these degeneracies have been studied numerically

and using conformal field theory methods [4,5,2,6-8].

The MR state is now believed theoretically to describe the phase observed at filling factor $\nu = 5/2$ in electron systems [9], while there is evidence that the RR $k=3$, $M=1$ state occurs at $\nu = 12/5$ [2]. For bosons, there is numerical evidence that the sequence of RR states is relevant to rotating cold, trapped, bosonic atoms [10,11]. Interest in non-Abelian statistics is presently being fueled by the possibility of topological quantum computation [12].

In this paper, we return to the question of finding and counting the zero-energy states of the $k+1$ -body interaction for bosons, for general k . We work in the lowest Landau level on the sphere, with N_ϕ quanta of magnetic flux piercing it [13]. The approach is a generalization of one used to prove linear independence of a proposed set of functions for the $k=2$ case in Ref. [5], and is very close to work [14] on the same counting problem for the limit $N_\phi \rightarrow \infty$, which has been reviewed, extended, and applied in Ref. [15]. (The authors of the first paper in Ref. [14] discovered the relation of the symmetric functions that vanish when $k+1$ coordinates coincide to conformal blocks of $SU(2)$ current algebra at level k , which was discussed independently in Ref. [2]. From the quantum Hall point of view, the results on the $N_\phi \rightarrow \infty$ case describe edge excitations.) Our main results are: (i) a complete and linearly-independent set of explicit wavefunctions for zero-energy states of the $k+1$ -body interaction for any number of particles N and any number of flux quanta N_ϕ , or alternatively, any number n of quasiholes, as defined by

$$N_\phi = \frac{2}{k}N + M(N-1) + \frac{n}{k} - 2; \quad (1)$$

(ii) explicit generating functions that encode the number of zero-energy states for any N , n , decomposed according to orbital angular momentum L_z (and hence L also); (iii) an interesting recursive structure emerges, that relates the quasihole problem for k to the similar problem

for $k-1$ at a reduced number of particles and flux quanta, and might be useful in making further progress in understanding the quasihole states. The arguments are short and self-contained, and make no use of conformal field theory. In an appendix, we also consider quasihole wavefunctions of a natural form that generalizes those in Ref. [3], but for these our results are less conclusive.

We want to find the complete set of zero-energy states for the $k+1$ -body Hamiltonians in Ref. [2]. The cases with $M > 0$ can be found by multiplying those for $M = 0$ by the Laughlin-Jastrow factor $\prod_{i < j} (z_i - z_j)$ to the power M , so we can focus on the $M = 0$ case (hence, on bosons). Then the problem reduces to the following: find all the symmetric polynomials in N complex variables z_i that vanish when any $k+1$ of the z_i are equal. For the present, working on the sphere means that the polynomial is of degree at most N_ϕ in each z_i [13]. We will denote the vector space of symmetric polynomials with these properties by $V_{N, N_\phi, k}$ and a typical member of this space by $\tilde{\Psi}_{N, N_\phi, k}$ (the polynomials $\tilde{\Psi}$ can be related to wavefunctions Ψ on the sphere or plane as explained in Ref. [5]).

Consider a function $\tilde{\Psi}_{N, N_\phi, k}$. While it vanishes whenever $k+1$ particles coincide, it may not vanish when only k coincide. Let z_1, z_2, \dots, z_k tend to the same value, which we will denote by Z_1 . If it vanishes in this limit, then (since it is totally symmetric) it vanishes when any k coordinates coincide, and belongs to $V_{N, N_\phi, k-1}$. If

it does not vanish, then its value (corresponding to the ‘‘residue’’ in Ref. [5]) is a function of Z_1 and of z_{k+1}, \dots, z_N , and is symmetric in the latter. In this residue we may let another set of k coordinates, say z_{k+1}, \dots, z_{2k} coincide at Z_2 . If the function vanishes in this second limit, then the residue consists of a function of Z_1 times a member of $V_{N-k, N_\phi, k-1}$. If it does not vanish in the second limit, we obtain a second residue, to which the procedure can be applied again.

In this way the space of functions is divided into a nested sequence of subspaces within subspaces (called a ‘‘filtration’’ in Ref. [15]). Each such subspace contains all the wavefunctions that have a non-vanishing residue, which is a function of Z_1, Z_2, Z_3, \dots , (each of which are the locations of $(N-F)/k$ ‘‘clusters’’ of k particles), and of the remaining F coordinates z_{N-F+1}, \dots, z_N of the F ‘‘unclustered’’ particles; these residues vanish if any k of the latter set coincide. Each subspace also contains functions with vanishing residues, which are also members of a subspace with fewer clusters. Thus, the $(N-F)/k$ th subspace [$(N-F)/k = 0, 1, \dots$] can be defined as the space of functions that vanish identically if $(N-F)/k+1$ clusters of k particles each coincide at $Z_1, \dots, Z_{(N-F)/k+1}$, for all values of these coordinates and of the remaining z 's. We will show that the non-vanishing residues of functions in this subspace must take the following form:

$$\tilde{\Psi}_{N, N_\phi, k} \rightarrow \prod_{p < p'} (Z_p - Z_{p'})^{2k} \prod_{p'', l} (Z_{p''} - w_l) \prod_{p''', m} (Z_{p'''} - z_{N-F+m})^2 \tilde{\Psi}_{F, N_\phi - 2(N-F)/k, k-1}. \quad (2)$$

The notation and the justification for each of the factors is as follows. The first product is over pairs of cluster coordinates Z_p , labeled by $p, p', \dots, = 1, 2, \dots, (N-F)/k$ (the number of clusters). The function vanishes when any $k+1$ particles coincide, so it must vanish when any two clusters coincide, and since it is symmetric under exchanges of clusters, it must contain a factor $(Z_p - Z_{p'})^2$ for each pair of clusters. In fact, because the original wavefunctions are symmetric functions for particles on the sphere, it must vanish as the $2k$ th power; the proof of this is given in an appendix to this paper. The last factor $\tilde{\Psi}_{F, N_\phi - 2(N-F)/k, k-1}$ is a function only of the F ‘‘unclustered’’ coordinates, so is independent of the cluster coordinates. By assumption, it vanishes if any k of the remaining F coordinates coincide, so can be any function in $V_{F, N_\phi - 2(N-F)/k, k-1}$. The residue must also vanish if any of the unclustered particles approaches a cluster, so it must contain $Z_p - z_{N-F+m}$ for the p th cluster and the m th unclustered particle ($m = 1, 2, \dots, F$). In fact, it must vanish quadratically, which again is proved in the Appendix. This form implies that the maximum degree in the unclustered coordinates in the remaining factor is $N_\phi - (N-F)/k$, as recorded in the second subscript on

$\tilde{\Psi}_{F, N_\phi - 2(N-F)/k, k-1}$. Finally, the residue can contain a further factor of a symmetric polynomial in the cluster coordinates. As the degree in any of the cluster coordinates is at most kN_ϕ , it works out that the maximum degree in each cluster coordinate in this symmetric polynomial factor is n . Each quasihole effectively contains $1/k$ of a flux quantum or vortex [2]. In general, n need not be divisible by k (in Ref. [2], only the special cases N and n divisible by k were considered). For N_ϕ to be an integer, we require that $2N + n$ be divisible by k (hence for k even, n must be even), and $\dim V_{N, N_\phi, k}$ (the dimension of $V_{N, N_\phi, k}$) is nonzero only when $n \geq 0$. The symmetric polynomials in the cluster coordinates of this degree can be expressed using the product $\prod_{p'', l} (Z_{p''} - w_l)$ as a generating function (by expanding it in symmetric polynomials in the variables w_l). These are then ‘‘coherent states’’, in which the coordinates w_l can be interpreted as the positions of the quasiholes. The function vanishes when any k particles approach a w_l , as it should [16], and this can be taken as a definition of the quasihole coordinates for these quasiholes, which have charge $-1/2$ (in units of particle number).

Let us suppose that the problem has already been

solved for values $k' = k - 1$. Then we have a set of linearly independent functions $\tilde{\Psi}_{F, N_\phi - 2(N-F)/k, k-1}$ that span $V_{F, N_\phi - 2(N-F)/k, k-1}$. Then by expanding in the quasihole coordinates w_l , we have a set of non-vanishing residues that are functions of the cluster coordinates and the unclustered particle coordinates, and clearly are linearly independent. This implies that if for each of these residues, there is a wavefunction from which it arose by taking limits, then these wavefunctions (which are in $V_{N, N_\phi, k}$) are linearly independent. The existence of a wavefunction that reduces to each non-vanishing residue in the limit will be shown in a moment.

Completeness of the set of functions can also be proved by induction. As the form of the non-vanishing residues has been fixed, we must show that there is a one-one mapping from the wavefunctions to the non-vanishing residues. This is done by induction on the number of clusters. For the residues with no clusters ($F = N$), the residue reduces to $\tilde{\Psi}_{N, N_\phi, k-1}$, for which a complete set is assumed to have already been found. Then for the residues with one cluster ($F = N - k$), if there are two wavefunctions with the same residue, then the difference of these functions is a wavefunction with no clusters (since it vanishes when k particles coincide), and these are also in the space $V_{N, N_\phi, k}$. This argument can be repeated, and so by induction we find that there is (up to addition of a wavefunction for fewer clusters) at most one wavefunction for each non-zero residue, that is (assuming existence) the dimension of the space of non-vanishing residues and that of the space of wavefunctions are the same for each N, N_ϕ , and k .

To prove the existence of a zero-energy wavefunction

that tends to each non-vanishing residue in the limit, we construct them. We use the following labeling for the particles: The particle number N is partitioned into $m_\alpha \geq 0$ clusters of size $\alpha = 1, \dots, k$, that is $N = \sum_\alpha \alpha m_\alpha$ (there will be a set of wavefunctions for each such partition). One particle is assigned to each box in a Ferrers-Young diagram in which there are at most k columns, and which consists of m_α rows of length α for each α , arranged as usual with the row lengths weakly decreasing as one goes down the diagram. Then (following Ref. [15]) the particle coordinates will be written $z_{ij}^{(\alpha)}$, where i, j label rows and columns, respectively (as for a matrix, i increases down the columns, and j increases to the right along the rows) of the rectangular block consisting of all rows of length $\alpha = 1, \dots, k$. Thus for each $\alpha, j = 1, \dots, \alpha$, and $i = 1, \dots, m_\alpha$. The rows of the diagram are ordered with (α, i) above (β, i') , written $(\alpha, i) > (\beta, i')$, if either $\alpha > \beta$, or $\alpha = \beta$ and $i < i'$ (note the direction of the last inequality). The wavefunctions will be written down in a form that deals with all levels of the recursion at once. It will turn out that $(N - F)/k = m_k$, while the remaining m_α represent, similarly, the numbers of clusters of sizes α smaller than k . Similarly, the wavefunctions involve not only quasihole coordinates $w_l^{(k)}$ ($l = 1, \dots, n_k = n$), which can be identified with the w_l in the residues, but also further coordinates $w_l^{(\alpha)}$ for all $\alpha = 1, \dots, k - 1$, which play a corresponding role in the parametrization of the wavefunctions for smaller k that appear in the recursion. Here for each α the range is $l = 1, \dots, n_\alpha$, with $n_1 \leq n_2 \leq \dots \leq n_k$.

The wavefunctions are

$$\tilde{\Psi}_{N, N_\phi, k}(z_1, \dots, z_N; \{w_l^{(1)}\}, \dots, \{w_l^{(k)}\}) = \mathcal{S}_z \left\{ \prod_{(\alpha, i) > (\beta, i')} \prod_{j=1, \dots, \beta} (z_{i,j}^{(\alpha)} - z_{i',j}^{(\beta)}) (z_{i,j+1}^{(\alpha)} - z_{i',j}^{(\beta)}) \cdot \prod_\alpha \prod_{i,j} \prod_l (z_{ij}^{(\alpha)} - w_l^{(\alpha)}) \right\}. \quad (3)$$

In this expression, \mathcal{S}_z is the symmetrizer over all the N particles, and $z_{i, \alpha+1}^{(\alpha)} = z_{i,1}^{(\alpha)}$. In each product $\prod_l^{(\alpha, j)}$, l runs over

$$l = n_{j-1} + (1 - \delta_{j,1}) \sum_{\beta: j-1 \leq \beta < \alpha} m_\beta + 1, \dots, n_j + \sum_{\beta: j-1 < \beta < \alpha} m_\beta \quad (4)$$

(where $n_0 = 0$). To ensure that each function is of the same degree $N_\phi = (2N + n)/k - 2$ in all coordinates z_i , we require

$$2 \sum_{\beta: \beta \geq \alpha} m_\beta + n_\alpha - n_{\alpha-1} = (2N + n)/k \quad (5)$$

for all $\alpha = 1, \dots, k$. Also, for each α , the wavefunctions are of the same degree m_α in $w_l^{(\alpha)}$ for all $l = 1, \dots, n_\alpha$.

The structure of the $z - z'$ factors inside the symmetrizer is based on that in Ref. [15], and for the factors

that connect two rows (α, i) , (β, i') with $\alpha = \beta$ is the same as in the ground states (i.e. the case $m_k = N/k$, $m_\alpha = 0$ for $\alpha < k$) in Ref. [2]. These particle-particle factors are of two types: one type connects two particles (and vanishes when their coordinates coincide) whenever they are in the same column in the diagram. The other type connects two particles in different rows only if the one in the higher row is one column to the right of the lower one; when the rows are of the same length, this is interpreted cyclically along the rows. The following ar-

guments are a version of the arguments for existence of a wavefunction mapping to each residue in Ref. [15], and generalize an argument for the ground state wavefunctions in Ref. [2].

The wavefunctions in eq. (3) vanish when any $k + 1$ coordinates are equal, because in each term inside the symmetrizer, at least two must be in the same column of the diagram (i.e. have the same value of j). They also vanish if k particles coincide at Z_1 say, unless the particles lie in the same row (which must be of length k). First, these particles must be in distinct columns for the wavefunction to have a possibility of not vanishing, and hence one must be in column k , and so in the first m_k rows. Then in order for the function not to vanish as the particles come together, the one in column $k - 1$ must be in the same or a higher row, and so on, until in the first column the particle will be in the same or a higher row as that in column k . If the row is higher, the wavefunction will vanish because of the cyclic connection of the first and last columns. Hence all k must lie in the same row. Repeating this argument as another k particles tend to Z_2 , and so on, one finds that in the clustered limit, when there are m_k clusters with k coinciding coordinates in each, the particles in each cluster came from the same row in each term that contributes a non-vanishing term to the residue. The residue vanishes if any k of the remaining coordinates coincide, and vanishes as $(Z_i - Z_j)^{2k}$ if any two clusters of k coincide. The wavefunctions also vanish if k particles coincide with a $w_l^{(k)}$. Again, the k particles must be in the same row, and then exactly one of them appears in a factor $(z - w_l^{(k)})$, so the wavefunction vanishes, and the residue does likewise. Finally, the residue vanishes quadratically as $(Z_i - z')^2$ for any of the remaining unclustered coordinates z' , as each of them is connected to exactly two particles in each row of length k . This establishes that the residue is of the form in eq. (2), with $\tilde{\Psi}_{F, N_\phi, k'}$ being a function in the remaining z 's and w 's of the same form as in eq. (3), with parameters as indicated. Then arguing by induction on k establishes that there is a wavefunction for each non-vanishing residue, as claimed.

Taking all the arguments together, we can conclude that the wavefunctions in eq. (3), though written as over-complete sets of ‘‘coherent states’’ using auxiliary coordinates $w_l^{(\alpha)}$, span a complete set of linearly-independent zero-energy wavefunctions. Because the residues are invariant under permutations of the $w^{(\alpha)}$'s among themselves (for fixed α), no states are lost if the wavefunctions are symmetrized over the $w^{(\alpha)}$'s also, by applying $\prod_\alpha \mathcal{S}_{w^{(\alpha)}}$. Then the functions can be expanded as linear combinations of polynomials in the z 's times products over α of symmetric polynomials in the $w_l^{(\alpha)}$'s for each α , and the polynomials in z for each distinct product of symmetric polynomials in the w 's form a complete and linearly independent set of wavefunctions.

For $k = 1$, the wavefunctions are the Laughlin quasi-hole wavefunctions. For $k = 2$ they are the same complete set of linearly-independent functions found in Ref. [5] (and for N even and $n = 2$ quasiholes, in Ref. [1]). In the latter, the unclustered (unpaired) particles were viewed as fermions occupying zero modes, but we now see that they can also be viewed as Laughlin quasihole wavefunctions for $\nu = 1/2$ involving F particles and n_1 quasiholes. For general k , they agree with Ref. [2] for $n = k$, N divisible by k .

We can now find the ‘‘number of states’’ $\#_{N, N_\phi, k} = \dim V_{N, N_\phi, k}$, by counting the non-vanishing residues (or by counting the wavefunctions directly). In these first results, we use only the recursive structure. We expand the symmetric polynomials in the residues in eq. (2) in the w_l 's, so as to obtain linearly-independent residues; the number of linearly-independent such functions is a binomial coefficient [5], and this then yields

$$\#_{N, N_\phi, k} = \sum_{F: F \equiv N \pmod{k}} \binom{\frac{N-F}{k} + n}{n} \#_{F, N'_\phi, k'}, \quad (6)$$

where $N'_\phi = N_\phi - 2(N - F)/k = (2F + n)/k - 2$, and $k' = k - 1$. This has the general structure found for $k = 2$ in Ref. [5]. For general k , the structure was conjectured in Ref. [2], and results were obtained in [6,7] (although the form of the ‘‘spatial degeneracy’’ factors, i.e. the binomial coefficients in this formula, seems never to have been derived generally). In the literature, this form is usually written with the notation [6,7]

$$\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_k = \#_{F, N'_\phi, k'}, \quad (7)$$

when $n + 2F$ is divisible by k . Remarkably, we have found that these coefficients, which represent the ‘‘internal’’ degeneracy of the quasiholes, are themselves related to a problem of counting the full degeneracy of zero energy quasihole states, but at a different value of k . This now allows an inductive solution for these coefficients, recursively in k .

First, we can write a recursion relation for the coefficients:

$$\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_k = \sum_{F': F' \equiv F \pmod{k'}} \binom{\frac{F-F'}{k'} + n'}{n'} \left\{ \begin{matrix} n' \\ F' \end{matrix} \right\}_{k'}, \quad (8)$$

where again $k' = k - 1$, and

$$n' = (k'n - 2F)/k. \quad (9)$$

Next we mention that for $k = 1$, the residues above are just the wavefunctions, and the function from the problem for $k' = 0$ is simply unity. Those wavefunctions are the Laughlin quasihole functions. This implies that $\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_1 = \delta_{F,0}$. In this case, the statistics is Abelian,

and the only degeneracy is due to the different possible locations of the quasiholes. The recursion relation then yields for $k = 2$,

$$\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_2 = \binom{n/2}{F}, \quad (10)$$

as found in Ref. [5]. When these numbers, or their sum over F , are larger than 1, they describe the degeneracy of states for quasiholes even when their positions are fixed, a characteristic of a non-Abelian state.

When these values are substituted into the recursion relation for $k = 3$, the result can be immediately rewritten

$$\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_3 = \sum_{a,b:a+2b=F} \binom{\frac{n-F}{3}}{a} \binom{\frac{2n-(2a+b)}{3}}{b}, \quad (11)$$

which was found previously by Ardonne [7], by a much longer method using conformal field theory and results from Ref. [17].

The solution for arbitrary k can be found recursively, and parametrized in the same way as the wavefunctions, eq. (3), using the partitions of N and n . First, the total number of states can be written as

$$\#_{N,N_\phi,k} = \sum_{m_1,\dots,m_k:\sum_\alpha \alpha m_\alpha = N} \prod_\alpha \binom{m_\alpha + n_\alpha}{n_\alpha}. \quad (12)$$

Here the recursive structure is clear; notice that $m_k = (N - F)/k$, $n_{k-1} = n'$, $m_{k-1} = (F - F')/k'$, etc., so this agrees with preceding results. In practice we need the values of n_α for given N and n ; these are obtained by solving the conditions on the number of $w_l^{(\alpha)}$ at each level α in terms of the m_β 's:

$$n_\alpha = \alpha \left(\frac{2N + n}{k} - \sum_{\beta:\beta \geq \alpha} 2m_\beta \right) - \sum_{\beta < \alpha} 2\beta m_\beta; \quad (13)$$

for $\alpha = k$, this reduces to $n_k = n$. There is a similar form for the coefficients $\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_k$, as a sum over partitions of $F = N - km_k$, which agrees with the result of Ref. [7].

One further refinement is to count zero-energy eigenstates for each angular momentum value L_z on the sphere. Instead of $\dim V_{N,N_\phi,k}$, we can choose to calculate the trace of $q^{N N_\phi/2 - L_z}$, where q is an indeterminate, and L_z is the z -component of total angular momentum \mathbf{L} . We recall that for the single-particle wavefunctions, the function z^m contributes $N_\phi/2 - m$ to L_z , for $m = 0, \dots, N_\phi$ [13,5]. Thus (for L_z eigenstates) we have arranged that the exponent of q is simply the total degree of the wavefunction in the z 's. From the number of states of each L_z , one can extract the number of quasihole states of each value of \mathbf{L}^2 also. This information can be valuable in numerical studies.

Once again it is sufficient to count residues, this time of each degree. The total degree in the cluster coordinates Z_p , $p = 1, \dots, m_k$, is $km_k(m_k - 1)$ plus the total degree of the symmetric polynomial in the Z 's produced by expanding the factor $\prod_{p,l}(Z_p - w_l)$, plus the contributions of the factors that connect with the unclustered particles. The number of symmetric polynomials of degree d in m_k variables and of degree at most n_k in each variable is given by the coefficient of the q^d term in the (Gauss) q -binomial coefficient (a polynomial in q), defined by

$$\binom{m_k + n_k}{n_k}_q = \frac{(m_k + n_k)_q!}{(m_k)_q!(n_k)_q!}, \quad (14)$$

where the q -factorial $(n)_q! = (n)_q(n-1)_q \cdots (1)_q$, and the q -deformed integers are

$$(n)_q = 1 + q + q^2 + \cdots + q^{n-1} = \frac{1 - q^n}{1 - q}. \quad (15)$$

This combinatorial result can be understood from its own generating function, which is

$$\frac{1}{(1-x)(1-xq)\cdots(1-xq^{m_k})} = \sum_{n_k=0}^{\infty} \binom{m_k + n_k}{n_k}_q x^{n_k}; \quad (16)$$

this identity is Heine's q -binomial formula [18]. It is easily seen that the left-hand side of eq. (16) is the generating function for symmetric polynomials in m_k variables, of degree at most n_k in each variable: The symmetric polynomials in m_k variables Z_p are (freely) generated by the elementary symmetric polynomials $e_1 = \sum_p Z_p$, $e_2 = \sum_{p < p'} Z_p Z_{p'}$, \dots , $e_{m_k} = \prod_p Z_p$, which are of total degrees $1, \dots, m_k$, respectively, and of degree at most 1 in each variable. Each factor $(1 - xq^m)^{-1}$ gives a series that contains one term $x^r q^{mr}$ with coefficient 1, corresponding to each distinct possible factor e_m^r . Similar formulas apply to each level in the recursion, with α in place of k . In all these expressions, the limit $q \rightarrow 1$ reproduces earlier or well-known results, with ordinary binomials, factorials, etc., in place of their q -analogues.

Since the q -binomials count the effects of the factors containing $w_l^{(\alpha)}$'s, we now only require the total degree (summed over α) of the minimal degree polynomial in the Z 's of all levels, that is the term of highest possible degree in all w 's. For the given partition described by m_α , this minimal total degree is

$$\begin{aligned} \sum_\alpha \alpha m_\alpha (m_\alpha - 1) + \sum_{\alpha < \beta} 2\alpha m_\alpha m_\beta, \\ = \sum_{\alpha, \beta} m_\alpha m_\beta M_{\alpha\beta} - N, \end{aligned} \quad (17)$$

where $M_{\alpha\beta} = \min(\alpha, \beta)$ [14,15].

The result is that the trace over $V_{N,N_\phi,k}$ is

$$\text{tr } q^{N N_\phi / 2 - L_z} = q^{-N} \sum_{m_1, \dots, m_k; \sum_\alpha m_\alpha = N} q^{\sum_{\beta, \gamma} m_\beta m_\gamma M_{\beta\gamma}} \prod_\alpha \binom{m_\alpha + n_\alpha}{n_\alpha}_q. \quad (18)$$

This expression counts the zero-energy states on the finite-size sphere for each L_z ; furthermore, $\text{tr } q^{-L_z}$ for $M \geq 0$ is independent of M . As a test, this formula has been compared with some rows for $k > 2$ in the tables of numbers of states for each L that have appeared in Ref. [6] (for $k = 3$) and recently in Ref. [8], with perfect agreement [19].

The corresponding result that generalizes $\left\{ \begin{matrix} n \\ F \end{matrix} \right\}_k$ for $q \neq 1$ was given in Ref. [7] as the ‘‘truncated character’’ for parafermions (but of course does not describe the total orbital angular momentum of the states). Those expressions now emerge from analyzing polynomials in z ’s.

It is possible to take the limit $N_\phi \rightarrow \infty$ with m_α fixed. In this limit, all $n_\alpha \rightarrow \infty$, and by assuming that $|q| < 1$,

$$\binom{m_\alpha + n_\alpha}{n_\alpha}_q \rightarrow \prod_{m=1}^{m_\alpha} \frac{1}{(1 - q^m)}. \quad (19)$$

Then the resulting generating function for the number of zero-energy wavefunctions with no upper limit on the degree in each variable is identical with the results of Refs. [14,15]. These references also explain how to shift the degree in q so that the subsequent limit $N \rightarrow \infty$ can be taken for each term of fixed degree in q . For the simplest case, one divides by $q^{N^2/k-N}$, and the limit counts the number of excitations at finite change in angular momentum of the edge of a very large drop of particles, similar to the results of Ref. [20] for the paired states (the $k = 2$ case). In this limit the leading term in the generating function when N is divisible by k is q^0 and corresponds to the vacuum of the conformal field theory for the edge, as there are no quasipoles in the interior. More generally, one may also take the N_ϕ and $N \rightarrow \infty$ limits with some quasipoles held fixed at the center of the drop, and arrange that the powers of q in general are the conformal weights of the states on the edge. For up to k quasipoles at the center, this was done in Refs. [14,15], and produces the character of the chiral conformal field theory for each sector of edge states (there are $k + 1$ distinct sectors for the $M = 0$ case treated here, but this number does depend on M [2]).

To conclude, we now have a satisfactory understanding of the trial quasipole wavefunctions of the RR states. We expect that the approach can be generalized to obtain analogous results for the quasipoles of other quantum Hall phases that have trial states that are zero-energy eigenstates of short-range interactions, such as bilayer states, spin-polarized ground states of spin-1/2 particles, and the various spin-singlet clustered states for particles with spin 1/2 or 1 that have been introduced in the literature; the results of Ref. [21] are relevant to the latter. We want to emphasize that ways of writing the quasipole wavefunctions other than the one presented here could still be of potential interest for the purpose of gaining useful physical insight, or for performing further calculations of the properties of the quasipoles.

Note added: After submission of the original version of this paper, some additional facts have come to light which it may be useful to describe here for completeness. First, I have become aware that an earlier, difficult paper by Feigin and Loktev obtains some of the results herein, including the counting formula, eq. (18). In addition, there is a connection with other recent work on the zero-energy states for the $k + 1$ -body Hamiltonian. Several authors [23–25] (some of whom concentrated on $k = 2$ only) have discovered that for the system on a torus or cylinder, as the radius is taken to zero, an effective description emerges that is combinatorially simple. [This builds on earlier corresponding results for the Laughlin Abelian ($k = 1$) states, and others [26].] For the cylinder (which topologically is the same as the sphere), we will consider the single-particle orbitals (corresponding to single-particle L_z eigenstates z^m on the sphere) to be labeled by $m + 1 = 1, \dots, N_\phi + 1$, arranged in sequence on a line. Then the recipe is that, for $M = 0$, the Hamiltonian stipulates that in a zero-energy state, any two neighboring orbitals can be occupied by a total of not more than k particles, which can be distributed in any way between the two orbitals. (For the $M = 1$ case, the first one relevant to fermions, this becomes no more than k in any group of $k + 2$ neighboring orbitals, with of course not more than one in each orbital because of the Pauli exclusion principle.) For the torus, the same Hamiltonian applies for orbitals with periodic boundary conditions, $N_\phi + 1 \equiv 1$, with ‘‘neighboring orbitals’’ interpreted cyclically. Clearly, either of these specifies a combinatorial problem in which one would like to count the total number of such states for each N , N_ϕ , k , and one would hope that the result for the sphere (cylinder) agrees with the formula, eq. (18), obtained here.

This type of problem is connected with a long history of results in combinatorics, for which see Andrews’ book [27]. The problem for bosons in the cylinder/sphere case can be related to counting partitions, by identifying the length of each row of (the Ferrers-Young diagram of) a partition with the orbital number $m + 1 = 1, 2, \dots$, of a corresponding particle. As partitions are defined to have rows of (not strictly) decreasing lengths, each partition corresponds to a unique many-particle basis state for a system of bosons, the number N of bosons being

the number of rows of non-zero length in the partition. (Similarly, for fermions there is a correspondence with partitions with all rows having distinct lengths. There is a correspondence between the two problems, obtained by adding $N - i$ to the length of the i th row of a partition for the boson case to obtain the corresponding partition for fermions.) Notice that the total number of boxes \mathcal{N} (the number being partitioned) is related to the total L_z by $NN_\phi/2 + N - L_z = \mathcal{N}$ for $M = 0$. Then in the limit $N_\phi \rightarrow \infty$, in which there is no upper limit on lengths of rows, the simplest case is the $k = 1$ case for bosons. The zero-energy states then correspond to partitions in which two neighboring rows have lengths differing by at least two. The number of these is described by a generating function that can be written in two ways; this is the celebrated Rogers-Ramanujan identity, which is thus connected combinatorially with the quasiholes of the Laughlin $\nu = 1/2$ state in the plane. The generalization to $k > 1$ is given by the Gordon-Andrews identities, see Ref. [27]. One side of each of these identities agrees with the $N_\phi \rightarrow \infty$ limit of our result, studied earlier in [14,15]. (There are further results also, in which some number l , $0 \leq l \leq k$ of quasiholes are fixed at the origin [14,15]. In the chain of orbitals, this corresponds to a restriction that the occupation of the first orbital is $\leq k - l$.)

To make full contact with our results, we require results for partitions with the additional restriction of a maximum length for the rows. For the $k = 1$ case, this “finitization” of the Rogers-Ramanujan identities was found in Ref. [28]. For $k > 1$, the number of partitions obeying the given rules, and the appropriate finitization of the Gordon-Andrews identities, was derived in detail by Warnaar [29] in a manner very close to the above interpretation as particles in orbitals. [The identities are also given in Ref. [22], in a “grand canonical” form in which one multiplies the results for fixed N by x^N (where x is another indeterminate) and then sums over N with N_ϕ fixed.] Here boundary conditions can be imposed at the two ends of the chain, that the occupation of the first orbital is $\leq k - l$, and that of the last is $\leq k - l'$. For the quantum Hall states on the sphere, these correspond to fixing l quasiholes at the north pole, l' at the south pole. One side of each of these identities (specialized to the case $l = l' = 0$) agrees with the formula (18) (times q^N). These results thus confirm that the counting produced by these simple one-dimensional Hamiltonians [26,23–25] reproduces that obtained from the two-dimensional analysis here.

For the torus, the periodic boundary condition on the one-dimensional system can also be handled. The ends of the open chain can be joined and the restriction of total occupation $\leq k$ for the pair of orbitals 1, N_ϕ can be imposed by taking Warnaar’s formulas, setting $l + l' = k$, and summing over l to obtain the total number of zero-energy states. It is an open problem to derive this using the wavefunctions on the torus in a fashion similar to

that of the present paper.

I am grateful to N.R. Cooper for many stimulating discussions and for pointing out several errors in the manuscript, and to R. Howe and I.B. Frenkel for helpful remarks. This work was supported by NSF grant no. DMR-02-42949.

Appendix A: The proof that a symmetric polynomial that is non-zero when $z_1 = z_2 = \dots = z_k$, and zero when in addition $z_{k+1} = z_1$, vanishes quadratically ($(z_{k+1} - z_1)^2$ (or faster) as $z_{k+1} \rightarrow z_1$, goes as follows. We will work on the sphere [13], using the homogeneous or spinor particle coordinates u_i, v_i , in terms of which $z_i = v_i/u_i$ for each i . We will use auxiliary coordinates like the w_l in eq. (2) to form coherent states; these correspond to spinors α_l, β_l with $w_l = \beta_l/\alpha_l$ (examples of such functions appear in the main text). In this form, any wavefunction is rotationally invariant if the rotation acts on all of the α_l, β_l pairs as well as on all the u_i, v_i pairs. That is, the wavefunction is a sum of products of factors of the forms $u_i v_j - v_i u_j$ (or similarly with α_l, β_l pairs in place of one or both particle spinors). This is because this singlet combination of two spinors is the only way to construct invariant functions out of the N spinors u_i, v_i and auxiliary spinors α_l, β_l . For our purposes, the polynomials must be homogeneous of degree N_ϕ in each u_i, v_i (corresponding to each particle having angular momentum $N_\phi/2$), and also symmetric under permutations of the particles. Then a wavefunction can be analyzed into terms according to the angular momentum \mathbf{L} of some set of $k + 1$ particles, schematically $\Psi = \sum_{L, L_z} \Psi_{L, L_z}(k + 1) \Psi_{L, -L_z}(N - k - 1)$, where $\Psi_{L, L_z}(k + 1)$ is a function of the first $k + 1$ coordinates only, and $\Psi_{L, -L_z}(N - k - 1)$ is a function of the remaining particle and all the auxiliary coordinates. These are combined so that the state Ψ is a singlet. If the state $\Psi_{L, L_z}(k + 1)$ for $L = L_{\max} = (k + 1)N_\phi/2$ is nonzero, then in particular its component with $L_z = L$ is nonzero, and this is represented by $\prod_i u_i^{N_\phi}$, so that it is nonzero when all $v_i = 0$ (i.e. $z_1 = \dots = z_{k+1} = 0$). Hence we require these $L = L_{\max}$ components to vanish. The rate at which the function vanishes as $z_{k+1} \rightarrow z_1 = z_2 = \dots = z_k = 0$ is determined by the largest angular momentum value L for which the component $\Psi_{L, L_z}(k + 1)$ is nonzero. By analyzing the totally symmetric states of $k + 1$ spins of magnitude $N_\phi/2$, one finds that the largest values are $L_{\max}, L_{\max} - 2, \dots$ (the multiplicities are one for these first two terms, but larger than one for subsequent terms). Thus the first subleading term contains factors $(u_i v_j - v_i u_j)^2$, or $(z_i - z_j)^2$, which is what we wanted to prove. This proof uses the sphere, but can presumably be repeated directly for the plane by reference to the center of mass as well as the total degree. However, this is unnecessary, as the counting of states for the plane subject to an up-

per limit N_ϕ on the degree in each variable is always the same as for the sphere.

From this result, it follows that if coordinates $z_{k+1}, z_{k+2}, \dots, z_{2k}$ approach $z_1 = \dots = z_k$, then the wavefunction must vanish quadratically in each of the second k coordinates, and so if $z_{k+1} = \dots = z_{2k}$, it vanishes as $(z_1 - z_{k+1})^{2k}$ as $z_{k+1} \rightarrow z_1$, as was also used in the form of the residues.

Appendix B: Here we describe some alternative quasihole wavefunctions, which are natural generalizations of the ground state and quasihole wavefunctions

$$\tilde{\Psi}_{N, N_\phi, k}(z_1, \dots, z_N; w_1, \dots, w_n) = \mathcal{S}_z \left\{ \prod_{i < j}^{(1)} (z_i - z_j)^2 \cdots \prod_{i < j}^{(k)} (z_i - z_j)^2 \prod_{i, l}^{(1)} (z_i - w_l) \cdots \prod_{i, l}^{(k)} (z_i - w_l) \right\}. \quad (20)$$

In the products $\prod^{(\alpha)}$ (for $\alpha = 1, \dots, k$), i, j run over $\sum_{\beta: \beta < \alpha} N_\beta + 1, \dots, \sum_{\beta: \beta \leq \alpha} N_\beta$, and also l runs over $\sum_{\beta: \beta < \alpha} \hat{n}_\beta + 1, \dots, \sum_{\beta: \beta \leq \alpha} \hat{n}_\beta$. To ensure that each function is of the same degree in all coordinates z_i , we impose $2N_\alpha + \hat{n}_\alpha = (2N + n)/k$ for all α (which are equivalent to the earlier conditions). Note that each function need not be of the same degree in all the w 's, and one should not symmetrize over the w 's. It is easy to see that when any $k + 1$ z 's are equal, the function vanishes, because in any term in the symmetrization over the z 's, at least two must be of the same "type" α . In the limit as $z_{(p-1)k+1}, \dots, z_{(p-1)k+k} \rightarrow Z_p$ (for $p = 1, \dots, N_k = m_k$), the function vanishes if a further set of k coordinates coincide with one another. Thus each of them reduces to the form of a non-vanishing residue with $F = N - kN_k$, but in which $\tilde{\Psi}_{F, N'_\phi, k'}$ is of the form (20) with the parameters as given, and which, unlike the residues above, still depends on the first $n - \hat{n}_k$ w_l 's as well as F of the z 's. In this function $\tilde{\Psi}_{F, N'_\phi, k'}$, the partition of N is replaced by a similar partition of F , with $F_\alpha = N_\alpha - N_k$ ($\alpha = 1, \dots, k'$), while the partition \hat{n}_α of n is unchanged, except that \hat{n}_k must be dropped.

The fact that $\tilde{\Psi}_{F, N'_\phi, k'}$ here depends on some of the same w 's as appear in front of it in the residue makes it nontrivial to verify that each nonvanishing residue can be obtained from some (linear combination of) functions of this form. One would like to expand the non-vanishing residues that result from eq. (20) in linearly-independent polynomials in w 's, in order to find the dimension of the space spanned by the coherent states. The clustered part of the residue produces a linearly-independent set of $\binom{(N-F)/k+n}{n}$ symmetric polynomials in the w 's, while the remaining factor $\tilde{\Psi}_{F, N'_\phi, k'}$ would be assumed by induction to be spanned by some linearly-independent set of $\#_{F, N'_\phi, k'}$ polynomials in the w 's. However, we were not able to establish that the total number of indepen-

dent polynomials in w 's in the products of these two sets is given by eq. (6). The reason is that the set of polynomials in w 's obtained by multiplying one member of one of these two sets of polynomials by a member of the other may not be linearly independent, even though both the original sets were. (This may be formulated mathematically as asking whether the set that spans the expansion of $\tilde{\Psi}_{F, N'_\phi, k'}$ is linearly-independent *over the algebra of symmetric polynomials*, not just over the complex numbers.) We were not able to establish the linear independence of the resulting set in general, or that the number of independent functions equals the required number, though in special cases of small systems it does [19]. If $\tilde{\Psi}_{F, N'_\phi, k'}$ could be expanded in S_n -harmonic polynomials in w 's, then the argument would go through. As usual, S_n denotes the symmetric (or permutation) group on n objects. The S_n -harmonic polynomials are a linearly-independent set of $n!$ polynomials in the n variables w_l which, when multiplied by arbitrary symmetric polynomials, span the full space of polynomials in w 's (thus, they form a basis for the latter vector space over the algebra of symmetric polynomials) [30]. Though, in the cases we checked, there are enough S_n -harmonic polynomials in each degree for this to work (and one must note that permutations of the w_l 's in the wavefunctions (20) leave the functions of z 's invariant, so only one member of each orbit under S_n in the S_n -harmonic polynomials can be used), we do not know if the functions can always be expanded in this way. Thus, while we suspect that the quasihole wavefunctions in eq. (20) do span the space $V_{N, N_\phi, k}$ for all N, N_ϕ, k , we have been unable to prove this so far.

-
- [1] G. Moore and N. Read, Nucl. Phys. **B 360**, 362 (1991).
 - [2] N. Read and E. Rezayi, Phys. Rev. B **59**, 8084 (1999).
 - [3] A. Cappelli, L.S. Georgiev, and I.T. Todorov, Nucl. Phys.

- B **599**, 499 (2001).
- [4] C. Nayak and F. Wilczek, Nucl. Phys. B**479**, 529 (1996).
- [5] N. Read and E. Rezayi, Phys. Rev. B **54**, 16864 (1996).
- [6] V. Gurarie and E. Rezayi, Phys. Rev. B **61**, 5473 (2000).
- [7] E. Ardonne, J. Phys. A **35**, 447 (2002).
- [8] N. Regnault and T. Jolicoeur, cond-mat/0601550.
- [9] R.H. Morf, Phys. Rev. Lett. **80**, 1505 (1998); E.H. Rezayi and F.D.M. Haldane, Phys. Rev. Lett. **84**, 4685 (2000).
- [10] N.R. Cooper, N.K. Wilkin, and J.M.F. Gunn, Phys. Rev. Lett. **87**, 120405 (2001).
- [11] E.H. Rezayi, N. Read, and N.R. Cooper, Phys. Rev. Lett. **95**, 160404 (2005).
- [12] M.H. Freedman, A. Kitaev, M.J. Larsen, and Z. Wang, quant-ph/0101025.
- [13] F.D.M. Haldane, Phys. Rev. Lett. **51**, 605 (1983).
- [14] A.V. Stoyanovsky and B.L. Feigin, Funct. Anal. Appl. **28**, 55 (1994); B. Feigin, M. Jimbo, R. Kedem, S. Loktev, and T. Miwa, J. Alg. **279**, 147 (2004).
- [15] E. Ardonne, R. Kedem, and M. Stone, J. Phys. A **38**, 617 (2005).
- [16] E. Ardonne, N. Read, E. Rezayi, and K. Schoutens, Nucl. Phys. B **607**, 549 (2001).
- [17] K. Schoutens, Phys. Rev. Lett. **79**, 2608 (1997).
- [18] This formula can be proved by induction on m_k , or by other methods; see e.g. V. Kac and P. Cheung, *Quantum Calculus* (Springer, New York, NY, 2002).
- [19] N.R. Cooper, private communication.
- [20] M. Milovanovic and N. Read, Phys. Rev. B **53**, 13559 (1996).
- [21] E. Ardonne, R. Kedem, and M. Stone, J. Phys. A **38**, 9183 (2005); math.RT/0504364.
- [22] B.L. Feigin and S.A. Loktev, Funct. Anal. Appl. **35**, 44 (2001).
- [23] F.D.M. Haldane, talk at APS March Meeting, Baltimore, March, 2006.
- [24] E.J. Bergholtz, J. Kailasvuori, E. Wikberg, T.H. Hansson, and A. Karlhede, cond-mat/0604251.
- [25] A. Seidel and D.-H. Lee, cond-mat/0604465.
- [26] E.H. Rezayi and F.D.M. Haldane, Phys. Rev. B. **50**, 17199 (1994); E.J. Bergholtz and A. Karlhede, Phys. Rev. Lett. **94**, 26802 (2005); cond-mat/0509434; A. Seidel, H. Fu, D.-H. Lee, J.M. Leinaas, J. Moore, Phys. Rev. Lett. **95**, 266405 (2005).
- [27] G.E. Andrews, *The Theory of Partitions*, (Cambridge University Press, Cambridge, 1984), especially Ch. 7.
- [28] G.E. Andrews, Scripta Math. **28**, 297 (1970), or see pp. 50, 157 in Ref. [27].
- [29] S.O. Warnaar, Commun. Math. Phys. **184**, 203 (1997).
- [30] R. Howe, private communication.