

Generalized Statistics Framework for Rate Distortion Theory with Bregman Divergences

R. C. Venkatesan
Systems Research Corporation
Aundh, Pune 411007, India
ravi@systemsresearchcorp.com

Abstract—A variational principle for the rate distortion (RD) theory with Bregman divergences is formulated within the ambit of the generalized (nonextensive) statistics of Tsallis. The Tsallis-Bregman RD lower bound is established. Alternate minimization schemes for the generalized Bregman RD (GBRD) theory are derived. A computational strategy to implement the GBRD model is presented. The efficacy of the GBRD model is exemplified with the aid of numerical simulations.

I. INTRODUCTION

The generalized (nonextensive) statistics of Tsallis [1,2] has recently been the focus of much attention in statistical physics, and allied disciplines¹. Nonextensive statistics generalizes the extensive Boltzmann-Gibbs statistics, and has found much utility in complex systems possessing long range correlations, fluctuations, ergodicity, chirality and fractal behavior. By definition, the Tsallis entropy is defined in terms of discrete variables as

$$S_q(x) = -\frac{1 - \sum_x p^q(x)}{1 - q}, \text{ where, } \sum_x p(x) = 1. \quad (1)$$

The constant q is referred to as the *nonextensivity parameter*. Given two independent variables x and y , one of the fundamental consequences of nonextensivity is demonstrated by the *pseudo-additivity* relation

$$S_q(xy) = S_q(x) + S_q(y) + (1 - q) S_q(x) S_q(y). \quad (2)$$

Here, (1) and (2) imply that extensive statistics is recovered as $q \rightarrow 1$. Taking the limit $q \rightarrow 1$ in (1) and evoking l'Hospital's rule, $S_q(x) \rightarrow S(x)$, the Shannon entropy. The *jointly convex* generalized Kullback-Leibler divergence (K-Ld) is of the form [3]

$$I_q(p(x) \| r(x)) = \sum_x p(x) \frac{\left(\frac{p(x)}{r(x)}\right)^{q-1} - 1}{q-1}. \quad (3)$$

In the limit $q \rightarrow 1$, the extensive K-Ld is readily recovered. Akin to the Tsallis entropy, the generalized K-Ld obeys the *pseudo-additivity* relation [3].

Defining the q -deformed logarithm and the q -deformed exponential as [4]

$$\ln_q(x) = \frac{x^{1-q} - 1}{1 - q}, \quad (4)$$

and

$$e_q^x = \begin{cases} [1 + (1 - q)x]^{1/(1-q)}; & 1 + (1 - q)x \geq 0 \\ 0; & \text{otherwise,} \end{cases} \quad (5)$$

respectively. The Tsallis entropy (1), and, the generalized K-Ld (3) may be written as

$$S_q(p(x)) = -\sum_x p(x)^q \ln_q p(x), \quad (6)$$

and,

$$I_q(p(x) \| r(x)) = -\sum_x p(x) \ln_q \frac{r(x)}{p(x)}, \quad (7)$$

respectively. Employing the relation [4]

$$\ln_q\left(\frac{x}{y}\right) = y^{q-1} (\ln_q x - \ln_q y), \quad (8)$$

the generalized K-Ld (7) may be expressed as the generalized mutual information

$$\begin{aligned} I_q(X; \tilde{X}) &= -\sum_{x, \tilde{x}} p(x, \tilde{x}) \ln_q \left(\frac{p(\tilde{x})}{p(\tilde{x}|x)}\right) \\ &= \sum_{\tilde{x}} \frac{1}{p^{q-1}(\tilde{x})} S_q(X) - \underbrace{\sum_{\tilde{x}} p(\tilde{x}) S_q(X | \tilde{X} = \tilde{x})}_{\langle S_q(X | \tilde{X} = \tilde{x}) \rangle_{p(\tilde{x})}}. \end{aligned} \quad (9)$$

Here, $\langle \bullet \rangle$ denotes the expectation value. The seminal paper on source coding within the framework of nonextensive statistics by Landsberg and Vedral [5], has provided the impetus for a number of investigations into the use of nonextensive information theory within the context of coding problems. The works of Yamano [6,7] represent a sample of some of the prominent efforts in this regard. *The source coding theorem in a nonextensive setting has been derived by Yamano [8].*

A statistical physics model for the variational problem encountered in rate distortion (RD) theory [9, 10] and the information bottleneck method [11], derived within a generalized statistics framework, has been recently established [12]. A nonextensive Blahut-Arimorto (BA) alternate minimization scheme [13] has been derived. The noteworthy result of the study in [12] is that the nonextensive RD curves possesses a

¹A continually updated bibliography of works related to nonextensive statistics may be found at <http://tsallis.cat.cbpf.br/biblio.htm>.

lower threshold for the minimum compression information in the distortion-compression plane, as compared to equivalent RD curves derived on the basis of the Boltzmann-Gibbs-Shannon framework. As was the case in [12], this paper employs values of q in the range $0 < q < 1$. This paper utilizes the statistical physics theory presented in [12] to formulate a generalized Bregman RD (GBRD) model. The GBRD model extends previous works by Rose [14] and Banerjee *et. al.* [15, 16] to achieve a principled and practical strategy to evaluate RD functions.

II. RATIONALE FOR THE GBRD MODEL

A. Background information

The RD problem in terms of discrete random variables is stated as follows: given a discrete random variable $X \in \Xi$ called the source or the codebook, and, another discrete random variable $\tilde{X} \in \tilde{\Xi}$ which is a compressed representation of X (also referred to as the *quantized codebook* and/or the *reproduction alphabet*), the information rate distortion function that is to be obtained is the minimization of the mutual information $I_{q \geq 1}(X, \tilde{X})$ over all conditional probabilities $p(\tilde{x}|x)$.

The crux of the RD problem is the numerical determination of the RD function using the BA scheme. The actual implementation of the BA scheme is sometimes impractical owing to a lack of knowledge of the optimal support of the *quantized codebook* \tilde{X} . Exact analytical solutions exist only for a few cases consisting of a combination well behaved sources and distortion measures. An initial attempt to achieve a practical and tractable solution to the RD problem was performed by Rose [14], for the case of Euclidean square distortion functions. Therein, it was demonstrated that for sources whose support is a bounded set, the RD function either equals the *Shannon lower bound*, or, the optimal support for the *quantized codebook* is finite thus permitting the use of a numerical procedure called the *mapping method*. The pioneering work of Rose was generalized by Banerjee *et. al* [15, 16] to include a wider class of distortion functions using Bregman divergences. One of the significant features in these works involved the formulation of a *Shannon-Bregman lower bound*.

Motivated by the recent results [12], this paper provides two means to solve the GBRD problem for sources with bounded support. First, the analytical solution may be obtained from a *Tsallis-Bregman lower bound* (Section IV). Next, for a finite reproduction alphabet, the RD function may be numerically obtained by a computational methodology derived from generalized statistics (Section III).

B. Bregman Divergences

Definition 1 (Bregman divergences): Let ϕ be a real valued strictly convex function defined on the convex set $S \subseteq \mathfrak{R}$, the domain of ϕ such that ϕ is differentiable on $\text{int}(S)$, the interior of S . The Bregman divergence $B_\phi : S \times \text{int}(S) \mapsto \mathfrak{R}^+$ is defined as $B_\phi(z_1, z_2) = \phi(z_1) - \phi(z_2) - (z_1 - z_2, \nabla\phi(z_2))$, where $B_\phi(z_1, z_2) = \phi(z_1) - \phi(z_2) - (z_1 - z_2, \nabla\phi(z_2))$ is the gradient of ϕ evaluated at z_2 .

A number of Bregman divergences have been tabulated in [15]. The generalized K-Ld in (3), also referred to as the Csizsár generalized K-Ld is not a Bregman divergence. Since the seminal work by Naudts [17], Bregman divergences have been the object of much research in nonextensive statistics. Defining $\phi(p) = -p \ln_q \left(\frac{1}{p} \right)$, the Bregman generalized K-Ld is defined as

$$\begin{aligned} d_\phi(\mathbf{p}, \mathbf{r}) &= \sum_{i=1}^d \frac{p_i}{(q-1)} \left(p_i^{q-1} - r_i^{q-1} \right) - \sum_{i=1}^d (p_i - r_i) r_i^{q-1} \\ &= I_{q,B}(\mathbf{p} \parallel \mathbf{r}) \end{aligned} \quad (10)$$

Setting $q - 1 \rightarrow \kappa$, (10) is consistent with Eq. (35) in [17].

III. THE GBRD MODEL

This Section provides a strategy to jointly obtain the optimal support $\tilde{\mathcal{X}}_s^2$ of the *quantized codebook* with cardinality $|\tilde{\mathcal{X}}_s| = k$, and, the conditional probability $p(\tilde{x}|x)$ that characterizes the RD problem. This is accomplished by a joint optimization of

$$\min_{\tilde{\mathcal{X}}_s, p(\tilde{x}|x)} L_{GBRD}^q = \min_{\substack{\tilde{\mathcal{X}}_s, p(\tilde{x}|x) \\ |\tilde{\mathcal{X}}_s|=k}} \left\{ I_q(X; \tilde{X}) + \beta \langle d_\phi(x, \tilde{x}) \rangle_{p(x, \tilde{x})} \right\}. \quad (11)$$

The optimal Lagrange multiplier β , hereafter referred to as the *inverse temperature*, depends upon the optimal tolerance level of the expectation of $D = \langle d_\phi(x, \tilde{x}) \rangle_{p(x, \tilde{x})} = \sum_{x, \tilde{x}} p(x, \tilde{x}) d_\phi(x, \tilde{x})$.

A. Constraint terms

Generalized statistics has utilized a number of constraints to define expectations. The original Tsallis (OT) constraints of the form $\langle A \rangle = \sum_i p_i A_i$ [1], were convenient owing to their similarity to the maximum entropy constraints. These were abandoned because of difficulties encountered in obtaining an acceptable form for the partition function, with the exception of a few specialized cases.

The OT constraints were subsequently replaced by the Curado-Tsallis (C-T) constraints $\langle A \rangle_q^{C-T} = \sum_i p_i^q A_i$. The C-T constraints were later replaced by the normalized Tsallis-Mendes-Plastino (T-M-P) constraints $\langle A \rangle^{T-M-P} = \sum_i \frac{p_i^q}{\sum_i p_i^q} A_i$. The dependence of the expectation value on the normalization *pdf* $\sum_i p_i^q$, renders the T-M-P constraints to be *self-referential*. This paper, like [12], utilizes a recent methodology [18] to “rescue” the OT constraints, and, has linked the OT, C-T, and, T-M-P constraints.

B. The nonextensive variational principle

Keeping $\tilde{\mathcal{X}}_s$ fixed, taking the variational derivative of (11) with respect to $p(\tilde{x}|x)$ while enforcing $\sum_{\tilde{x}} p(\tilde{x}|x) = 1$ with

²Calligraphic symbols indicate sets

the normalization Lagrange multiplier $\lambda(x)$, yields

$$p(\tilde{x}|x) = p(\tilde{x}) \left[\frac{(1-q)}{q} \left\{ \tilde{\lambda}(x) + \beta d_\phi(x, \tilde{x}) \right\} \right]^{1/(q-1)},$$

$$\tilde{\lambda}(x) = \frac{\lambda(x)}{p(x)} - p(x)^{(1-q)}. \quad (12)$$

Multiplying the terms in the square brackets in (12) by the conditional probability $p(\tilde{x}|x)$, and summing over \tilde{x} yields

$$\frac{q}{q-1} \sum_{\tilde{x}} p(\tilde{x}) \left(\frac{p(\tilde{x}|x)}{p(\tilde{x})} \right)^q + \beta \underbrace{\sum_{\tilde{x}} d_\phi(x, \tilde{x}) p(\tilde{x}|x)}_{\beta \langle d_\phi(x, \tilde{x}) \rangle_{p(\tilde{x}|X=x)}} + \tilde{\lambda}(x) \sum_{\tilde{x}} p(\tilde{x}|x) = 0. \quad (13)$$

Evoking $\sum_{\tilde{x}} p(\tilde{x}|X=x) = 1$, yields

$$\tilde{\lambda}(x) = \frac{q}{(1-q)} \aleph_q(x) - \beta \langle d_\phi(x, \tilde{x}) \rangle_{p(\tilde{x}|X=x)}. \quad (14)$$

Here, $\aleph_q(x) = \sum_{\tilde{x}} p(\tilde{x}) \left(\frac{p(\tilde{x}|x)}{p(\tilde{x})} \right)^q$. The conditional pdf $p(\tilde{x}|x)$ acquires the form

$$p(\tilde{x}|x) = \frac{p(\tilde{x}) \{1 - (q-1)\beta^* d_\phi(x, \tilde{x})\}^{1/(q-1)}}{\mathfrak{S}_{RD}^{1/(1-q)}},$$

$$\mathfrak{S}_{RD} = q \aleph_q(x) + (q-1) \beta \langle d_\phi(x, \tilde{x}) \rangle_{p(\tilde{x}|X=x)}, \quad (15)$$

$$\beta^* = \frac{\beta}{\mathfrak{S}_{RD}},$$

where β^* is the *effective inverse temperature*. Transforming $q \rightarrow 2 - q^*$ in the numerator and evoking (5), (15) is expressed in the form of a q -deformed exponential

$$p(\tilde{x}|x) = \frac{p(\tilde{x}) \exp_{q^*}(-\beta^* d_\phi(x, \tilde{x}))}{\tilde{Z}(x, \beta^*)}. \quad (16)$$

The partition function evaluated at each instance of the source distribution is

$$\tilde{Z}(x, \beta^*) = \mathfrak{S}_{RD}^{\frac{1}{(1-q^*)}} = \sum_{\tilde{x}} p(\tilde{x}) \exp_q[-\beta^* d_\phi(x, \tilde{x})] \quad (17)$$

The term $\{1 - (1 - q^*) \beta^* d(x, \tilde{x})\}$ in the numerator of (16) is a manifestation of the *Tsallis cut-off condition* [18]. This implies that solutions of (16) are valid when $\beta^* d(x, \tilde{x}) < 1/(1 - q^*)$. The *effective nonextensive RD Helmholtz free energy* is

$$F_{RD}^q(\beta^*) = \frac{-1}{\beta^*} \left\langle \ln_q \tilde{Z}(x, \beta^*) \right\rangle_{p(x)} = \frac{1}{\beta^*} \left\langle \frac{\mathfrak{S}_{RD} - 1}{q - 1} \right\rangle_{p(x)}. \quad (18)$$

Solution of (16) may be viewed from two distinct perspectives, i.e. the *canonical perspective* and the *parametric perspective*. Owing to the *self-referential* nature of the *effective inverse temperature* β^* , the analysis and solution of (16) within the context of the *canonical perspective* is a formidable undertaking. For practical applications, the parametric perspective is utilized by evaluating the conditional pdf $p(\tilde{x}|x)$, employing the nonextensive BA algorithm, for a-priori specified $\beta^* \in [0, \infty]$. Note that within the context of the *parametric perspective*, the *self-referential nature* of β^*

vanishes. The *inverse temperature* β and the *effective inverse temperature* β^* relate as

$$\beta = \frac{q \aleph_q(x) \beta^*}{\left[1 - \beta^* (q-1) \langle d_\phi(x, \tilde{x}) \rangle_{p(\tilde{x}|X=x)} \right]}. \quad (19)$$

C. Support estimation step

The procedure in Section III.B. is carried out for a given β , corresponding to a single point on the RD curve. Reproduction alphabets with optimal support are obtained by keeping $p(\tilde{x}|x)$ fixed, and solving

$$\min_{\tilde{X}_s} L_{GBRD}^q = \min_{\tilde{X}_s} \left\{ I_q(X; \tilde{X}) + \beta \langle d_\phi(x, \tilde{x}) \rangle_{p(x, \tilde{x})} \right\}. \quad (20)$$

The solution to (20) is the optimal estimate/predictor [15, 16]

$$\tilde{x}^* = \langle x \rangle_{p(x|\tilde{X}=\tilde{x})} = \sum_x p(x|\tilde{X}=\tilde{x}) x. \quad (21)$$

Algorithm 1 depicts the *pseudo-code* for the greedy joint non-convex optimization procedure that constitutes the solution to the GBRD model.

IV. THE TSALLIS-BREGMAN LOWER BOUND

Theorem 1 For OT constraints, the nonextensive RD function with source $X \sim p(x)$ and a Bregman divergence d_ϕ is always lower bounded by the Tsallis-Bregman lower bound defined by

$$R_{qL}(D) = \sum_{\tilde{x}} \frac{1}{p^{q-1}(\tilde{x})} S_q(X) + \sup_{\beta \geq 0} \left\{ -\beta D + \langle \ln_{q^*} \gamma_{L, \beta^*}(x) \rangle_{p(x)} \right\}, \quad (22)$$

where γ_{L, β^*} is a unique function satisfying

$$\int_{\text{dom}(\phi)} p(t) \gamma_{L, \beta^*}(t) \exp_{q^*}(-\beta^* d_\phi(t, \mu)) dt = 1; \forall \mu \in \text{dom}(\phi). \quad (23)$$

To highlight the critical dependence of the *Tsallis-Bregman lower bound* upon the type of constraint employed to solve the GBRD variational problem in Section III B, and enunciate certain aspects of q -deformed algebra and calculus, a slightly weaker bound (see Appendix C, [16]) is stated and proved.

Theorem 2 The generalized RD function is defined by

$$R_q(D) = \sup_{\beta \geq 0, \gamma^* \in \Lambda_{\beta^*}} \left\{ -\beta D + \int_x p(x) \ln_{q^*} \gamma^*(x) dx \right\}, \quad (24)$$

where Λ_{β^*} is the set of all admissible partition functions. Further, for each $\beta \geq 0$, a necessary and sufficient condition for $\gamma^*(x)$ to attain the supremum in (24) is a probability density $p(\tilde{x})$ related to $\gamma^*(x)$ as

$$\gamma^*(x) = \left(\int_{\tilde{x}} p(\tilde{x}) \exp_q(-\beta^* d_\phi(x, \tilde{x})) d\tilde{x} \right)^{-1}. \quad (25)$$

Proof

$$\begin{aligned}
L_{GBRD}^q [p(\tilde{x}|x)] &= \\
&- \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) \ln_q \left(\frac{p(\tilde{x})}{p(\tilde{x}|x)} \right) dx d\tilde{x} + \\
&+ \beta \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) d_\phi(x,\tilde{x}) dx d\tilde{x} \\
&\stackrel{(a)}{=} \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) \ln_{q^*} \left(\frac{p(\tilde{x}|x)}{p(\tilde{x})} \right) dx d\tilde{x} + \\
&+ \beta \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) d_\phi(x,\tilde{x}) dx d\tilde{x} \\
&\stackrel{(b)}{=} \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) \ln_{q^*} \left(\frac{\exp_{q^*}(-\beta^* d_\phi(x,\tilde{x}))}{\tilde{Z}(x,\beta^*)} \right) dx d\tilde{x} + \\
&+ \beta \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) d_\phi(x,\tilde{x}) dx d\tilde{x} \\
&\stackrel{(c)}{=} - \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) \tilde{Z}^{(q-1)}(x,\beta^*) (\beta^* d_\phi(x,\tilde{x}) + \\
&\ln_{q^*} \tilde{Z}(x,\beta^*)) dx d\tilde{x} + \beta \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) d_\phi(x,\tilde{x}) dx d\tilde{x} \\
&\stackrel{(d)}{=} - \int_{x,\tilde{x}} p(x)p(\tilde{x}|x) \ln_q \tilde{Z}(x,\beta^*) dx d\tilde{x} \\
&= - \int_x p(x) \ln_q \tilde{Z}(x,\beta^*) dx \\
&\stackrel{(e)}{=} - \int_x p(x) \ln_q \left(\int_{\tilde{x}} p(\tilde{x}) \exp_{q^*}(-\beta^* d_\phi(x,\tilde{x})) d\tilde{x} \right) dx \\
&= L_{GBRD}^q [p(\tilde{x})]. \tag{26}
\end{aligned}$$

Here, (a) follows from $\ln_q(x) = -\ln_{q^*}(1/x)$; $q^* = 2 - q$, (b) follows from (16) where the partition function in the continuum is $\tilde{Z}(x,\beta^*) = \int_{\tilde{x}} p(\tilde{x}) \exp_{q^*}(-\beta^* d_\phi(x,\tilde{x})) d\tilde{x}$, (c) follows from (8), (d) follows from (15), and, (e) follows from (17). Analogous to the mapping method of [14], a mapping from the unit interval to the reproduction alphabet is sought such that the Lebesgue measure over the unit interval results in an optimal $p(\tilde{x})$ over the reproduction alphabet. Thus, for each instance of a probability measure corresponding to $p(\tilde{x})$ on $\tilde{\mathcal{X}}$ there exists a unique map $\tilde{x} : [0,1] \mapsto \tilde{\mathcal{X}}$ that maps the Lebesgue measure (μ) to $p(\tilde{x})$ such that for any function f defined over \tilde{x} , $\int_{\tilde{x}} f(\tilde{x}) p(\tilde{x}) d\tilde{x} = \int_{u \in [0,1]} f(\tilde{x}(u)) d\mu(u)$. Thus, (26) yields

$$\begin{aligned}
L_{GBRD}^q [\tilde{x}] &= \\
&= - \int_x p(x) \ln_q \left(\int_{\tilde{x}} p(\tilde{x}) \exp_{q^*}(-\beta^* d_\phi(x,\tilde{x})) d\tilde{x} \right) dx, \\
&= - \int_x p(x) \ln_q \left(\int_{u \in [0,1]} \exp_{q^*}(-\beta^* d_\phi(x,\tilde{x}(u))) d\mu(u) \right) dx. \tag{27}
\end{aligned}$$

To obtain the optimality condition, the q -deformed terms characteristic to generalized statistics are to be operated on by the *dual Jackson derivative* instead of the usual Newtonian derivative [4]. To obtain expressions analogous to the case of Newton-Leibniz calculus, the *dual Jackson derivative* defined as

$$\begin{aligned}
D^{(q)} f(x) &= \frac{1}{1+(1-q)f(x)} \frac{df}{dx}; 1+(1-q)f(x) \neq 0 \\
\Rightarrow D^{(q)} \ln_q \tilde{Z}(x,\beta^*) &= \frac{1}{\tilde{Z}(x,\beta^*)} \frac{d\tilde{Z}(x,\beta^*)}{dx} \tag{28}
\end{aligned}$$

is employed to enforce the optimality condition. The optimality condition for (27) becomes

$$D^{(q)} L_{GBRD}^q = - \int_x \frac{p(x) \frac{d}{dx} \left(\exp_{q^*}(-\beta^* d_\phi(x,\tilde{x}(u))) \right) dx}{\int_{u \in [0,1]} \exp_{q^*}(-\beta^* d_\phi(x,\tilde{x}(u))) u d\mu}. \tag{29}$$

Evoking *Borel isomorphism*, and, assuming that the optimal support $\tilde{\mathcal{X}}_s$ contains a non-empty open ball B_ϵ , $\epsilon > 0$ implies

that

$$\begin{aligned}
D^{(q)} L_{GBRD}^q &= \int_x \frac{p(x) \exp_{q^*}(-\beta^* d(x,\tilde{x})) dx}{N(x,\beta^*)} = k_o \forall \tilde{x} \in B_\epsilon \subset \tilde{\mathcal{X}}_s, \\
\Rightarrow \int_{\tilde{x} \in B_\epsilon} \int_x \frac{p(x)p(\tilde{x}) \exp_{q^*}(-\beta^* d(x,\tilde{x})) dx d\tilde{x}}{N(x,\beta^*)} &= k_o \int_{\tilde{x} \in B_\epsilon} p(\tilde{x}) d\tilde{x}, \\
\Rightarrow \int_{\tilde{x} \in B_\epsilon} \int_x p(x)p(\tilde{x}|x) dx d\tilde{x} &= k_o \int_{\tilde{x} \in B_\epsilon} p(\tilde{x}) d\tilde{x} \Rightarrow 1 = k_o, \tag{30}
\end{aligned}$$

holds true for all $\tilde{x} \in B_\epsilon$. Defining $\gamma_{\beta^*}^*(x) = \left(\int_{\tilde{x}} p(\tilde{x}) \exp_{q^*}(-\beta^* d_\phi(x,\tilde{x})) d\tilde{x} \right)^{-1}$, (30) yields

$$\int_x p(x) \gamma_{\beta^*}^*(x) \exp_{q^*}(-\beta^* d(x,\tilde{x})) dx = 1; \forall \tilde{x} \in B_\epsilon. \tag{31}$$

Using the relation $-\ln_q(x) = \ln_{q^*}(1/x)$, (26) becomes

$$L_{GBRD}^q [p(\tilde{x})] = \int_x p(x) \ln_{q^*} \gamma_{\beta^*}^*(x) dx. \tag{32}$$

Thus, $\gamma_{\beta^*}^*(x)$ satisfies (25) and attains the supremum in (24) for a given β and corresponding β^*

$$R_q(D_\beta) = -\beta D_\beta + \int_x p(x) \ln_{q^*} \gamma_{\beta^*}^*(x) dx, \tag{33}$$

where D_β is the distortion value at which the supremum in (24) is attained for a given β . Thus, the *Tsallis-Bregman lower bound* is

$$R_{qL}(D_\beta) = -\beta D_\beta + \int_x p(x) \ln_{q^*} \gamma_{\beta^*}^*(x) dx = R_q(D_\beta). \tag{34}$$

V. NUMERICAL SIMULATIONS

The efficacy of the GBRD model is computationally investigated by drawing a sample of 1000 two-dimensional data points, from three spherical Gaussian distributions with centers (2,3.5), (0,0), (0,2) (the *quantized codebook*). The priors and standard deviations are 0.3,0.4,0.3, and, 0.2,0.5,1.0, respectively. *To test effectiveness of the support estimation step, the quantized codebook is shifted from the true means to positions at the edges of the spherical Gaussian distributions.* The computational procedure described in Algorithm 1 is repeatedly solved for each value of β^* , till a reproduction alphabet with optimal support is obtained. This consistently coincides with the true mean, for a negligible error. This effect is particularly pronounced, and rapidly achieved, for regions of low and intermediate values of β^* , thus providing implicit proof of the relation between soft clustering and RD with Bregman divergences [15].

Fig.1 depicts the RD curves for extensive RD with Bregman divergences [15, 16] and the GBRD model, with the constituent discrete points overlaid upon them. A Euclidean square distortion (a Bregman divergence) is employed. Each curve has been generated for values of $\beta \in [.1, 2.5]$ (the extensive case), and $\beta^* \in [.1, 100]$ (the GBRD cases), respectively. *Note that for the GBRD cases, the slope of the RD curve is $-\beta$ and not $-\beta^*$. Note that all GBRD curves inhabit the non-achievable (no compression) region of the extensive RD model with Bregman divergences. Further, GBRD models possessing a lower nonextensivity parameter q inhabit the non-achievable regions of GBRD models possessing a higher value of q . It*

Algorithm 1 GBRD Model

Input

1. $X \sim p(x)$ over $\{x_i\}_{i=1}^n \subset \text{dom}(\phi) \subseteq \mathbb{R}^m$.
2. Bregman divergence d_ϕ , $|\tilde{\mathcal{X}}_s| = k$, effective variational parameter $\beta^* \in [0, \infty]$, each β^* is a single point on the RD curve.

Output

1. $\tilde{\mathcal{X}}_s^* = \{\tilde{x}_j\}_{j=1}^k$, $P^* = \left\{ \left\{ \tilde{x}_j | x_i \right\}_{j=1}^k \right\}_{i=1}^n$ that locally optimizes (11), (R_β, D_β) tradeoff at each β^*
2. Value of $R_q(D)$ where its slope equals $-\beta = \frac{1 - \beta^*(q-1) \langle d_\phi(x, \tilde{x}) \rangle_{p(\tilde{x}|x=\cdot)}}{q \mathcal{N}_q(x) \beta^*}$.

Method

1. Initialize with some $\{\tilde{x}_j\}_{j=1}^k \subset \text{dom}(\phi)$.

2. Set up outer β^* loop.

repeat

Blahut-Arimoto loop (16) for single value of β^*

repeat

for $i=1$ to n do

for $j=1$ to k do

$$p(\tilde{x}_j | x_i) \leftarrow \frac{p(\tilde{x}_j) \exp_{q^*}(-\beta^* d_\phi(x_i, \tilde{x}_j))}{Z(x_i, \beta^*)}$$

end for

end for

for $j=1$ to k do

$$p(\tilde{x}_j) \leftarrow \sum_{i=1}^n p(\tilde{x}_j | x_i) p(x_i)$$

end for

until convergence

Support Estimation Step ($\tilde{\mathcal{X}}_s$ using (21))

for $j=1$ to k do

$$\tilde{x}_j \leftarrow \sum_{i=1}^n p(x_i | \tilde{x}_j) x_i$$

end for

until convergence

Calculate D_β and R_β for the β^* .

advance β^*

$$\beta^* = \beta^* + \delta\beta^*$$

is observed that the GBRD model undergoes compression and clustering more rapidly than the equivalent extensive RD model with Bregman divergences. A primary cause for such behavior is the rapid increase in β^* for marginal increases in β , as depicted in Fig. 2 and obtained from (19).

ACKNOWLEDGEMENT

This work was supported by *RAND-MSR contract CSM-DI & S-QIT-101107-2005*. Gratitude is expressed to A. Plastino, S. Abe, and, K. Rose for helpful discussions.

REFERENCES

- [1] C. Tsallis. Possible Generalizations of Boltzmann-Gibbs Statistics. *J. Stat. Phys.*, **542**, pp 479-487, 1988.
- [2] M. Gell-Mann and C. Tsallis (Eds.). *Nonextensive Entropy-Interdisciplinary Applications*. Oxford University Press, New York, 2004.
- [3] C. Tsallis. Generalized Entropy-Based Criterion for Consistent Testing. *Phys. Rev. E*, **58**, pp 1442-1445, 1998.
- [4] E. Borges. A Possible Deformed Algebra and Calculus Inspired in Nonextensive Thermostatistics. *Physica A*, **340**, pp. 95-111, 2004. Manuscript available at <http://arxiv.org/abs/cond-mat/0304545>.

- [5] P.T. Landsberg and V. Vedral. Distributions and Channel Capacities in Generalized Statistical Mechanics. *Phys. Lett. A*, **247**, pp. 211-217, 1998.
- [6] T. Yamano. Information Theory based on Nonadditive Information Content. *Phys. Rev. E*, **63**, pp. 046105-046111, 2001.
- [7] T. Yamano. Generalized Symmetric Mutual Information Applied for the Channel Capacity. *Phys. Rev. E.*, To Appear. Manuscript available at <http://arxiv.org/abs/cond-mat/0102322>.
- [8] T. Yamano. Source Coding Theorem based on a Nonadditive Information Content. *Physica A*, **305**,1, pp. 190-195, 2002.
- [9] T. Cover and J. Thomas. *Elements of Information Theory*. John Wiley & Sons, New York, NY, 1991.
- [10] T. Berger. *Rate Distortion Theory*. Prentice-Hall, Englewood Cliffs, NJ, 1971.
- [11] N. Tishby, F. C. Pereira, W. Bialek. The Information Bottleneck Method. *Proceedings of the 37th Annual Allerton Conference on Communication, Control and Computing*, pp 368-377, 1999.
- [12] R. C. Venkatesan. Generalized Statistics Framework for Rate Distortion Theory. *Physica A*, To Appear, 2007. Preliminary manuscript available at <http://arxiv.org/abs/cond-mat/0611567>.
- [13] R. E. Blahut. Computation of Channel Capacity and Rate Distortion Functions. *IEEE Trans. on Inform. Theory*, **IT18**, pp. 460473, 1972.
- [14] K. Rose. A Mapping Approach to Rate Distortion Computation and Analysis. *IEEE Trans. on Inform. Theory*, **40**,6 pp. 19391952, 1994.
- [15] A. Banerjee, S. Merugu, I. Dhillon, and, J. Ghosh. Clustering with Bregman Divergences. *Journal of Machine Learning Research (JMLR)*, **6**, pp 1705-1749, 2005.
- [16] A. Banerjee. Scalable Clustering Algorithms *Ph.D. dissertation, The University of Texas at Austin, 2005*.
- [17] J. Naudts. Continuity of a Class of Entropies and Relative Entropies. *Rev. Math. Phys.*, **16**, pp. 809-824 , 2004. Manuscript available at <http://arxiv.org/abs/cond-mat/0208038>.
- [18] G. L. Ferri, S. Martinez, and A. Plastino. Equivalence of the Four Versions of Tsallis Statistics. *J. Stat. Phys.*, **04**, pp. 04009-04024 , 2004. Manuscript available at <http://arxiv.org/abs/cond-mat/0503441>.




Fig. 1. Rate Distortion Curves for GBRD Model and Extensive RD with Bregman Divergences




Fig. 2. Curves for β v/s β^* for GBRD Model