

Comment on “Dynamical Heterogeneities in a Supercooled Lennard-Jones Liquid”

In two recent interesting letters [1] evidence was presented for the existence of a growing dynamic correlation length when we approach the glass transition from the liquid phase (a similar divergence is present also in the off-equilibrium dynamics below T_c [2]). Here we would like to point out that this phenomenon can be easily predicted using the replica approach of reference [3]. Let us call $\rho(x, t)$ the density at time t ($\rho(x, t) = \sum_i \delta(x_i(t) - x)$). It is convenient to consider the quantity $Q_{z_1}(x, t) = \rho(x - z_1/2, 0)\rho(x + z_1/2, t)$. Using it we can construct the correlation $C(x - y, t) = \langle Q_{z_1}(x, t)Q_{z_2}(y, t) \rangle$ that should be weakly depend on the values of z_1 and z_2 for $|z_1|, |z_2| \ll x - y$. The data in [1] can be read as evidence for the existence of a correlation length that controls the decay of C at large spatial argument. This length reaches a maximum at a time t comparable with the relaxation time of the system and this time grows for lower temperatures. Let us take the mode coupling point of view, assuming an incipient divergence of the relaxation time and breaking of ergodicity at a temperature T_c . Exactly at the mode coupling transition we would sample at large times a single ergodic component and obtain

$$C_\infty(x) = \lim_{t \rightarrow \infty} C(x - y, t) = \langle \rho(x + z_1/2)\rho(y + z_2/2) \rangle_c \langle \rho(x - z_1/2)\rho(y - z_2/2) \rangle_c \quad (1)$$

where the expectation value is taken inside the same ergodic component and c stands for connected. As soon as we approach T_c and the times become large, the dynamic correlation function goes, in the appropriate time window, to the critical one. Our burden is to prove that there are (within the approximation used) long range correlations at the mode coupling point.




FIG. 1. We plot $\partial^2 W / \partial \epsilon^2$ as a function of the coupling ϵ for the model in [3] and $T = T_c(1 - 0.1)$ (dashed line), $T = T_c$ (full line) and $T = T_c(1 + 0.5)$ (dotted line). Notice the divergence of the $\partial^2 W / \partial \epsilon^2$ for $\epsilon \rightarrow 0$ for $T = T_c$ in the high branch of the curve.

In [3] it was shown that the mode coupling transition

may be characterized as follows. Consider a system that thermalizes at temperature T with an attractive coupling ϵ with a reference configuration which is itself a fixed equilibrium configuration at temperature T . We call W the free-energy as a function of ϵ . There are three possibilities for the expectation value of $(\partial W / \partial \epsilon)|_{\epsilon=0}$: (a) A non zero stable value below the ideal glass transition ($T < T_K$) (b) Flatly a zero value above the Mode Coupling transition ($T > T_c$) (c) A metastable non zero value in the region $T_K < T < T_c$. The metastable state exists also for non zero coupling and is destabilized on the spinodal line. At T_c the spinodal point is at $\epsilon = 0$. The situation is sketched in fig 1. It is a general feature of spinodal points that the second derivative of the free energy, and its associated correlation length are divergent. This is precisely the length associated to $C_\infty(x)$. The learned reader may notice that we are just finding what one should expect from the “principle of marginal stability” [5] which states that the mode coupling transition point is critical. Similar conclusions follow also from the geometrical description of [4]. It should also be evident that in principle this correlation length could be computed directly in the mode coupling approach. Unfortunately such a computation is not so simple and the method of [3] provides a useful short cut. We stress that the divergence is present only in the average of the product of two correlation functions inside the same ergodic component. No divergent length would be seen in equilibrium correlation functions, whose dominant contribution comes from configurations in different ergodic components.

-
- [1] W.Kob, C.Donati, S.J.Plimpton, P.H. Poole and S.C. Glotzer, Phys. Rev. Lett. **79** (1997) 2827. C. Donati, J.F. Douglas, W. Kob, S.J. Plimpton, P.H. Poole and S.C. Glotzer Phys. Rev. Lett. **80** (1998) 2338
 - [2] G. Parisi, cond-mat/9801034
 - [3] S. Franz, G. Parisi, Phys. Rev. Lett. **79** (1997) 2486, and cond-mat/9711215 (Physica A in press).
 - [4] J. Kurchan and L. Laloux J. Phys. A: Math. Gen. **29** (1996) 1929-1948
 - [5] A. Crisanti, H. Horner and H.J. Sommers, Z. Phys. B **92** 257 (1993)

Silvio Franz(*) and Giorgio Parisi(**)

(*) Abdus Salam ICTP Strada Costiera 11, P.O. Box 563, 34100 Trieste (Italy)

(**) Università di Roma “La Sapienza” Piazzale A. Moro 2, 00185 Rome (Italy)