Geometrization of perfect fluid in 5-D Kaluza-Klein theory.

Sergey S. Kokarev* 150000, Russia, Yaroslavl, YSPU, r.409

Abstract

General formulation of geometrization matter problem by scalar field $\varphi = \sqrt{-G_{55}}$ with the help of possibilities of classical 5-D Kaluza-Klein theory is given. Mathematical integrability conditions for such geometrization for the case of perfect fluid are derived.

1 Introduction

When GR had been built and gravitational interactions within its frame had been geometrized, Einstein himself considered this problem as half solved: energy-momentum tensor, that appeared in righthand part of field equations, in his mind, would had geometrical nature too [1].

Such a possibility naturally appears whithin the frame of multidimensional geometrical models type of Kaluza-Klein theory, of which the first (5-dimensional) version has been proposed by T.Kaluza in 1921 [2, 3]. Extracomponents, given by "scalar" sector of multidimensional metric have been used for a number of purposes. In present article the possibility of description of matter sources in terms of geometrical scalar field $\varphi = \sqrt{-G_{55}}$ in effective 4-D space-time is analyzed. In present time such an approach is worked out by Wesson and others (see Ref. in [7, 8, 9, 10]), but some steps to the problem have been made by the number of authors earlier [4, 5].

2 Statement of the problem

The starting point of our investigation is vacuum 5-D Einstein equations, that we shall write in the form:

$${}^{5}R_{AB} - \frac{1}{2}G_{AB}{}^{5}R = 0, \tag{1}$$

where ${}^5R_{AB}$ – 5-D Ricci tensor, 5R – 5-D curvature scalar, G_{AB} – 5-D metric, which in 4-D presentation has the following kind:

$$G_{AB} = \begin{pmatrix} \tilde{g}_{\mu\nu} - (4k/c^4)\varphi^2 A_{\mu} A_{\nu} & (2\sqrt{k}/c^2)\varphi^2 A_{\mu} \\ \hline (2\sqrt{k}/c^2)\varphi^2 A_{\nu} & -\varphi^2 \end{pmatrix}$$
 (2)

Expression (2) for G_{AB} can be derived with the help of method of 1+4-splitting of 5-D manifold, which is outlined in monography [5]. "Vector" sector $G_{5\mu}$ is identified

^{*}e-mail: sergey@yspu.yaroslavl.ru

with an electromagnetic potential: $G_{5\mu} = (2\sqrt{k}/c^2)\varphi^2 A_{\mu}$, that lets us consider 5-D Kaluza-Klein models as unified theory of gravitational and electromagnetic interactions.

Under the identification of metric of 4-D space-time section $\tilde{g}_{\mu\nu}$ with observable 4-D metric $g_{\mu\nu}$ there is a possibility of conformal transformation of the form:

$$\tilde{g}_{\mu\nu} = F(\varphi)g_{\mu\nu},\tag{3}$$

where $F(\varphi)$ – arbitrary function of scalar field.

For identification of components of multidimensional metric with 4-D values it is necessary to impose the following conditions:

$$\frac{\partial \tilde{g}_{\mu\nu}}{\partial x^5} = 0, \qquad \text{(cylindricity condition)} \tag{4}$$

and restrictions on coordinate transformations:

$$x'^{\mu} = x'^{\mu}(x^0, x^1, x^2, x^3); \tag{5}$$

$$x^{5} = x^{5} + f(x^{0}, x^{1}, x^{2}, x^{3}).$$
(6)

Such transformations on the one hand conserve cylindricity conditions (4) and 4-D covariancy of $\tilde{g}_{\mu\nu}$, A_{μ} , φ , and, on the other hand, are 4-D general coordinate transformations – (5) and gauge transformations of vector potentials A_{μ} – (6).

The method of 1+4-splitting lets us split 5-D Einstein equations (1) in a ten 4-D Einstein equation:

$${}^{4}\tilde{R}_{\mu\nu} - 1/2\tilde{g}_{\mu\nu}{}^{4}\tilde{R} = \varphi^{2}\kappa T_{\mu\nu}^{(em)} + \frac{1}{\varphi}(\varphi_{;\tilde{\nu};\tilde{\mu}} - \tilde{g}_{\mu\nu}\tilde{\nabla}^{2}\varphi); \tag{7}$$

four equations, corresponding to Maxwell theory:

$$\tilde{F}^{\mu\nu}_{;\tilde{\nu}} - \frac{2\varphi_{,\alpha}}{\varphi} \tilde{F}^{\alpha\mu} = 0 \tag{8}$$

and the 15-th equation, that connects invariants of gravitational and electromagnetic fields:

$${}^{4}\tilde{R} + \frac{3}{8\pi}\kappa\varphi^{2}\tilde{F}_{\alpha\beta}\tilde{F}^{\alpha\beta} = 0. \tag{9}$$

In (7)—(9) values with "tilde" are related to starting metric $\tilde{g}_{\mu\nu}$.

In present article all investigations are carried out, at first, without an electromagnetic field $(F_{\mu\nu}=0)$, and secondly, conformal factor in (3) is taken in the form:

$$F(\varphi) = \varphi^{2n},\tag{10}$$

where n – arbitrary real constant. Then equations (7) and (9) with account of conformal transformation (10) can be put to the form:

$${}^{4}R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}{}^{4}R = (1+2n)\phi_{;\mu;\nu} - (2n^{2}+2n-1)\phi_{,\mu}\phi_{,\nu} -$$
(11)

$$-g_{\mu\nu}((1+2n)\nabla^2\phi + (n^2+n+1)(\nabla\phi)^2);$$

$$n\nabla^2\phi + n^2(\nabla\phi)^2 - 1/6^4R = 0, (12)$$

where $\phi = \ln \varphi$. Equations (8) are satisfied identically.

So, when all outlined above conditions are satisfied, then vacuum Einstein equations (1) (after 1+4-splitting procedure) take the form of nonvacuum 4-D Einstein equations (11), (12).

Let us assume now, that we have some exact solutions of 4-D nonvacuum Einstein equations with energy-momentum tensor of perfect fluid in right-hand $side(\kappa = 1)$:

$$R_{\mu\nu} - 1/2g_{\mu\nu}R = (p+\varepsilon)u_{\mu}u_{\nu} - pg_{\mu\nu} \equiv T_{\mu\nu}^{(hd)}$$
(13)

If scalar field ϕ and constant n are so, that

$$(p+\varepsilon)u_{\mu}u_{\nu} - pg_{\mu\nu} = \tag{14}$$

$$(1+2n)\phi_{;\mu;\nu} - (2n^2+2n-1)\phi_{\mu}\phi_{\nu} - g_{\mu\nu}((1+2n)\nabla^2\phi + (n^2+n+1)(\nabla\phi)^2).$$

and equation (12) is satisfied, then we can say, that such $T_{\mu\nu}^{(hd)}$ can be geometrized within the frame of 5-D Kaluza-Klein theory and 4-D metric, generated by this sources, has a purely 5-D origin. Below we shall derive restrictions on the geometry of 4-D space-time and movement of matter there, under which the equation (14) is satisfied.

3 Integrability conditions

Equations (14) can be considered as a system of differential equations in partial derivatives of the second order for components of the gradient $\phi_{,\mu}$, taking $u_{\mu}, \varepsilon, p, g_{\mu\nu}$ as prescribed functions. For integrability analysis of this system it is convenient to rewrite it in following more general form:

$$\phi_{:\mu:\nu} = k\phi_{\mu}\phi_{\nu} + au_{\mu}u_{\nu} + bg_{\mu\nu},\tag{15}$$

where:

with $g_{\mu\nu}$ equation (11).

$$k = -\frac{k_2}{k_1}; \quad a = \frac{p+\varepsilon}{k_1};$$

$$b = \frac{-p - k_3(\nabla\phi)^2}{k_1} = c + \overline{k}(\nabla\phi)^2;$$

$$c = -\frac{p}{k_1}; \quad \overline{k} = -\frac{k_3}{k_1};$$

$$(16)$$

 $k_1 = 1 + 2n \neq 0;$ $k_2 = -2n^2 - 2n + 1;$ $k_3 = 3n^2 + 3n$

The equation (15) can be derived from (14) with using the relation:

$$\nabla^2 \phi = -(1+2n)(\nabla \phi)^2.$$

which, in turn, follows from linear combination of equation (12) and contracted

(17)

Integrability conditions for system (18) imply identical fulfillment of the relationship

$$2\phi_{;\mu;[\nu;\lambda]} = R^{\sigma}_{\cdot\mu\nu\lambda}\phi_{\sigma},\tag{18}$$

if into the left-hand side expressions from right-hand side of (15) are inserted, with replacement of second covariant derivatives $\phi_{;\mu;\nu}$ through (15) again. Identical fulfillment (18) is necessary for the existence of a scalar field ϕ , satisfied the equation (14).

Omitting intermediate calculations, we perform (18) in the form:

$$R^{\sigma}_{\cdot \mu\nu\lambda}\phi_{\sigma} = u_{\mu}(u_{\nu}a_{\lambda} - u_{\lambda}a_{\nu}) + kau_{\mu}(u_{\lambda}\phi_{\nu} - u_{\nu}\phi_{\lambda}) + (c_{\lambda}g_{\mu\nu} - c_{\nu}g_{\mu\lambda}) +$$
(19)

$$\xi_1(\phi_{\lambda}g_{\mu\nu} - \phi_{\nu}g_{\mu\lambda}) + \xi_2(u_{\lambda}g_{\mu\nu} - u_{\nu}g_{\mu\lambda}) + a(u_{\mu;\lambda}u_{\nu} - u_{\mu;\nu}u_{\lambda} + u_{\mu}(u_{\nu;\lambda} - u_{\lambda;\nu}))$$

where
$$\xi_1 = 2\overline{k}(k(\nabla\phi)^2 + b) - kb; \quad \xi_2 = 2\overline{k}a(\nabla\phi \cdot \vec{u})$$
 (20)

4 Tetrads formalism

For obtaining of invariant form of integrability condition (15) it is convenient to go to tetrad component of all tensors. As a tetrad basis we take Lorentz tetrad $\vec{t}, \vec{x}, \vec{y}, \vec{z}$, The first of noted vectors can be identified with 4-velocity of matter \vec{u} :

$$q = \vec{u} \otimes \vec{u} - \vec{x} \otimes \vec{x} - \vec{y} \otimes \vec{y} - \vec{z} \otimes \vec{z}. \tag{21}$$

Curvature tensor $R_{\mu\nu\lambda\sigma}$ is represented in such basis in the following kind:

$$R_{\mu\nu\lambda\sigma} = a_{ij} X_{\mu\nu}^{(i)} X_{\lambda\sigma}^{(j)} \quad i, j = \overline{1, 6}, \tag{22}$$

where $X_{\mu\nu}^{(i)}$ – simple bivectors, that forms the basis of 6-dimensional linear bivector space and are built from basic tetrad vectors:

$$X_{\mu\nu}^{(1)} = 2u_{[\mu}x_{\nu]}; \quad X_{\mu\nu}^{(2)} = 2u_{[\mu}y_{\nu]}; X_{\mu\nu}^{(3)} = 2u_{[\mu}z_{\nu]}; \quad X_{\mu\nu}^{(4)} = 2x_{[\mu}y_{\nu]}; X_{\mu\nu}^{(5)} = 2z_{[\mu}x_{\nu]}; \quad X_{\mu\nu}^{(6)} = 2y_{[\mu}z_{\nu]}.$$

$$(23)$$

Matrix a_{ij} – is symmetrical and its elements are tetrad components of curvature tensor, for example:

$$R_{\mu\nu\lambda\sigma}u^{\mu}x^{\nu}u^{\lambda}x^{\sigma} = R_{(0)(1)(0)(1)} = a_{11} \quad \dots$$
 (24)

Einstein equations (13) in tetrad form take the following kind:

$$a_{35} = a_{24};$$
 $a_{14} = a_{36};$ $a_{26} = a_{15};$ $a_{12} = -a_{56};$ $a_{13} = -a_{46};$ $a_{23} = -a_{45};$ (25)

$$-a_{44} - a_{55} - a_{66} = \varepsilon;$$
 $a_{66} - a_{22} - a_{33} = p;$ $a_{55} - a_{11} - a_{33} = p;$ $a_{44} - a_{11} - a_{22} = p.$

In the right-hand side of (15) all tensors can be decomposed by a tetrad basis as follows:

$$\vec{\nabla}\phi = \phi_0 \vec{u} + \phi_1 \vec{x} + \phi_2 \vec{y} + \phi_3 \vec{z}; \tag{26}$$

$$\vec{\nabla}\varepsilon = \varepsilon_0 \vec{u} + \varepsilon_1 \vec{x} + \varepsilon_2 \vec{y} + \varepsilon_3 \vec{z}; \tag{27}$$

$$\vec{\nabla}a = \frac{1 + p_{\varepsilon}}{k_1}\vec{\varepsilon}; \quad \vec{\nabla}c = -\frac{p_{\varepsilon}}{k_1}\vec{\nabla}\varepsilon. \tag{28}$$

Here $p_{\varepsilon} = dp/d\varepsilon$.

For decomposition of the covariant derivative of vector u_{μ} we use the well known expression [5]:

$$u_{;\mu;\nu} = F_{\mu}u_{\nu} + \omega_{\mu\nu} + \sigma_{\mu\nu} - \frac{\theta}{3}h_{\mu\nu},$$
 (29)

where

$$F_{\mu} = u_{\mu;\nu} u^{\nu}$$

- the acceleration vector of a comoving reference frame, $F_{\nu}u^{\nu}=0$;

$$\omega_{\mu\nu} = u_{[\mu;\nu]} + u_{[\mu}F_{\nu]} \tag{30}$$

– the antisymmetric tensor of angular velocity of the reference frame, $\omega_{\mu\nu}u^{\nu}=0$;

$$\sigma_{\mu\nu} = u_{(\mu;\nu)} - u_{(\mu}F_{\nu)} + \frac{\theta}{3}h_{\mu\nu} \tag{31}$$

– the symmetric shear tensor (traceless part of strain tensor) of the reference frame, $\sigma_{\mu\nu}u^{\nu}=0$;

$$\theta = u^{\mu}_{;\mu}; \tag{32}$$

- the stretch scalar of the reference frame (trace of strain tensor);

$$h_{\mu\nu} = u_{\mu}u_{\nu} - g_{\mu\nu} \tag{33}$$

- the metric of the local 3-D space section, orthogonal to u_{μ} .

Tensors of the reference frame F, ω σ in a chosen tetrad basis are given by following expressions:

$$\vec{F} = F_1 \vec{x} + F_2 \vec{y} + F_3 \vec{z}; \tag{34}$$

$$\omega_{\mu\nu} = \omega_4 X_{\mu\nu}^{(4)} + \omega_5 X_{\mu\nu}^{(5)} + \omega_6 X_{\mu\nu}^{(6)}; \tag{35}$$

$$\sigma_{\mu\nu} = \sigma_2 Y_{\mu\nu}^{(2)} + \sigma_3 Y_{\mu\nu}^{(3)} - (\sigma_2 + \sigma_3) Y_{\mu\nu}^{(4)} + \sigma_8 Y_{\mu\nu}^{(8)} + \sigma_9 Y_{\mu\nu}^{(9)} + \sigma_{10} Y_{\mu\nu}^{(10)}$$
(36)

In the last expression $Y_{\mu\nu}^{(i)}$ – diadic basis of 10-dimensional linear space of symmetric tensors, formed by basic vectors:

$$Y_{\mu\nu}^{(1)} = u_{\mu}u_{\nu}; Y_{\mu\nu}^{(2)} = x_{\mu}x_{\nu}; Y_{\mu\nu}^{(3)} = y_{\mu}y_{\nu}; Y_{\mu\nu}^{(4)} = z_{\mu}z_{\nu}; Y_{\mu\nu}^{(5)} = 2u_{(\mu}x_{\nu)}; Y_{\mu\nu}^{(6)} = 2u_{(\mu}y_{\nu)}; Y_{\mu\nu}^{(7)} = 2u_{(\mu}z_{\nu)}; Y_{\mu\nu}^{(8)} = 2x_{(\mu}y_{\nu)}; Y_{\mu\nu}^{(9)} = 2x_{(\mu}z_{\nu)}; Y_{\mu\nu}^{(9)} = 2y_{(\mu}z_{\nu)}. (37)$$

The tensor $h_{\mu\nu}$ has the following form:

$$h = \vec{x} \otimes \vec{x} + \vec{y} \otimes \vec{y} + \vec{z} \otimes \vec{z}. \tag{38}$$

Integrability conditions (15), being rewritten in tetrad form are invariant under 4-D general coordinates transformations. Freedom in the choosing of tetrad basic vectors is reduced to local space rotations of triad $\vec{x}, \vec{y}, \vec{z}$, and can be used for the simplification of the obtained in the next section integrability equations. For the most generality we suppose that triad $\vec{x}, \vec{y}, \vec{z}$ is arbitrary oriented.

5 Integrability conditions in tetrads representation.

Equating scalar coefficient in (15) under the similar combinations of basic vectors in right-hand and in the left-hand sides we get the following system of scalar equations, that is equivalent to the original system (15):

$$a_{11}\phi_1 + a_{12}\phi_2 + a_{13}\phi_3 = e_1 + \lambda\phi_1 - f_1;$$

$$\begin{aligned} a_{12}\phi_1 + a_{22}\phi_2 + a_{23}\phi_3 &= e_2 + \lambda\phi_2 - f_2; \\ a_{13}\phi_1 + a_{23}\phi_2 + a_{33}\phi_3 &= e_3 + \lambda\phi_3 - f_3; \\ a_{14}\phi_1 + a_{24}\phi_2 + a_{34}\phi_3 &= 2W_4; \\ a_{15}\phi_1 + a_{25}\phi_2 + a_{35}\phi_3 &= 2W_5; \\ a_{16}\phi_1 + a_{26}\phi_2 + a_{36}\phi_3 &= 2W_6; \\ a_{11}\phi_0 + a_{14}\phi_2 - a_{15}\phi_3 &= \Delta_2 + S; \\ a_{12}\phi_0 + a_{24}\phi_2 - a_{25}\phi_3 &= \Delta_8 + W_4; \\ a_{13}\phi_0 + a_{34}\phi_2 - a_{35}\phi_3 &= \Delta_9 - W_5; \\ a_{14}\phi_0 + a_{44}\phi_2 - a_{45}\phi_3 &= p_e e_2 - \xi_1\phi_2; \\ a_{15}\phi_0 + a_{45}\phi_2 - a_{55}\phi_3 &= -p_e e_3 + \xi_1\phi_3; \\ a_{16}\phi_0 + a_{46}\phi_2 - a_{56}\phi_3 &= 0; \\ a_{12}\phi_0 - a_{14}\phi_1 + a_{16}\phi_3 &= \Delta_8 - W_4; \\ a_{22}\phi_0 - a_{24}\phi_1 + a_{26}\phi_3 &= \Delta_3 + S; \\ a_{23}\phi_0 - a_{34}\phi_1 + a_{36}\phi_3 &= \Delta_{10} + W_6; \\ a_{24}\phi_0 - a_{44}\phi_1 + a_{46}\phi_3 &= -p_e e_1 + \xi_1\phi_1; \\ a_{25}\phi_0 - a_{45}\phi_1 + a_{56}\phi_3 &= 0; \\ a_{26}\phi_0 - a_{46}\phi_1 + a_{66}\phi_3 &= p_e e_3 - \xi_1\phi_3; \\ a_{13}\phi_0 + a_{15}\phi_1 - a_{16}\phi_2 &= \Delta_9 + W_5; \\ a_{23}\phi_0 + a_{25}\phi_1 - a_{26}\phi_2 &= \Delta_{10} - W_6; \\ a_{33}\phi_0 + a_{35}\phi_1 - a_{36}\phi_2 &= \Delta_4 + S; \\ a_{34}\phi_0 + a_{45}\phi_1 - a_{46}\phi_2 &= 0; \\ a_{35}\phi_0 + a_{55}\phi_1 - a_{56}\phi_2 &= p_e e_1 - \xi_1\phi_1; \\ a_{36}\phi_0 + a_{56}\phi_1 - a_{66}\phi_2 &= -p_e e_2 + \xi_1\phi_2. \end{aligned}$$

The obtained equations connect components of a curvature tensor of 4-D space with a characteristics of motion of a matter there. Besides, this connections are additional to Einstein equations (13). If space-time allows the identical fulfillment of the obtained integrability equations and scalar field satisfy the equation (12), then this space-time has 5-D nature and 5-dimensionally geometrized matter.

(39)

Here

 $S = \frac{-p_{\varepsilon}}{k_1} \varepsilon_0 + \xi_1 \phi_0 + \xi_2 - a\theta/3.$

6 Example: potential motion in the Friedman flat cosmological model.

Let us suppose, that expansion $\phi_{,\mu}$ over basic tetrad vectors has the following form:

$$\phi_{,\mu} = \phi_0 u_\mu. \tag{40}$$

This means, that a vector field u_{μ} is orthogonal to hypersurface $\phi = const.$ For simplicity we restrict our attention to the case of motion in the Friedman flat cosmological model:

$$ds^{2} = dt^{2} - e^{2\lambda}(dx^{2} + dy^{2} + dz^{2}). \tag{41}$$

Nonzero components of curvature tensor are:

$$a_{11} = a_{22} = a_{33} = \ddot{\lambda} + \dot{\lambda}^2, \quad -a_{44} = -a_{55} = -a_{66} = \dot{\lambda}^2.$$
 (42)

Tetrad Einstein equations (25) take the following form:

$$3\dot{\lambda}^2 = \varepsilon; \quad -2\ddot{\lambda} - 3\dot{\lambda}^2 = p. \tag{43}$$

4-velocity u_{μ} has the following components:

$$u_0 = 1, \quad u_1 = u_2 = u_3 = 0.$$
 (44)

From the tensors characteristics of the reference frame only stretch coefficient is nonzero:

$$\theta = 3\dot{\lambda}.\tag{45}$$

Integrability conditions of the previous section are reduced to only:

$$a_{11}\phi_0 = S.$$
 (46)

Taking into account that $\phi_0 = \dot{\phi}$, writing out abbreviated designations by formulae (20) and (39) and expressing $\dot{\varepsilon}$ from Einstein equations (43), condition (46) can be put to the form:

$$\mu_1 \dot{\phi}^3 + \dot{\phi}(\dot{\lambda}^2(\mu_2 - 1) - \ddot{\lambda}) - \mu_3 \dot{\lambda} \ddot{\lambda} = 0, \tag{47}$$

where

$$\mu_1 = \overline{k}(k+2\overline{k}); \quad \mu_2 = 3(2\overline{k}/k_1 + p_{\varepsilon}k/k_1); \quad \mu_3 = 2(3p_{\varepsilon} - 1)/k_1$$

and state equation is taken in the following form:

$$p = p_{\varepsilon}\varepsilon, \quad p_{\varepsilon} = const.$$
 (48)

Equation (12), that in metric (41) has the form:

$$\ddot{\phi} + 3\dot{\lambda}\dot{\phi} + (1+2n)\dot{\phi}^2 = 0 \tag{49}$$

has the following first integral:

$$\dot{\phi} = \frac{e^{-3\lambda}}{\overline{C} + (1+2n) \int e^{-3\lambda} dt},\tag{50}$$

where \overline{C} – is a constant integration, of which general solution can be easily obtained:

$$\phi = \frac{1}{1+2n} \ln(\overline{C} + (1+2n) \int e^{-3\lambda} dt) + C_1$$
 (51)

With the substitution of $\dot{\phi}$ from (50) to (49) the expression in the left-hand must be equal zero identically.

Einstein equations for a chosen class state equation (48) can be easily integrated:

$$\lambda = \begin{cases} (2/3(1+p_{\varepsilon})) \ln[(3/2)(1+p_{\varepsilon})t + \overline{C}_0], & p_{\varepsilon} > -1; \\ C_0 t, & p_{\varepsilon} = -1. \end{cases}$$

For $p_{\varepsilon} = -1$ $(p + \varepsilon = 0)$, omitting all intermediate calculations, we obtain that (46) identically satisfied under n = 1, -2. In this case $\phi = C_0 t$. Case $p_{\varepsilon} = 1$ $(p = \varepsilon)$ will be considered in subsection (3) of Conclusion. For $p_{\varepsilon} \neq \pm 1$ we obtain the following square equation for the index of conformal transformation n:

$$\alpha_1 n^2 + \alpha_1 n + \alpha_2 = 0, (52)$$

where

$$\alpha_1 = 27p_{\varepsilon}^3 + 63p_{\varepsilon}^2 + 33p_{\varepsilon} + 5; \tag{53}$$

$$\alpha_2 = 18p_{\varepsilon}^2 + 24p_{\varepsilon} - 10.$$

Its solution is:

$$n = \frac{3p_{\varepsilon}(\sigma - 1) - (3\sigma + 1)}{2(3p_{\varepsilon} + 1)}, \quad \sigma = \pm 1$$
(54)

For dust $(p_{\varepsilon} = 0)$ we have n = 1, -2, for radiation $(p_{\varepsilon} = 1/3)$ n = 0, -1.

Corresponding 5-D vacuum metrics that in all cases can be obtained by the inverse conformal transformation are only of the following two types:

$$dI_1^2 = dt^2 - dx^2 - dy^2 - dz^2 - t^2(dx^5)^2; (55)$$

$$dI_2^2 = dt^2 - t(dx^2 + dy^2 + dz^2) - \frac{1}{t}(dx^5)^2$$
(56)

Its relations with considered 4-D metrics are shown in the table:

	$p_{\varepsilon} = -1$	$p_{\varepsilon} = 0$	$p_{\varepsilon} = 1/3$
(1)	n = 1	n = -2	n = -1
(2)	n = -2	n = 1	n = 0

Cases with n=1 have been considered in [11, 12], the case with n=0 have been analyzed in Wesson's work [7] .

7. Conclusion

Completing our account we make some remarks:

1. The approach, proposed here, can be called "4-dimensional", because 4-D metric and an energy-momentum tensor are given. From integrability conditions scalar field ϕ and index n can be found, or, that is the same, can be found 5-D vacuum space-time which geometrizes beforehand given matter. As an example of another approach, "5-dimensional", we can consider the approach of Wesson, where known 5-D vacuum solution is used. 5-D Einstein equations are split in to parts: one - 4-D Einstein tensor, other - a combination

of derivatives of the scalar field, which is declared as an effective energy-momentum tensor of induced matter. Type of this tensor in such approach is, in general, arbitrary. In both [8, 9] obtained energy-momentum tensor was anisotropic. All investigations in Wesson's group are carried out under n=0. Note, that using of the conformal transformation alleviates statement made in [10]: there is no rigid necessity to introduce the 5-th coordinate into metric to obtain state equation for effective matter other than radiationlike.

- 2. The problem of the geometrization of matter in Kaluza-Klein theory has many formal analogies with the known problem of isometric embedding of 4-D Riemannian space into flat space of more dimensions [6].
- 3. The restriction on index $n \neq -1/2$ involves the fact that in this case the second derivatives of a scalar field in equations (14) vanish. Equations (14) became algebraic with respect to the gradient ϕ_{μ} and integrability conditions will be

$$\phi_{\mu,\nu} - \phi_{\nu,\mu} = 0,$$

where ϕ_{μ} is algebraically expressed through u_{μ} , p and ε . In the considered example a peculiar case $p_{\varepsilon} = 1$ $(p = \varepsilon)$ is realized namely under n = -1/2 from both 5-D metrics (55,56).

4. Cosmological models of open and closed types have been considered under n = 1 in [5](p.230-234) from the viewpoint of "5-dimensional" approach.

References

- [1] A. Einstein, Physic and reality, M. "Nauka", p.159 (1974) (In Russian)
- [2] T. Kaluza, Sitzungsber. d. Berl. Akad., 966-971 (1921).
- [3] Yu. S. Vladimirov, Space-time: explicit and hidden dimensions, M.Nauka, 1989 (In Russian)
- [4] E. Schmutzer, Exp. Techn, Phys., 28 pp.395-402, 499-508 (1980); 29 pp.129-136, 337-341, 463-480 (1981).
- [5] Yu. S. Vladimirov Reference frames in gravitation theory, M.Energoizdat, 1982 (In Russian).
- [6] L. P. Eisenhart, Riemannian geometry, Izd.in.lit., 1948 (In Russian).
- [7] P. S. Wesson, Astroph. Journal, **394** p.19-24 (1992).
- [8] P. S. Wesson, Astroph. Journal, **420** L49-L52 (1994).
- [9] P. S. Wesson, *Phys.Let.*, **B 276** 299-302 (1992).
- [10] P. S. Wesson, J. Ponce de Leon, P. Lim, H. Liu, Int. J. of Mod. Phys. D2 163-170 (1993).
- [11] S. S. Kokarev, Izv. VUZov (Physics), N1, p.111-117 (1995).(In Russian)
- [12] S. S. Kokarev, In abstracts of shool-ceminar "Multid. gravity and cosmology", Yaroslavl, 1994., p. 19. (In Russian).