

# On nonlinear parametric quantisation of gravity and cosmological models

Charles Wang

Department of Physics, Lancaster University

Lancaster LA1 4YB, UK

E-mail: c.wang@lancaster.ac.uk

## Abstract

A generalisation of the recently formulated nonlinear quantisation of a parameterised theory is presented in the context of quantum gravity. The parametric quantisation of a Friedmann universe with a massless scalar field is then considered in terms of the analytic solutions of the resulting evolution equations.

PACS numbers: 04.20.Fy, 04.60.Ds, 04.60.Kz, 11.10.Lm

In a recent paper [1] the quantisation of a physical system with finite degrees of freedom subject to a Hamiltonian constraint was re-examined. A new scheme, called parametric quantisation, was introduced in that work that advocates the idea of treating a chosen time variable as a constrained classical variable coupled to other dynamical variables to be quantised. The consideration was motivated by the cosmological approach to quantum gravity [2] whose dynamical structure is analogous to that of the parameterised theory of a relativistic particle in an  $(n+1)$ -dimensional pseudo-Riemannian spacetime with a metric tensor  $\gamma_{\mu\nu}$  in coordinates  $q^\mu$ ,  $\mu, \nu = 0, 1, \dots, n$ , described by a Lagrangian of the following generic form:

$$L(q^\mu, \dot{q}^\mu, N) = \frac{1}{2N} \gamma_{\mu\nu}(q^\lambda) \dot{q}^\mu \dot{q}^\nu - NV(q^\lambda). \quad (1)$$

Here  $q^\mu = q^\mu(t)$  described the particle trajectory parameterised by  $t$ , whose choice is related to the function  $N(t) > 0$ , and  $\dot{q}^\mu := \frac{dq^\mu}{dt}$ . Classically this generates the Lagrange's equations of motion

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}^\mu} \right) - \frac{\partial L}{\partial q^\mu} = 0 \quad (2)$$

subject to the Hamiltonian constraint

$$\mathcal{H}(q^\mu, \dot{q}^\mu, N) := -\frac{\partial L}{\partial N} = \frac{1}{2N^2} \gamma_{\mu\nu} \dot{q}^\mu \dot{q}^\nu + V = 0. \quad (3)$$

The equivalent canonical description is obtained from the “constrained Hamiltonian”:

$$H(q^\mu, p_\mu, N) = \dot{q}^\mu p_\mu - L = N\mathcal{H} \quad (4)$$

in terms of the conjugate momenta  $p_\mu := \frac{\partial L}{\partial \dot{q}^\mu} = \frac{1}{N} \gamma_{\mu\nu} \dot{q}^\nu$  where  $\mathcal{H}$  can be re-express as

$$\mathcal{H}(q^\mu, p_\mu) = \frac{1}{2} \gamma^{\mu\nu} p_\mu p_\nu + V. \quad (5)$$

The canonical equations of motion are given by

$$\frac{dq^\mu}{dt} = \frac{\partial H}{\partial p_\mu}, \quad \frac{dp_\mu}{dt} = -\frac{\partial H}{\partial q^\mu} \quad (6)$$

subject to the Hamiltonian constraint

$$\mathcal{H}(q^\mu, p_\mu) = 0. \quad (7)$$

A quantisation scheme may be set up by choosing one of the variables, say  $q^0$  with its conjugate momentum  $p_0$ , as a classical variable that interacts with the remaining quantised variables  $q^a, a = 1, 2, \dots, n$ , in a semi-classical fashion [1]:

$$i \frac{\partial \psi}{\partial t} = \hat{H} \psi \quad (8)$$

$$\frac{dq^0}{dt} = \left\langle \frac{\partial \hat{H}}{\partial p_0} \right\rangle, \quad \frac{dp_0}{dt} = - \left\langle \frac{\partial \hat{H}}{\partial q^0} \right\rangle \quad (9)$$

$$\langle \hat{\mathcal{H}} \rangle = 0 \quad (10)$$

where  $\psi = \psi(q^a, t)$  and  $\langle \hat{O} \rangle$  denotes the expectation value of any operator  $\hat{O}$ .<sup>1</sup> The operators  $\hat{H}$  and  $\hat{\mathcal{H}}$  are obtained by substituting  $p_a \rightarrow \hat{p}_a := -i \frac{\partial}{\partial q^a}$  into  $H$  and  $\mathcal{H}$  respectively followed by a suitable factor ordering. In this letter, equations (8), (9) and (10) will be cast into a more compact form that will prove to be advantageous in generalising parametric quantisation to a field theoretical framework. This reformulation is done by constructing an “unconstrained Hamiltonian”:

$$h(q^a, p_a, q^0, \dot{q}^0, N) := \dot{q}^a p_a - L = H - \dot{q}^0 \frac{\partial L}{\partial \dot{q}^0} \quad (11)$$

with a partial Legendre transformation of  $L$  by leaving out the “ $\dot{q}^0 \rightarrow p_0$ ” transform as performed in (5). With this the classical particle dynamics obeys the set of equations:

$$\frac{dq^a}{dt} = \frac{\partial h}{\partial p_a}, \quad \frac{dp_a}{dt} = -\frac{\partial h}{\partial q^a} \quad (12)$$

plus the Hamiltonian constraint taking the form:

$$\mathcal{H}(q^a, p_a, q^0, \dot{q}^0, N) := \frac{1}{N} \left( h + \dot{q}^0 \frac{\partial L}{\partial \dot{q}^0} \right) = 0. \quad (13)$$

Using the operators  $\hat{h}(q^a, \hat{p}_a, q^0, \dot{q}^0, N)$  and  $\hat{\mathcal{H}}(q^a, \hat{p}_a, q^0, \dot{q}^0, N)$  obtained by substituting  $p_a \rightarrow \hat{p}_a$  into  $h$  and  $\mathcal{H}$  given in (11) and (13) with suitable factor ordering, the system of equations (8), (9) and (10) now becomes

$$i \frac{\partial \psi}{\partial t} = \hat{h} \psi \quad (14)$$

$$\langle \hat{\mathcal{H}} \rangle = 0. \quad (15)$$

---

<sup>1</sup>Units in which  $c = \hbar = 16\pi G = 1$  are adopted throughout.

For any given positive  $N(t)$ , equations (14) and (15) constitute a system of evolution for the wavefunction  $\psi(q^a, t)$  and classical variable  $q^0(t)$  whose initial data may be specified arbitrarily. It is evident that the presence of (15) makes the quantum theory nonlinear. (See e.g. [6, 7] for other examples of nonlinear quantum theories.)

This formulation suggests a parametric quantisation of gravity whose formal treatment is outlined as follows. Start from the standard ADM Lagrangian for general relativity [5]:

$$L[g_{ij}, \dot{g}_{ij}, N_\mu] = \int N \sqrt{g} (K_{ij} K^{ij} - K^2 + R) d^3x \quad (16)$$

in terms of the 3-metric  $g_{ij}(x^k, t)$ ,  $g = \det(g_{ij})$ , lapse function  $N_0(x^k, t) := N(x^k, t)$ , shift vector  $N_i(x^k, t)$ , extrinsic curvature  $K_{ij}$ ,  $K = g^{ij} K_{ij}$  and intrinsic scalar curvature  $R$  of the evolving 3-geometry. ( $\mu = 0, 1, 2, 3; i, j, k = 1, 2, 3$ .) A natural way of isolating the “true” gravitational degrees of freedom is to transform from  $g_{ij}(x^k, t)$  to a set of embedding variables:  $\vartheta^\mu(x^k, t)$ ,  $\mu = 0, 1, 2, 3$ , and unconstrained variables:  $\varphi^a(x^k, t)$ ,  $a = 1, 2$  [3]. Here  $\vartheta^0$  specifies time slicing and  $\vartheta^a$  sets spatial coordinate condition. We then have  $L = L[\varphi^a, \varpi_a, \vartheta^\mu, \dot{\vartheta}^\mu, N_\mu]$ . The idea now is to regard  $\vartheta^\mu$  as constrained classical variables coupled “semi-classically” to the quantised true degrees of freedom carried by  $\varphi^a$ .

In view of the discussions above we introduce the unconstrained Hamiltonian

$$h[\varphi^a, \varpi_a, \vartheta^\mu, \dot{\vartheta}^\mu, N] := \int \dot{\varphi}^a \varpi_a d^3x - L \quad (17)$$

with the conjugate momenta

$$\varpi_a(x) := \frac{\delta L}{\delta \dot{\varphi}^a(x)}. \quad (18)$$

Conversely we may write  $L[\varphi^a, \varpi_a, \vartheta^\mu, \dot{\vartheta}^\mu, N_\mu] = \int \dot{\varphi}^a \varpi_a d^3x - h$  where  $\dot{\varphi}^a = \frac{\delta h}{\delta \varpi_a}$  and derive from which the super-Hamiltonian  $\mathcal{H} = \mathcal{H}^0$  and super-momenta  $\mathcal{H}^i$  as follows:

$$\mathcal{H}^\mu[\varphi^a, \varpi_a, \vartheta^\nu, \dot{\vartheta}^\nu, N_\nu; x] := - \frac{\delta L[\varphi^a, \varpi_a, \vartheta^\nu, \dot{\vartheta}^\nu, N_\nu]}{\delta N_i(x)} \quad (19)$$

where  $\mu, \nu = 0, 1, 2, 3; a = 1, 2, 3$ .

By analogy with (14) and (15) the nonlinear parametric quantisation of gravity may be formulated with the following system of equations:

$$i \frac{\partial \psi}{\partial t} = \hat{h} \psi \quad (20)$$

$$\langle \hat{\mathcal{H}}^\mu \rangle = 0. \quad (21)$$

that generates the nonlinear evolution of the wavefunction  $\psi[\varphi^a; t]$  and embedding variables  $\vartheta^\mu(x^k, t)$ , using the operators  $\hat{h}[\varphi^a, \hat{\varpi}_a, \vartheta^\mu, \dot{\vartheta}^\mu, N]$  and  $\hat{\mathcal{H}}^\mu[\varphi^a, \hat{\varpi}_a, \vartheta^\nu, \dot{\vartheta}^\nu, N_\nu; x]$  in terms of  $\hat{\varpi}_a = -i \frac{\delta}{\delta \varphi^a}$  with some suitable factor ordering.

It is an ongoing research programme to further investigate the quantisation of gravity using (20) and (21). The rest of this letter will be devoted to the parametric quantisation of a Friedmann universe with a massless scalar field based on the much simpler set of equations (14) and (15). Such a cosmological model has the Lagrangian [1, 4]:

$$L(\phi, \dot{\phi}, R, \dot{R}, N) = -\frac{6R}{N} \dot{R}^2 + \frac{R^3}{2N} \dot{\phi}^2 + 6NKR \quad (22)$$

in terms of the massless scalar field  $\phi(t)$ , scale factor  $R(t)$  and lapse function  $N(t)$  that enter into the Robertson-Walker metric

$$g = -N^2 dt^2 + R^2 \sigma. \quad (23)$$

Here, as usual,  $\sigma$  is the metric on the homogeneous and isotropic 3-space of constant curvature  $K$ , with  $K = 1, 0, -1$  corresponding to the closed, flat and open cases respectively. Regarding  $R$  as the classical time variable we obtain from (11) and (13)

$$h = 6R \frac{\dot{R}^2}{N} + \frac{Np^2}{2R^3} - 6NKR \quad (24)$$

$$\mathcal{H} = -6R \frac{\dot{R}^2}{N^2} + \frac{p^2}{2R^3} - 6KR \quad (25)$$

where

$$p := \frac{\partial L}{\partial \dot{\phi}} = \frac{R^3}{N} \dot{\phi}. \quad (26)$$

Comparing with the generic Lagrangian in (1) for  $n = 1$  we see that the current model has the nonzero metric components and potential as follows:

$$\gamma_{00} = -12R, \quad \gamma_{11} = R^3 \quad (27)$$

$$V = -6KR. \quad (28)$$

By substituting  $p \rightarrow \hat{p} := -1 \frac{\partial}{\partial \phi}$  into  $h$  the operators

$$\hat{h} = -\frac{N}{2R^3} \frac{\partial^2}{\partial \phi^2} + 6R \frac{\dot{R}^2}{N} - 6NKR \quad (29)$$

$$\hat{\mathcal{H}} = -\frac{1}{2R^3} \frac{\partial^2}{\partial \phi^2} - 6R \frac{\dot{R}^2}{N^2} - 6KR \quad (30)$$

are constructed that will act on a wavefunction  $\psi(\phi, t)$  (of weight  $\frac{1}{2}$  with respect to the 1-dimensional metric  $\gamma_{11}$ ) whose norm is normalised according to:

$$\langle \psi, \psi \rangle := \int_{-\infty}^{\infty} |\psi(\phi, t)|^2 d\phi = 1. \quad (31)$$

On parametric quantisation the evolution equations for the wavefunction  $\psi$  and classical variable  $R(t)$  follow from (14) and (15) to be: <sup>2</sup>

$$1 \frac{\partial \psi}{\partial t} = -\frac{N}{2R^3} \frac{\partial^2 \psi}{\partial \phi^2} \quad (32)$$

$$-6R \frac{\dot{R}^2}{N^2} + \frac{P^2}{2R^3} - 6KR = 0 \quad (33)$$

where

$$P^2 := -\left\langle \frac{\partial^2}{\partial \phi^2} \right\rangle. \quad (34)$$

We now solve these equations in the  $N = R$  gauge. First the wave equation (32) admits the following general solution:

$$\psi(\phi, t) = \int_{-\infty}^{\infty} \frac{A(k)}{\sqrt{2\pi}} e^{i[k\phi - F(t)k^2]} dk \quad (35)$$

---

<sup>2</sup>The term  $6R \frac{\dot{R}^2}{N^2} - 6KR$  in  $\hat{h}$  is dropped as it contributes only to an overall phase in  $\psi$ . However this term must be retained in applying (15).

where  $A(k)$  is an arbitrary complex-valued function and

$$F(t) := \int_{t_0}^t \frac{dt'}{2R(t')^2} \quad (36)$$

for a chosen reference time  $t_0$ . If at  $t = t_0$  the wavefunction is given by the normalised gaussian packet with a variance  $\delta_0$  of the form:

$$\psi_0(\phi) := \psi(\phi, t_0) = (2\pi)^{-\frac{1}{4}} \delta_0^{-\frac{1}{2}} e^{-\frac{(\phi-\phi_0)^2}{4\delta_0^2}} e^{ik_0(\phi-\phi_0)} \quad (37)$$

then

$$A(k) = (2\pi)^{-\frac{1}{4}} \sigma^{-\frac{1}{2}} e^{-\frac{(k-k_0)^2}{4\sigma^2}} e^{-ik\phi_0} \quad (38)$$

where  $k_0$  is the mean and  $\sigma = \frac{1}{2\delta_0}$  is the variance of the  $k$ -distribution (in the “momentum space”). Substituting (38) into (35) we obtain the following:

$$\psi(\phi, t) = (2\pi)^{-\frac{1}{4}} \left( \frac{\delta_0^2 + 1F(t)}{\delta_0} \right)^{-\frac{1}{2}} e^{-\frac{\phi-\phi_0-2k_0F(t)}{4(\delta_0^4+F(t)^2)/\delta_0^2}} e^{i \left[ k_0(\phi-\phi_0) + \frac{\phi-\phi_0-2k_0F(t)}{4(\delta_0^4+F(t)^2)/F(t)} - k_0^2 F(t) \right]} \quad (39)$$

This is a moving wave packet with the time-dependent standard deviation given by

$$\delta(t) := \left| \frac{\delta_0^2 + 1F(t)}{\delta_0} \right| = \sqrt{\frac{\delta_0^4 + F(t)^2}{\delta_0^2}} \quad (40)$$

Clearly this yields the probabilistic density

$$|\psi(\phi, t)|^2 = \frac{1}{\sqrt{2\pi}\delta(t)} e^{-\frac{(\phi-\phi_0-2k_0F(t))^2}{2\delta(t)^2}}. \quad (41)$$

Furthermore, from (34)

$$P^2 = k_0^2 + \sigma^2 \quad (42)$$

which is time-independent. With suitable choice of the origin of the parameter time  $t$ ,  $R(t)$  can now be determined by solving (33) to yield

$$R(t)^2 = \begin{cases} \frac{P \sin 2t}{2\sqrt{3}} & (K = 1) \\ \frac{Pt}{\sqrt{3}} & (K = 0) \\ \frac{P \sinh 2t}{2\sqrt{3}} & (K = -1) \end{cases}. \quad (43)$$

Substituting (43) into (36) we can finally determine  $F(t)$  to be

$$F(t) = \begin{cases} \frac{\sqrt{3}}{2P} \ln \frac{\tan t}{\tan t_0} & (K = 1) \\ \frac{\sqrt{3}}{2P} \ln \frac{t}{t_0} & (K = 0) \\ \frac{\sqrt{3}}{2P} \ln \frac{\tanh t}{\tanh t_0} & (K = -1) \end{cases}. \quad (44)$$

It's my pleasure to thank Profs R W Tucker, C J Isham and M A H MacCallum for fruitful conversations. Continued support from EPSRC and BAE Systems enabling this work is gratefully acknowledged.

# References

- [1] Wang C 2003 *Class. Quantum Grav.* **20** 3151 (gr-qc/0304101)
- [2] Misner C W 1969 *Phys. Rev. D* **186** 1319
- [3] Kuchař K V 1992 *Proc. 4th Canadian Conf. on General Relativity and Relativistic Astrophysics*  
eds G Kunstatter, D E Vicent and J G Williams (Singapore: World Scirntific)
- [4] Blyth W F and Isham C J 1975 *Phys. Rev. D* **11** 768
- [5] Carlip S 2001 *Rep. Prog. Phys.* **64** 885
- [6] Kibble T W B 1978 *Commun. Math. Phys.* **64** 73
- [7] Weinberg S 1989 *Ann. Phys., NY* **194** 336