

Time in quantum gravity by weakening the Hamiltonian constraint

Hrvoje Nikolić

*Theoretical Physics Division, Rudjer Bošković Institute, P.O.B. 180, HR-10002 Zagreb, Croatia.**
(Dated: November 20, 2018)

We replace the usual Hamiltonian constraint of quantum gravity $H|\psi\rangle = 0$ by a weaker one $\langle\psi|H|\psi\rangle = 0$. This allows $|\psi\rangle$ to satisfy the time-dependent functional Schrödinger equation. In general, only the phase of the wave function appears to be time independent. The resulting quantum theory has the correct classical limit and thus provides a viable theory of quantum gravity that solves the problem of time without introducing additional nongravitational degrees of freedom.

PACS numbers: 04.60.Ds

The classical Hamiltonian of the Einstein theory of gravity vanishes when the metric satisfies the Einstein equations [1, 2, 3, 4]. This is a consequence of the fact that the Einstein theory of gravity is a theory invariant with respect to arbitrary time reparametrizations. It is widely believed that this classical Hamiltonian constraint

$$H = 0 \quad (1)$$

should be quantized such that every physical state ψ satisfies the Wheeler-DeWitt (WDW) [5] equation $\hat{H}\psi = 0$, where \hat{H} is the corresponding quantum mechanical operator. On the other hand, the functional Schrödinger equation takes the form

$$\hat{H}\psi = i\hbar\partial_t\psi. \quad (2)$$

Owing to the WDW equation, the right-hand side of (2) should vanish, which implies that ψ does not depend on time. However, if ψ does not depend on time, then it is not clear how it can be consistent with the fact that classical quantities do depend on time. This is the problem of time in quantum gravity [3, 4].

The problem can be partially resolved when the interaction of gravity with matter is introduced. For example, solving the WDW equation by using the WKB approximation, one finds that the matter wave functional satisfies a “many-fingered time” Tomonaga-Schwinger equation [6, 7]. However, this does not solve the problem of time beyond the WKB approximation. In particular, this does not lead to any time dependence in the case of pure gravity without matter, which is in contradiction with the classical theory of gravity.

In this Letter we propose a simple general solution of the problem of time in quantum gravity. For simplicity, we study the case of pure gravity without matter, but the results are easily generalized to any theory with a vanishing classical Hamiltonian, including the case of gravity interacting with matter. The basic idea is to allow ψ to depend on time by satisfying (2), but, at the same time, to require that this should not contradict the Hamiltonian constraint (1) in the classical limit. As a byproduct, we also find that this approach partially resolves another important problem of quantum gravity – the problem of

operator ordering. It appears that the requirement of unitary time evolution of ψ leads to a restriction on the operator ordering of the Hamiltonian.

Let $\psi(g, t) = \langle g|\psi(t)\rangle$ be a time-dependent wave function, where g represents the canonical degrees of freedom. (In the case of gravity, g is the 3-metric.) Instead of the WDW equation $\hat{H}|\psi\rangle = 0$, we require a weaker form of the quantum Hamiltonian constraint:

$$\langle\psi|\hat{H}|\psi\rangle = 0. \quad (3)$$

This corresponds to the requirement that the average value of \hat{H} vanishes, which is consistent with the classical Hamiltonian constraint (1).

Note that such a weaker form of the constraint is similar to the Gupta-Bleuler quantization of the classical Lorentz-condition constraint $\partial_\mu A^\mu = 0$ of electrodynamics [8, 9]. In quantum electrodynamics, it is not consistent to require $\partial_\mu \hat{A}^\mu|\psi\rangle = 0$. However, one can write $\hat{A}^\mu = \hat{A}^{\mu(+)} + \hat{A}^{\mu(-)}$, where $\hat{A}^{\mu(+)}$ ($\hat{A}^{\mu(-)}$) is the positive (negative) frequency part. The Gupta-Bleuler method consists in requiring $\partial_\mu \hat{A}^{\mu(+)}|\psi\rangle = 0$, which is equivalent to $\langle\psi|\partial_\mu \hat{A}^{\mu(-)} = 0$. This implies $\langle\psi|\partial_\mu \hat{A}^\mu|\psi\rangle = 0$, which is analogous to (3).

The condition (3) can be written as

$$\int \mathcal{D}g \psi^* \hat{H} \psi = 0. \quad (4)$$

By writing $\psi = Re^{iS/\hbar}$ (where R and S are real), using (2), and requiring the conservation of probability

$$\frac{d}{dt} \int \mathcal{D}g \psi^* \psi = 0, \quad (5)$$

one finds that (4) is fulfilled if

$$\partial_t S = 0. \quad (6)$$

In other words, the Hamiltonian constraint implies that the wave function takes the form

$$\psi(g, t) = R(g, t) e^{iS(g)/\hbar}. \quad (7)$$

The phase S is time independent, but the square of the absolute value $|\psi|^2 = R^2$ may depend on time.

Let us investigate the idea above in more detail. To simplify the analysis, we write the Hamiltonian of classical gravity in the condensed notation

$$H = G_{AB}(g)\pi^A\pi^B + V(g), \quad (8)$$

where $g \equiv \{g_A\}$ and $G_{AB} = G_{BA}$. Explicitly,

$$G_{AB}\pi^A\pi^B \equiv \kappa \int d^3x G_{ijkl}\pi^{ij}\pi^{kl}, \quad (9)$$

$$V \equiv -\kappa^{-1} \int d^3x \sqrt{|g^{(3)}|} R^{(3)}. \quad (10)$$

Here π^{ij} ($i, j = 1, 2, 3$) are the canonical momenta corresponding to the canonical coordinates g_{ij} , $g^{(3)}$ is the determinant of g_{ij} , $R^{(3)}$ is the curvature constructed from the metric g_{ij} , $\kappa = 8\pi G_N$, and

$$G_{ijkl} = \frac{\sqrt{|g^{(3)}|}}{2} (g_{ik}g_{jl} + g_{jk}g_{il} - g_{ij}g_{kl}) \quad (11)$$

is the supermetric.

In the quantum case, the momenta π^A become the derivative operators

$$\hat{\pi}^A = -i\hbar \frac{\delta}{\delta g_A} \equiv -i\hbar \partial^A. \quad (12)$$

Since G_{AB} depends on g , different orderings of the operators in the first term of (8) are not equivalent. Therefore, we study a quantum variant of (8) of the form

$$\hat{H} = \frac{1}{a+b} [aG_{AB}\hat{\pi}^A\hat{\pi}^B + b\hat{\pi}^AG_{AB}\hat{\pi}^B] + V, \quad (13)$$

where a and b are arbitrary constants. Eq. (13) certainly does not correspond to the most general ordering of \hat{H} , but represents a large class of possible orderings. This class of orderings will be sufficient to demonstrate the fact that the unitarity condition (5) imposes restrictions on the otherwise arbitrary choice of ordering. Using (12), (13) becomes

$$\hat{H} = -\hbar^2 [G_{AB}\partial^A\partial^B + cG_{AB}{}^{,A}\partial^B] + V, \quad (14)$$

where $G_{AB}{}^{,A} \equiv \partial^AG_{AB}$ and $c = b/(a+b)$. Using (7), one finds that the complex equation (2) is equivalent to a set of two real equations

$$G_{AB}S^{,A}S^{,B} + V + Q = 0, \quad (15)$$

$$\partial_t R^2 + \partial^A(R^2 2G_{AB}S^{,B}) + (c-1)2R^2 G_{AB}{}^{,A}S^{,B} = 0, \quad (16)$$

where

$$Q = -\hbar^2 \left(G_{AB} \frac{R^{,AB}}{R} + cG_{AB}{}^{,B} \frac{R^{,A}}{R} \right). \quad (17)$$

Now we see that (5) is satisfied if the last term in (16) vanishes, because then the left-hand side of (5) reduces to an integral over a total derivative, which vanishes. This implies that we must take $c = 1$ which is equivalent to $a = 0$, so (13) reduces to

$$\hat{H} = \hat{\pi}^A G_{AB} \hat{\pi}^B + V. \quad (18)$$

This fixes a unique ordering among all the orderings that belong to the class of orderings defined by (13). Note that this ordering provides that not only norms $\langle \psi | \psi \rangle$, but also all scalar products $\langle \psi_1 | \psi_2 \rangle$, are time independent. This is because (2) and (18) imply

$$\frac{d}{dt} \int \mathcal{D}g \psi_1^* \psi_2 = \hbar \int \mathcal{D}g \partial^A [G_{AB}(\psi_1^* i \overleftrightarrow{\partial}^B \psi_2)], \quad (19)$$

which vanishes because the integral over a total derivative vanishes.

Let us now consider the classical limit $\hbar \rightarrow 0$. Eqs. (15) and (16) with $c = 1$ become

$$G_{AB}S^{,A}S^{,B} + V = 0, \quad (20)$$

$$\partial_t R^2 + \partial^A(R^2 2G_{AB}S^{,B}) = 0. \quad (21)$$

Eq. (20) is the classical Hamilton-Jacobi equation equivalent to the classical Hamiltonian constraint (1), provided that we make the classical identification $\pi^A = S^{,A}$. The classical time derivative of the metric $\dot{g}_A \equiv \partial_t g_A$ is given by the classical Hamilton equation

$$\dot{g}_A = \frac{\partial H}{\partial \pi^A} = 2G_{AB}\pi^B, \quad (22)$$

so Eq. (21) can be written as

$$\partial_t \rho + \partial^A(\rho \dot{g}_A) = 0, \quad (23)$$

where $\rho = R^2$. Eq. (23) is naturally interpreted as a local (in the configuration space) probability conservation in a classical statistical ensemble of metric configurations described by the probability density $\rho(g, t)$. This conservation can be written in terms of a total derivative as

$$\frac{d\rho}{dt} = 0. \quad (24)$$

Note that the analog of (23) for the WDW equation corresponds to ρ that does not explicitly depend on time, i.e., satisfies $\partial_t \rho = 0$. On the other hand, there is no reason for a classical probability density to satisfy this constraint. The initial classical probability density $\rho(g, t_0)$ is arbitrary, while its further time evolution is determined by classical dynamics, which, in general, leads to an explicitly time-dependent solution ρ of Eq. (23). In this sense, the conventional stronger form of the Hamiltonian

constraint does not have the correct classical limit, while our weaker form of the Hamiltonian constraint does.

Note that the interpretation of ρ as a classical probability density in the limit $\hbar \rightarrow 0$ is particularly natural from the point of view of the Bohmian interpretation of quantum gravity, which is usually formulated within the conventional WDW approach (see, e.g., [10] and references therein). However, in order to interpret ρ as a classical probability density in the limit $\hbar \rightarrow 0$, it is not necessary to adopt the Bohmian interpretation of the full quantum theory.

For those who are skeptical about such a probabilistic interpretation of ρ in the classical limit, there are also other, more conventional, ways of looking at the classical limit. One possibility is to construct a solution $\psi(g, t)$ peaked around a classical solution $g_A^{\text{class}}(t)$, so that

$$R^2(g, t) \approx \delta(g - g^{\text{class}}(t)), \quad (25)$$

where $\delta(g)$ is a short symbol for $\prod_A \delta(g_A)$. Then the average value of g_A is

$$\langle g_A \rangle = \int \mathcal{D}g \psi^*(g, t) g_A \psi(g, t) \approx g_A^{\text{class}}(t). \quad (26)$$

We see that it is necessary that R depends on time in order to obtain the correct classical limit with a time-dependent metric. Another approach to the classical limit is through the Ehrenfest theorem. The Schrödinger equation (2) implies

$$\begin{aligned} \partial_t \langle \psi | g_A | \psi \rangle &= \langle \psi | i\hbar^{-1} [\hat{H}, g_A] | \psi \rangle, \\ \partial_t \langle \psi | \hat{\pi}^A | \psi \rangle &= \langle \psi | i\hbar^{-1} [\hat{H}, \hat{\pi}^A] | \psi \rangle. \end{aligned} \quad (27)$$

Using the fact that

$$\begin{aligned} i\hbar^{-1} [\hat{H}, g_A] &= \{ \hat{H}, g_A \}_{\text{PB}} + \mathcal{O}(\hbar), \\ i\hbar^{-1} [\hat{H}, \hat{\pi}^A] &= \{ \hat{H}, \hat{\pi}^A \}_{\text{PB}} + \mathcal{O}(\hbar), \end{aligned} \quad (28)$$

we see that the classical limit $\hbar \rightarrow 0$ applied to (27) leads to the classical equations of motion for the averaged values, provided that $|\psi\rangle$ satisfies a time-dependent Schrödinger equation (2).

So far, we have not discussed a very important issue – the issue of reparametrization invariance of our quantum theory. For that purpose, note first that the Hamiltonian $H = \int d^3x \mathcal{H}$, known also under the name super-hamiltonian, should be distinguished from the canonical Hamiltonian

$$H^{\text{can}} = \int d^3x [N\mathcal{H} + N_i \mathcal{H}^i], \quad (29)$$

where $N = 1/\sqrt{|g^{00}|}$ is the lapse function, $N_i = g_{0i}$ is the shift function, and $\mathcal{H}^i = -2\pi^{ij}{}_{|j}$ is the supermomentum density. Strictly speaking, one should replace (2) with

$$\hat{H}^{\text{can}} \psi = i\partial_t \psi. \quad (30)$$

In this sense, our previous results correspond to the choice of gauge $N = 1$, $N_i = 0$, because only then $H = H^{\text{can}}$. However, it is easy to modify all the equations such that they are correct for all choices of N and N_i . Integrating by parts, one finds

$$\int d^3x N_i \mathcal{H}^i = \int d^3x 2[N_{i|j} - \Gamma_{kj}^k N_i] \pi^{ij} \equiv F_A \pi^A. \quad (31)$$

Redefining G_{AB} in (9) and V in (10) such that the factor N is included, the generalization of (18) is

$$\hat{H}^{\text{can}} = \hat{\pi}^A G_{AB} \hat{\pi}^B + \frac{1}{2}(F_A \hat{\pi}^A + \hat{\pi}^A F_A) + V. \quad (32)$$

With this ordering, the generalizations of (15) and (16) are

$$G_{AB} S^{,A} S^{,B} + F_A S^{,A} + V + Q = 0, \quad (33)$$

$$\partial_t R^2 + \partial^A [R^2 (2G_{AB} S^{,B} + F_A)] = 0. \quad (34)$$

In the classical limit, (33) becomes the Hamilton-Jacobi equation for the classical constraint $H^{\text{can}} = 0$, while (34), together with $\pi^A = S^{,A}$ and the generalization of (22)

$$\dot{g}_A = \frac{\partial H^{\text{can}}}{\partial \pi^A} = 2G_{AB} \pi^B + F_A, \quad (35)$$

becomes the conservation equation (23) equivalent to (24). In addition, we also quantize the classical super-momentum constraint $\mathcal{H}^i = 0$ in the usual way, by postulating

$$\hat{\mathcal{H}}^i \psi = 0. \quad (36)$$

This constraint provides the reparametrization invariance of the quantum theory with respect to the coordinate transformations of the form $x^i \rightarrow x'^i = f^i(x^1, x^2, x^3)$. The constraint (36) implies that (30) is equivalent to (2) with a redefined Hamiltonian

$$\hat{H} = \int d^3x N \hat{\mathcal{H}}. \quad (37)$$

The resulting Schrödinger equation (2) is covariant with respect to time reparametrizations of the form $t \rightarrow t'(t)$. The variables N and N_i are not quantized because they are not dynamical, i.e., because they do not possess the corresponding canonical momenta.

We still do not have the full reparametrization invariance with respect to general coordinate transformations of the form $x^\mu \rightarrow x'^\mu = f^\mu(x^0, x^1, x^2, x^3)$, $\mu = 0, 1, 2, 3$. We only have the reparametrization invariance with respect to coordinate transformations that do not mix space coordinates with time coordinates, i.e., with respect to transformations that preserve the foliation of spacetime into spacelike hypersurfaces Σ . It is, of course, impossible to obtain a complete symmetry between space

and time coordinates in a canonical approach. Even the classical canonical theory requires the global spacetime topology of the form $\Sigma \times \mathbf{R}$, which introduces asymmetry between space and time. However, the WDW approach treats the space and time constraints in a more symmetrical way than we do, by postulating $\mathcal{H}\psi = 0$, $\mathcal{H}^i\psi = 0$. In this sense, the WDW approach may seem more appealing than our approach. On the other hand, as we have seen, the WDW approach does not have the correct classical limit, while our approach does. If a classical theory is verified experimentally, then it is more important for a quantum theory to have the correct classical limit than to possess all the symmetries that the classical theory possesses. Indeed, the existence of various quantum anomalies teaches us that the quantum theory does *not* need to possess all the symmetries that the classical theory does.

Note also that the existence of a preferred foliation of spacetime is not necessarily a problem if one takes the viewpoint according to which the Einstein theory of gravity may correspond to an effective theory for an underlying more fundamental theory that possesses a preferred time even at the classical level. This speculation is supported by the fact that the low-energy phenomena of various nonrelativistic systems, such as certain kinds of fluids [11, 12], may be described by generally covariant equations.

Finally, let us note one new way of looking at the problem of general covariance in quantum gravity. The WDW equations for ψ contain neither space nor time derivatives and therefore they are covariant. Our Schrödinger equation contains a time derivative but does not contain space derivatives, so it is not covariant. However, in our approach, S does not depend on time, so (15) and (33) take the same covariant form as the analogous equations in the WDW approach. (If S were time dependent, then the right-hand sides of (15) and (33) would be equal to $-\partial_t S$.) The nonvanishing partial time derivatives appear only in (16) and (34), which makes these equations noncovariant. However, there is a way of writing the noncovariant equation (34) in a covariant form. All we have to do is to assume that the classical equations (35) and $\pi^A = S^{,A}$ are, in some sense, also valid in the quantum case. With this assumption, (34) can be written as (23) which is equivalent to (24). Since ρ does not depend on x^i (recall that ψ is a functional, not a function, of $g_{ij}(x^1, x^2, x^3)$), (24) can be written in the covariant form as

$$\frac{d\rho}{dx^\mu} = 0. \quad (38)$$

Fortunately, there is a way of making sense of equations (35) and $\pi^A = S^{,A}$ in the quantum domain; they correspond to the already mentioned Bohmian interpretation of quantum gravity. In this interpretation, $g_A(t)$ is a deterministic hidden variable. Note that, owing to the

nonclassical term Q in (33), the equation of motion (35) is not covariant; the dynamics of $g_{\mu\nu}(x^0, x^1, x^2, x^3)$ is described by a modification of the classical Einstein equation covariant only with respect to transformations that preserve the foliation of spacetime [10]. In a way, the non-covariance is translated from the equations that determine the wave function ψ to the equations that determine unobservable hidden variables. In other words, we have a remarkable and surprising result that the Bohmian interpretation should be invoked in order to write the equations that determine ψ in a covariant form. It remains to be seen in future research if this result has deeper consequences on the role of the Bohmian interpretation in quantum gravity.

To summarize, in this Letter we have shown that time in pure quantum gravity without additional nongravitational degrees of freedom can be introduced by replacing the usual Hamiltonian constraint $\hat{H}|\psi\rangle = 0$ by a weaker one $\langle\psi|\hat{H}|\psi\rangle = 0$. The results are easily generalized to any theory with a vanishing classical Hamiltonian, including the case of gravity interacting with matter. Contrary to the conventional approach, based on the WDW equation, which cannot explain why metric can depend on time in the classical limit, our approach does explain that. Unfortunately, with our approach, we do not completely solve the problem of the full reparametrization invariance in quantum gravity, but at least we provide some hints for a solution of this problem. In particular, it appears that the Bohmian interpretation might have a previously unnoted role in establishing the general covariance of quantum gravity.

This work was supported by the Ministry of Science and Technology of the Republic of Croatia under Contract No. 0098002.

* Electronic address: hrvoje@thphys.irb.hr

- [1] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman and Company, New York, 1973).
- [2] K. Kuchar, in *Relativity, Astrophysics, and Cosmology*, edited by W. Israel (D. Reidel Publishing Company, Dordrecht/Boston, 1973).
- [3] T. Padmanabhan, *Int. J. Mod. Phys. A* **4**, 4735 (1989).
- [4] C. J. Isham, gr-qc/9210011.
- [5] B. DeWitt, *Phys. Rev.* **160**, 1195 (1967).
- [6] T. Banks, *Nucl. Phys. B* **249**, 332 (1985).
- [7] C. Kiefer and T. P. Singh, *Phys. Rev. D* **44**, 1067 (1991).
- [8] S. S. Schweber, *An Introduction to Relativistic Quantum Field Theory* (Harper & Row, New York, 1961).
- [9] L. H. Ryder, *Quantum Field Theory* (Cambridge University Press, Cambridge, 1984).
- [10] A. Shojai and F. Shojai, *Class. Quant. Grav.* **21**, 1 (2004); gr-qc/0311076.
- [11] M. Visser, *Class. Quant. Grav.* **15**, 1767 (1998).
- [12] G. E. Volovik, *Phys. Rep.* **351**, 195 (2001).