

General relativity and quintessence explain the Pioneer anomaly

J.P. Mbelek

Service d'Astrophysique, C.E. Saclay
F-91191 Gif-sur-Yvette Cedex, France

July 4, 2018

Abstract

The anomalous time depending blueshift, the so-called "Pioneer anomaly", that was detected in the radio-metric data from Pioneer 10/11, Ulysses and Galileo spacecraft may not result from a real change of velocity. Rather, the Pioneer anomaly may be understood within the framework of general relativity as a time depending gravitational frequency shift accounting for the time dependence of the density of the dark energy when the latter is identified with quintessence. Thus, instead of being in conflict with Einstein equivalence principle, the main Pioneer anomaly appears merely as a new validation of general relativity in the weak field and low velocity limit.

1 Introduction

Since 1998, Anderson *et al.* have continuously reported an anomalous Doppler shift derived from a twenty years study of radio-metric data from Pioneer 10/11, Ulysses and Galileo spacecraft [1]. The observed effect mimics a constant acceleration acting on the spacecraft with magnitude $a_P = (8.74 \pm 1.33) \times 10^{-8} \text{ cm s}^{-2}$, directed toward the sun to within the accuracy of the Pioneers' antennas and a steady frequency drift $\frac{d\Delta\nu}{dt} \simeq 6 \times 10^{-9} \text{ Hz/s}$ which equates to a "clock acceleration" $\frac{d(\Delta\nu/\nu)}{dt} = \frac{a_P}{c}$. An independant analysis of radio Doppler tracking data from the Pioneer 10 spacecraft for the time period 1987 - 1994 confirms the previous observations [2]. Besides, it

has been noted that the magnitude of a_P compares nicely to the Modified Newtonian Dynamics (MOND) acceleration constant $a_0 \sim cH_0$ [3], where H_0 is the Hubble parameter at present cosmological epoch. Actually, the reported anomaly cannot really be related to MOND. Indeed, the gravitational pulling of the sun up to 100 AU still yields an acceleration greater than a_0 by at least three orders of magnitude, equating a_0 only at about 3000 AU. Hence, Newtonian dynamics up to GR corrections should apply to spacecraft. The same argument was put forward by M. Milgrom [4] to reject the claim that MOND fails in the laboratory [5]. In a further study, Anderson *et al.* found that the small difference of magnitude of a_P for Pioneer 10 and Pioneer 11 is related to their difference of spin-rate history [6]. Removing the spin-rate change contribution, yields an apparent anomalous acceleration $a_P = (7.84 \pm 0.01) \times 10^{-8} \text{ cm s}^{-2}$ of the same amount with a great accuracy during a long time interval for both Pioneer 10/11. Since this has found a conventional explanation [7], it points out in favor of an external origin for the main Pioneer anomaly at the expense of any possible internal cause (*e. g.*, heat systematics). As yet, this also excludes any significant drag force acting on the spacecraft (*e. g.*, the mirror world explanation [8]) as an explanation of the anomaly. Indeed, the velocities of the Pioneer 10 and 11 relative to the sun differ by about 5% in the time interval of interest. Moreover, Whitmire and Matese [9] have carried out a statistical analysis by comparing the mean original energy obtained from the Comet Catalogue of Marsden and Williams [10], with and without corrections due to an hypothetical Pioneer anomalous acceleration, to that predicted by galactic tidal theory. They conclude that the implied higher binding energy is incompatible with the established evidence that the galactic tide is dominant in making Oort cloud comets observable. The whole work of these authors is based on the major assumption that the Pioneer anomaly reveals a real acceleration. As they emphasized, systematic outgassing would not hide the effects of such anomalous acceleration if real. There-

fore, the alternative that the Pioneer anomaly does not result from a real change in velocity deserves to be investigated. Indeed, a direct interpretation of the observational data from the spacecraft implies merely a time dependent frequency blueshift of the photons. However, any true Doppler shift would involve an accompanying acceleration. Hence, given our knowledge of photons frequency shift, the only other relativistic effect that can be confused, at the solar system scale, with a real Doppler shift is the gravitational frequency shift. In the weak field and low velocity limit, this should involve a time dependent gravitational potential instead of a spatial dependent one, in order to avoid an induced anomalous acceleration for test bodies. Of course, the origin of a time dependent gravitational potential would not be relevant if not justified on the physical ground. Actually, although unfamiliar, we show that a time dependent gravitational potential follows from our present knowledge of the matter-energy content of the universe. Indeed, quintessence is one possible form of dark energy that explains the recent discovery of the accelerating universe in term of a scalar field whose energy density is time dependent and dominant over that of the ordinary matter at present cosmological epoch. Now, a time dependent matter-energy density will naturally generate a time dependent potential. Given that the energy density of the quintessence is on average the same everywhere depending only on time, this should holds even at small scale, not only at the cosmological level. The idea of a time dependent gravitational potential have been revived recently by A. F. Rañada [11] and K. Trencovski [12]. However, both attempts introduce only phenomenologically the time dependent potential (*e. g.* by choosing its sign by convenience and trying to match exactly a_P with cH_0). As a common feature, none of both approaches refer to a field equation to derive the time dependent potential. Above all, none rely intirely on GR. Besides, we note that the link between the time dependent potential and dark energy (whether this pertains to a true cosmological constant or quintessence is not expressed at all) is either put by hand [11] or just

invoked by words [12] by the authors. The aim of this paper is to demonstrate that the Pioneer anomaly can find a natural explanation within the framework of general relativity (GR) when the gravitational contribution of quintessence is properly accounted for. Then, the Pioneer anomaly interprets as a gravitational blueshift in excess induced locally by the energy density of quintessence which provides a time dependent contribution to the gravitational potential. Furthermore, because of its weakness, the Pioneer anomaly seen as a time dependent gravitational potential actually entails no contradiction with respect to the four classical tests of GR. Also, it is worth noticing that the analysis that is carried out in the following involves the Hubble constant only through the time dependence and non locality of the energy density of the quintessence. In any case, the matter content of gravitational bound systems like galaxies or the solar system are almost decoupled from the general expansion of the universe whose effect is quite negligible on the stars or planetary orbits (see [13] and [14], section 16.1 H). Indeed, the exact mathematical treatments of gravitational bound systems in an expanding universe that are based on GR solely (with or without a true cosmological constant) always lead to conclude to a negligible effect out of an ensemble of dust particles [15, 16, 17] or a star [18]. Thus, although the net effect is not exactly zero, it cannot account for the Pioneer anomaly in the framework of GR (see *e. g.*, [19]) without reconsidering the matter-energy content of the universe.

2 Time varying potential from quintessence

Let us consider the dark energy as quintessence, that is a neutral scalar field coupled to ordinary matter only through its gravitational influence. We show that this involves a time varying contribution to the gravitational potential even for gravitationally bound systems. As one knows, unlike the ordinary matter (luminous or dark matter) whose mass density tends to decrease continuously driven by the expansion

of the universe or clumps locally under the gravitational pulling, the energy density of the dark energy remains constant on average everywhere at any given cosmic time (quintessence models) if not a universal constant (a property of spacetime itself) in the case of a true cosmological constant. For that reason, we will not assume a static metric for the solar system hereafter. Instead, in addition to the point like mass of the sun and the other heavenly bodies (whose effects as usual are seen as perturbations), we will also deal with the gravitational effect of the uniform mass density, ρ_Q , of the dark energy which behaves like a cosmological "constant" $\Lambda = 8\pi G\rho_Q/c^2$. By choosing the coordinates such that the metric tensor be diagonal, the metric still expresses in the canonical form

$$ds^2 = \left(1 + 2\frac{V}{c^2}\right)c^2 dt^2 - \left(1 + 2\frac{V}{c^2}\right)^{-1} dr^2 - r^2 (d\theta^2 + \sin^2\theta d\varphi^2), \quad (1)$$

but with the potential V now depending on both spatial and time coordinates, where $x^0 = ct$, $x^1 = r$ is the radius from the sun, $x^2 = \theta$ is the polar angle and $x^3 = \varphi$ is the azimuthal angle. The function V is derived from Einstein equations $R_{\mu\nu} - \frac{1}{2}(R + 2\Lambda)g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}$, in the limit of the weak and slowly varying gravitational field. We proceed as usual by setting $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, where all the $h_{\mu\nu}$'s are much less than unity. Thus, the Einstein equations reduce to $R_{00} = \frac{4\pi G}{c^2}(\rho_{matter} + \rho_Q + 3\frac{P_Q}{c^2})$ in the weak field and low velocity limit. Expanding g_{00} and g_{11} in the first order in V/c^2 and R_{00} in the first order of the derivatives of the $h_{\mu\nu}$'s (see e. g., [14], section 15.1 and [20]) but without discarding as usual the time derivatives, one gets $R_{00} = \frac{1}{2}\eta^{\alpha\beta}(h_{\beta 0,0\alpha} - h_{00,\beta\alpha} + h_{0\alpha,\beta 0} - h_{\beta\alpha,00}) = \frac{1}{c^2}\left(\frac{1}{c^2}\frac{\partial^2 V}{\partial t^2} + \Delta V\right)$. Hence, the equation to solve reads

$$\frac{1}{c^2}\frac{\partial^2 V}{\partial t^2} + \Delta V = 4\pi G\left(\rho_{matter}(\vec{r}) + \rho_Q + 3\frac{P_Q}{c^2}\right), \quad (2)$$

where the equation of state of the quintessence is given by $P_Q = w_Q \rho_Q c^2$ with equation of state parameter $w_Q \simeq -1$. Usually, one solves (2) by assuming a static

potential. This yields (up to an integration constant)

$$V = V_N - \frac{1}{6} \Lambda c^2 r^2, \quad (3)$$

where the Newtonian potential $V_N = V_N(\vec{r})$ is a solution of Gauss equation

$$\Delta V_N = 4\pi G \rho_{matter} \quad (4)$$

and the quadratic extrapotential term is related to the dark energy density. However, because of the fine-tuning that would be required both for the cancellation of the vacuum energy (quantum field theory problem) and the coincidence problem (why the density of the dark energy and that of the ordinary matter are of the same order just today, whereas both rates of evolution are quite different), quintessence models are prevailing in the literature at the expense of a true cosmological constant [21]. A quintessence model involves a dynamical cosmological constant driven by an almost homogeneous scalar field (spatial fluctuations cancel out on average) [22], Q , with a suitable self-interaction potential (most often chosen as an inverse power law of Q). This means that in some sens, though not exactly, Λ is defined as a function of time in such models. So, the previous solution (3) which is purely static does not hold any more. Thus, while in the case of a true positive cosmological constant the vacuum solution yields a de Sitter spacetime, spacetime is rather asymptotically Minkowskian in the case of quintessence whose energy density decreases monotonously toward zero with respect to time. Since $\rho_Q = \rho_Q(t)$ is non zero everywhere, Birkhoff's theorem cannot rigorously be invoked to infer a static spacetime. Let us emphasize that the generalization of Birkhoff's theorem to the case with a Λ -term (cosmological constant) is restricted to the case when the latter is indeed a true universal constant [23]. Besides, it is clear that making a gauge transformation in order to cancel out any time dependent potential term will not be helpful, since this is associated with a change of reference frame. Now, we need to

study the motion of spacecraft in the same reference frame as for the planets. So, let us look for a solution of the form $V = V_N(\vec{r}) + V_Q(t)$. Equation (2) then splits into equation (4) and the following

$$\frac{d^2 V_Q(t)}{dt^2} = -8\pi G \rho_Q(t) c^2. \quad (5)$$

At present cosmological epoch, $\rho_Q(t)$ is a slowly varying function as compared to the time scale that concerns the Pioneer 10/11 spacecraft. Therefore, the solution of equation (5) reads in the first approximation

$$V_Q(t) = -\frac{3}{2} \Omega_Q H_0^2 c^2 (t - t_0)^2 - 3\Omega_Q H_0^2 c^2 (t_0 - t_i)(t - t_0) + V_Q(t_0), \quad (6)$$

where t_i is the cosmic time at which $V_Q(t)$ passes by its maximum value and $\Omega_Q = \Lambda(t_0)c^2/3H_0^2$ denotes the density parameter associated with the quintessence at present cosmological epoch. Let us point out that one passes from the non-Newtonian extra potential of relation (3) to relation (6) by making the substitution $r \rightarrow c(t - t_i)$ in (3) then multiplying the whole result by 3 (because of the isotropy of the 3-space). As regards the phenomenology, it seems reasonable to assume that the linear term of eq(6) is of the order of the quadratic one early in the past but finally dominates over it at present epoch. These conditions are satisfied for $t_i \simeq t_0/2$ and imply $V_Q(t_0) \simeq V_Q(0)$. Present cosmological observations yield $\Omega_Q \simeq 0.7$ and $H_0 t_0 \simeq 0.96_{-0.06}^{+0.09}$ for a flat universe [24]. Hence, relation (6) yields

$$\frac{V_Q(t)}{c^2} \simeq -H_0^2 (t - t_0)^2 - H_0 (t - t_0) + \frac{V_Q(t_0)}{c^2}. \quad (7)$$

According to recent observations [25], the cosmic jerk when the expanding universe made the transition from deceleration to acceleration corresponds to a redshift $z_j = 0.46 \pm 0.13$ or equivalently a fractional look-back time $(t_0 - t_j)/t_0 \simeq 0.4$. So, the choice of parameter made above is consistent with the natural expectation that the potential energy $U_Q = m V_Q$ of a test particle of mass m decreases monotonously past t_j .

3 Interpretation of the observational data

3.1 The main Pioneer anomaly

Since $g_{00} = 1 + 2\left(\frac{V_N + V_Q}{c^2}\right)$, the resulting gravitational frequency shift of photons is derived from the following relation

$$\frac{\nu_B}{\nu_A} = \sqrt{\frac{g_{00}(A)}{g_{00}(B)}} \simeq 1 + Z_{ordinary} + Z_{anomalous}, \quad (8)$$

where $Z_{ordinary} = \frac{V_N(A) - V_N(B)}{c^2}$ represents the familiar gravitational frequency shift whereas $Z_{anomalous} = \frac{V_Q(t_A) - V_Q(t_B)}{c^2}$ is the time varying gravitational frequency shift related to the quintessential dark energy. For a small time of flight, $\Delta t = t_B - t_A$, of the photons of the communication signals between the spacecraft and the earth (Δt negligible as compared to the age of the universe), one gets in the first order approximation on account of relation (7)

$$Z_{anomalous} \simeq -\frac{1}{c^2} \frac{dV_Q}{dt} \left(\frac{t_A + t_B}{2} \right) \Delta t. \quad (9)$$

Since $\frac{dV_Q}{dt} < 0$, relation (9) implies a systematic blueshift in excess. For the time interval of interest for the Pioneer 10/11 spacecraft, the quadratic term of relation (7) is negligibly small with respect to the linear one. On account of relation (9) above, this involves an anomalous blueshift in excess

$$Z_{anomalous} = \frac{a_P \Delta t}{c}, \quad (10)$$

where $a_P \simeq H_0 c$ in accordance with observations.

The above analysis can be pushed further by taking into account the revolution and rotation of the earth. Then, one derives the annual and daily modulations following the same reasoning as before. We argue that these modulations are further evidences for the dependence of the spacecraft anomalous frequency shift with respect to time, in particular the time of flight of the photons of the communication signals.

3.2 The absence of a true anomalous acceleration

Moreover, the extrapotential V_Q will not effect the motion of the spacecraft or heavenly bodies being time dependent only. Indeed, in the weak fields and low velocity limit, the geodesic equation simplifies to

$$\begin{aligned}\frac{d^2\vec{r}}{dt^2} &= -\frac{1}{2}\vec{\nabla}g_{00} \\ &= -\vec{\nabla}V_N,\end{aligned}\tag{11}$$

where \vec{r} denotes the position vector of the test body.

3.3 The steady frequency drift

According to Anderson *et al.* (see [6], section V - B and C, p. 19), the "clock acceleration" $a_t = d(\Delta\nu/\nu)/dt$ cannot be attributed to a systematic drift in the atomic clocks of the DSN (Deep Space Network, the network of ground stations that are employed to track interplanetary spacecraft). We show that it follows from the path equation of photons $dP^\mu = -\Gamma_{\alpha\beta}^\mu P^\alpha dx^\beta$ for the time component $\mu = 0$, where $P^0 = \frac{h\nu}{c}$ and $\vec{P} = \frac{h\nu}{c}\vec{u}$ define the 4-momentum P^μ of a photon of frequency ν propagating in the direction of the unit vector \vec{u} . Since $\Gamma_{00}^0 \simeq \frac{1}{c^3}\frac{dV_Q}{dt} \simeq -\frac{a_P}{c^2}$ in weak field and low velocity limit, one finds after removing the gravitational frequency shift due to the Newtonian potential solely

$$a_t \simeq \frac{1}{\nu} \left[\frac{d\nu}{dt} - \left(\frac{d\nu}{dt} \right)_{V_Q=0} \right] \simeq \frac{a_P}{c},\tag{12}$$

as observed. Let us notice that relation (12) can be derived in the usual simple manner from the conservation of energy for the photon to which one assigns a gravitational mass $m = \frac{h\nu}{c^2}$ in accordance with the EP and a kinetic energy $E_c = h\nu$. Further, the other first order correction implied by the potential V_Q comes from the radial component $\mu = 1$. This brings an additional contribution to the deflection angle of the photons near the sun of the order $\Delta\delta \sim a_t \Delta t$ for a time of flight Δt .

Clearly, $\Delta\delta$ is by far less than the classical result $\delta = 1.75''$ by more than 7 orders of magnitude.

4 Discussion

The study that we have carried out in this paper does not pretend to bring the last word on the so-called Pioneer anomaly. We have just tried to clarify the subject as to the consistency of such an effect with general relativity. Although, the results seem encouraging in this respect, we still have at least two problems to face. Namely, whereas the slope of the observed anomalous frequency shift versus time appears almost constant between 20 AU and 80 AU (see *e. g.*, FIG. 1 of reference [1] for Pioneer 10 in the time period 1987-1995), it drops down to zero below 10 AU (see FIG. 6 or FIG. 7 of reference [6]). Nevertheless, Ulysses and Galileo which were moving below or about 5.4 AU (orbital radius of Jupiter) also have shown an almost apparent constant acceleration in their radio Doppler and ranging data but with a high correlation with solar pressure (respectively 0.888 and 0.99). These issues have never been addressed up to now by any theoretical study. Of course, the latter observations cannot fit within the framework of GR as a result of quintessence since the EP would be strongly violated. As a way out, we conjecture that mass loss toward the sun may potentially explain, at least in part if not all, the vanishing of a_P observed on both Pioneer 10/11 spacecraft below 10 AU. Indeed, since the temperature within the Pioneer 10/11 is almost stabilized to about 300 K while decreasing with respect to the heliocentric radius, outgassing out of the spacecraft may occur (note the large error bars on a_P below 10 AU for all four spacecraft). Also given that the mass of the Pioneer has decreased by almost 7% since the date of launch, a mass loss of about 2 ppm is still possible during the following ten years after launch. Because of the gravitational pulling of the sun, we expect that the ejected mass will be directed toward the sun. For both Pioneer 10/11, one just needs an

almost steady rate of mass loss directed toward the sun $\dot{m} \simeq 0.5$ gm per year during the ten years following the launch of the spacecraft but quickly decreasing past this time interval. As for the issue of the high correlation with the solar pressure in the case of Galileo and Ulysses, given the proximity of the sun, it is likely that the capture of dust particles of the solar wind may provide an answer. Let us recall that the Pioneer 10/11, Galileo and Ulysses missions were also dedicated to collect dust particles from the solar wind, all the four spacecraft being endowed with dust detectors on-board [27]. A more detailed study including precise estimates of the aforementioned effects will be presented elsewhere.

5 Conclusion

The study of the Pioneer 10/11 spin-rate change histories points out in favor of the possibility that the cause of the main Pioneer anomaly is external to the spacecraft. However, interpreting the main Pioneer anomaly as a true acceleration leads to some difficulties as regards its effect on Long-Period Comet Orbits and the Oort cloud. Especially, assuming a resisting medium as cause of the anomaly (with the drag acceleration proportional to the orbital speed or its square) not only necessitates a hole inside 10 or 20 AU to avoid large effects on short period comets such as Halley and Encke [26] (note also how huge such an anomalous acceleration would be relative to the earth-moon system, if real about the orbital radius of the earth) but above all also makes difficult to understand the analogous anomalous accelerations observed on Ulysses and Galileo data (orbital radius of both spacecraft less than 5.4 AU). The alternative view that we have adopted then consists to consider that the Pioneer anomaly may not be the result of a new force term. Indeed, the main Pioneer anomaly may be well understood within the framework of GR as a time depending gravitational frequency shift (with respect to the reference frame used by JPL for Doppler tracking of spacecraft). The latter is derived from an extra potential whose

source is the time depending energy density of quintessence like dark energy. Thus, the Pioneer anomaly may accommodate the EP as expressed by the relation $a_t = a_P/c$ between the clock "acceleration", a_t , and the anomalous "acceleration", a_P . Finally, the Pioneer anomaly seems to be rather a new validation of GR in the weak field and low velocity limit. Moreover, the interpretation of the Pioneer anomaly in the framework of GR favors clearly the case of quintessential dark energy at the expense of a true cosmological constant. This exhibits the potentiality of quintessence to help GR gives rise to some Machian behavior. The features emphasized above should make clear the very difference between our new approach and any other one published as yet on the subject.

References

- [1] J. D. Anderson *et al.*, 1998, *phys. Rev. Lett.* **81**, 2858.
- [2] C. B. Markwardt, gr-qc/0208046.
- [3] M. Milgrom, 1983, *ApJ* **270**, 365.
- [4] Private communication.
- [5] A. Abramovici and Z. Vager, 1986, *Phys. Rev.* **D 34**, 3240.
- [6] J. D. Anderson *et al.*, *Phys. Rev.* **D 65**, 082004.
- [7] J. P. Mbelek and M. Michalski, 2004, *Int. J. Mod. Phys.* **D 13**, 865.
- [8] R. Foot and R. R. Volkas, 2001, *Phys. Lett.* **B 517**, 13.
- [9] D. P. Whitmire and J. J. Matese, 2003, *Icarus* **165**, 219.
- [10] B. G. Marsden and G. V. Williams, 2001, *Catalog of Cometary Orbits*, 14th edition, Smithsonian Astrophysical Observatory, Cambridge, MA.

- [11] A. F. Rañada, 2003, Int. J. Mod. Phys. **D 12**, 1755 and gr-qc/0403013.
- [12] K. Trencovski, gr-qc/0402024, gr-qc/0403067 and gr-qc/0404060.
- [13] C. W. Misner, K. S. Thorne, J. A. Wheeler, 1973, Gravitation, Freeman and Company, pp. 719, 739 and 740.
- [14] W. Rindler, 2001, General relativity, Oxford university Press, Oxford.
- [15] P. D. Noerdlinger and V. Petrosian, 1971, ApJ **168**, 1.
- [16] W. B. Bonnor, 1996, MNRAS **282**, 1467.
- [17] J. L. Anderson, 1995, PRL **75**, 3602.
- [18] C. Gilbert, 1956, MNRAS **116**, 678.
- [19] A. L. Licht, gr-qc/0102103.
- [20] I. R. Kenyon, 1990, Relativity Special, General and Cosmological, Oxford university Press, Oxford, section 7.4 and appendix B.
- [21] S. Perlmutter, Physics Today April 2003, 53.
- [22] P. J. Steinhardt, Royal Society of London Transactions Series A **361**, 2497 and <http://feynman.princeton.edu/~steinh/>
- [23] W. Rindler, 1998, phys. Lett. **A 245**, 363, and references therein.
- [24] S. Perlmutter *et al.*, 1999, ApJ **517**, 565.
- [25] A. G. Riess *et al.*, 2004, ApJ **607**, 665.
- [26] J. D. Anderson, S. Turyshev and M. M. Nieto, AAS 201st Meeting, Session 45. The Solar System, Poster : Effect of the Pioneer Anomaly on Long-Period Comet Orbits, <http://www.aas.org/publications/baas/v34n4/aas201/516.htm>.

[27] M. Landgraf, astro-ph/9906300 ; H. Krüger *et al.*, astro-ph/0106322 ;
http://spaceprojects.arc.nasa.gov/Space_Projects/pioneer/PN10&11.html#instruments.