

The Letelier-Gal'tsov spacetime revisited

G. Clément^{a*}, D.V. Gal'tsov^{b†} and P.S. Letelier^{c‡}

^aLaboratoire de Physique Théorique LAPTH (CNRS), B.P.110, F-74941 Annecy-le-Vieux cedex, France

^bDepartment of Theoretical Physics, Moscow State University, 119899, Moscow, Russia

^cDepartamento de Matemática Aplicada-IMECC, Universidade Estadual de Campinas, 13083-970 Campinas, S.P., Brazil

November 27, 2018

Abstract

Contrary to a recent claim by Anderson [*The Mathematical Theory of Cosmic Strings*, I.O.P.Publishing, Bristol 2003], we show that the Letelier-Gal'tsov metric does represent a system of crossed straight infinite cosmic strings moving with arbitrary constant relative velocities.

In a recent book on “the mathematical theory of cosmic strings” [1], Anderson devotes a section to “the proper status of the Letelier-Gal'tsov ‘crossed-string’ metric”. This metric describes in a single coordinate patch the Gott [2] geometry that consists in the gluing of two separate spacetimes, each one representing a tilted moving string. After describing the construction of [3], Anderson argues that “the Letelier-Gal'tsov metric seems capable of describing an arbitrary number of open strings with essentially arbitrary shapes and in arbitrary states of relative motion”, which according to him “suggests something that was perhaps evident from the start, namely that [this metric] is just the static parallel string metric written in an obscure coordinate system”. He then proceeds to substantiate this claim by showing that the geodesic distance between two strings is actually constant. However, as we shall show here, his proof fails due to the omission of an integration constant in the approximate evaluation of a certain integral near a given string. Correcting some misprints contained in the original paper [3], we shall also show that the Letelier-Gal'tsov metric indeed represents a system of crossed straight cosmic strings moving with arbitrary but constant relative velocities.

The Letelier-Gal'tsov “multiple moving crossed cosmic strings spacetime” is obtained from the Minkowski spacetime

$$ds^2 = dt^2 - dz^2 - dZd\bar{Z} \quad (1)$$

by the singular coordinate transformation defined by the line integral

$$Z(\zeta) = X + iY = \int_{\zeta_0}^{\zeta} \prod_{i=1}^N (\xi - \alpha_i(t, z))^{-4\lambda_i} d\xi \quad (2)$$

($\zeta = x + iy$) with ζ_0 fixed. This line element can be rewritten as

$$ds^2 = dt^2 - dz^2 - e^{-4V} (d\zeta + Fdt + Gdz)(d\bar{\zeta} + \bar{F}dt + \bar{G}dz) \quad (3)$$

with

$$V = \sum_{i=1}^N \lambda_i \ln |\zeta - \alpha_i|^2, \quad (4)$$

$$F = \prod_{i=1}^N (\zeta - \alpha_i)^{4\lambda_i} \int_{\zeta_0}^{\zeta} \frac{d\xi}{\prod_{k=1}^N (\xi - \alpha_k)^{4\lambda_k}} \sum_{j=1}^N \frac{4\lambda_j \dot{\alpha}_j}{\xi - \alpha_j}, \quad (5)$$

$$G = \prod_{i=1}^N (\zeta - \alpha_i)^{4\lambda_i} \int_{\zeta_0}^{\zeta} \frac{d\xi}{\prod_{k=1}^N (\xi - \alpha_k)^{4\lambda_k}} \sum_{j=1}^N \frac{4\lambda_j \alpha'_j}{\xi - \alpha_j}. \quad (6)$$

*email: gclément@lapp.in2p3.fr

†email: galtsov@mail.phys.msu.ru

‡email: letelier@ime.unicamp.br

Consider for definiteness two moving strings of equal tension located at $\alpha_{\pm} = \pm\alpha(t, z)$. Then,

$$Z(\zeta) = \int_{\zeta_0}^{\zeta} (\xi^2 - \alpha^2)^{-4\lambda} d\xi, \quad (7)$$

and

$$F(\zeta) = 8\lambda\alpha\dot{\alpha}(\zeta^2 - \alpha^2)^{4\lambda}I(\zeta), \quad G(\zeta) = 8\lambda\alpha\alpha'(\zeta^2 - \alpha^2)^{4\lambda}I(\zeta), \quad (8)$$

with

$$I(\zeta) = \int_{\zeta_0}^{\zeta} (\xi^2 - \alpha^2)^{-4\lambda-1} d\xi. \quad (9)$$

Near $\zeta = +\alpha$, the derivative of I behaves as

$$\frac{\partial I}{\partial \zeta} = (2\alpha)^{-4\lambda-1}(\zeta - \alpha)^{-4\lambda-1} - (1 + 4\lambda)(2\alpha)^{-4\lambda-2}(\zeta - \alpha)^{-4\lambda} + \dots, \quad (10)$$

which integrates to

$$I = C_{\lambda}(\alpha, \zeta_0) - \frac{1}{4\lambda}(2\alpha)^{-4\lambda-1}(\zeta - \alpha)^{-4\lambda} - \frac{1 + 4\lambda}{1 - 4\lambda}(2\alpha)^{-4\lambda-2}(\zeta - \alpha)^{1-4\lambda} + \dots \quad (11)$$

There is no reason for the integration constant $C_{\lambda}(\alpha, \zeta_0)$, which measures the “finite part” of $I(\zeta)$, to vanish. It follows that

$$F(\zeta) = -\dot{\alpha} + 4\lambda C_{\lambda}(\alpha, \zeta_0)(2\alpha)^{1+4\lambda}\dot{\alpha}(\zeta - \alpha)^{4\lambda} - \frac{4\lambda}{1 - 4\lambda}\frac{\dot{\alpha}}{\alpha}(\zeta - \alpha) + \dots, \quad (12)$$

and a similar expansion for $G(\zeta)$. For $\lambda < 1/4$, the second term of (12) dominates the third (the corresponding term is absent from the first Eq. (14) of [3]).

At fixed z and t , the geodesic distance between the two strings is $|Z(\alpha) - Z(-\alpha)|$. From the above, $Z(\alpha)$ varies according to

$$dZ = (\zeta^2 - \alpha^2)^{-4\lambda}(d\zeta + F dt + G dz)|_{\zeta=\alpha} \simeq 8\lambda C_{\lambda}(\alpha, \zeta_0)\alpha(\dot{\alpha} dt + \alpha' dz), \quad (13)$$

showing that the strings are indeed tilted and moving. Note that Anderson, following exactly the same line of reasoning, obtained $dZ \simeq O(\zeta - \alpha)^{1-4\lambda}(\dot{\alpha} dt + \alpha' dz) \rightarrow 0$, because in his evaluation of $F(\zeta)$ near the string he omitted the integration constant C_{λ} and thus included only the first and third terms of (12).

One can directly evaluate $Z(\alpha)$ by computing $dZ(q\alpha)/d\alpha$ (q constant, $0 < q < 1$) and taking the limit $q \rightarrow 1$. Choosing for convenience $\zeta_0 = 0$ (“center-of-mass”),

$$Z_q(\alpha) \equiv Z(q\alpha) = \int_0^{q\alpha} (\xi^2 - \alpha^2)^{-4\lambda} d\xi. \quad (14)$$

The derivative is

$$\frac{dZ_q}{d\alpha} = q(q^2\alpha^2 - \alpha^2)^{-4\lambda} + 8\lambda\alpha \int_0^{q\alpha} (\xi^2 - \alpha^2)^{-4\lambda-1} d\xi. \quad (15)$$

Also, computing (14) by integrating by parts, one obtains

$$(1 - 8\lambda)Z_q = q\alpha(q^2\alpha^2 - \alpha^2)^{-4\lambda} + 8\lambda\alpha^2 \int_0^{q\alpha} (\xi^2 - \alpha^2)^{-4\lambda-1} d\xi = \alpha \frac{dZ_q}{d\alpha}. \quad (16)$$

This differential equation is integrated by

$$Z_q(\alpha) = K_{\lambda}(q)\alpha^{1-8\lambda}, \quad (17)$$

with $K_{\lambda}(q)$ an integration constant. This makes sense for $q = 1$ (direct computation gives $K_0(1) = 1$, $K_{1/8}(1) = -i\pi/2$), and thus gives the relative motion of the cosmic strings for all $\alpha(t, z)$. Comparing with (13), one obtains the relation between the integration constants C_{λ} and K_{λ}

$$C_{\lambda}(\alpha, 0) = \frac{1 - 8\lambda}{8\lambda} K_{\lambda}(1) \alpha^{-1-8\lambda}. \quad (18)$$

This dependence of $C_\lambda(\alpha, 0)$ on α can also be directly obtained from the definition (9) of I by a scaling argument.

Also, the relative string motion is, contrary to appearance, not arbitrary. In the special case of parallel strings, the system can be trivially reduced (by omitting the dz^2 in (3)) to a (2+1)-dimensional system of conical singularities. As conical singularities correspond to point particles, one must require for consistency [4] that these follow geodesics of the spacetime (this approach was used successfully in [5, 6] to study the dynamics of systems of black holes and conical singularities). More generally, it has been shown that the world-sheets of self-gravitating cosmic strings are totally geodesic submanifolds [7, 8]. In the present case, the spacetime is flat outside the cosmic strings, and the solution of the geodesic equations for the $Z_i \equiv Z(\zeta = \alpha_i)$ is straightforwardly given in terms of initial data Z_{0i} , v_i and w_i by

$$Z_i(t, z) = Z_{0i} + v_i t + w_i z. \quad (19)$$

In the case of our example of two strings in the center-of-mass frame, comparison of (17) and (19) leads to the solution

$$\alpha(t, z) = (\beta t + \gamma z + \delta)^{1/(1-8\lambda)}, \quad (20)$$

with β , γ and δ (initial “velocities”, “inclinations” and “positions”) complex integration constants (proportional to v , w , and Z_0).

The relative motion of two cosmic strings has been reduced, via (1), (7) and (20), to the (1+1)-dimensional free motion of a point particle. The world-line of this particle may always be Lorentz-transformed to the time axis in the case of a subluminal velocity ($v^2 < w^2$), or to the z -axis in the case of a superluminal velocity ($v^2 > w^2$). Correspondingly, it follows that the two-cosmic string system may be Lorentz-transformed to a static system of two crossed strings in the first case, or to a system of two moving parallel strings in the second case, in accordance with the results of [9].

Thus the metric (3) indeed describes a system of infinite straight strings arbitrarily oriented and moving with respect to each other. The strings are gravitationally interacting, so one could expect gravitational radiation to occur. But this space-time does not contain gravitational waves and does not exhibit radiative features. In the case of parallel strings, this can be easily understood as a consequence of the absence of gravitons in $2 + 1$ gravity. For two crossed strings undergoing a superluminal (subluminal) collision this follows from the possibility of finding a Lorentz frame in which the strings are moving parallel (static). In the limiting case of a “luminal” collision the absence of gravitational radiation follows from the existence of the above exact solution.

It is worth noting that this simplicity is lost once other string interactions are taken into account. Two crossed strings moving with a superluminal relative velocity do radiate a massless second rank antisymmetric tensor field (string axion) if the strings are charged with respect to this field. This effect was recently considered using perturbation theory [10]. Therefore a system of strings interacting with a two-form field can hardly be described by a simple solution like (3).

References

- [1] M.R. Anderson, “The Mathematical Theory of Cosmic Strings”, Institute of Physics Publishing, Bristol 2003.
- [2] J.R. Gott III, Phys. Rev. Lett. **66** (1991) 1126.
- [3] P.S. Letelier and D.V. Gal’tsov, Class. Quantum Grav. **10** (1993) L101.
- [4] S. Deser and R. Jackiw, Ann. Phys. (N.Y.) **153** (1984) 405.
- [5] G. Clément, Phys. Rev. D **50** (1994) R7119.
- [6] G. Clément and A. Fabbri, Class. Quantum Grav. **16** (1999) 323.
- [7] J.A.G. Vickers, Class. Quantum Grav. **4** (1987) 1.
- [8] W.G. Unruh, G. Hayward, W. Israel and D. McManus, Phys. Rev. Lett. **62** (1989) 2897.
- [9] D.V. Gal’tsov, Yu.V. Grats and P.S. Letelier, Ann. Phys. (N.Y.) **224** (1993) 90.
- [10] D.V. Gal’tsov, E.Yu. Melkumova and R. Kerner, Phys. Rev. D **70** (2004) 045009.