

Dirac Quasinormal modes of Schwarzschild black hole

Jiliang Jing*

*Institute of Physics and Department of Physics,
Hunan Normal University,
Changsha, Hunan 410081, P. R. China*

Abstract

The quasinormal modes (QNMs) associated with the decay of Dirac field perturbation around a Schwarzschild black hole is investigated by using continued fraction and Hill-determinant approaches. It is shown that the fundamental quasinormal frequencies become evenly spaced for large angular quantum number and the spacing is given by $\omega_{\lambda+1} - \omega_{\lambda} = 0.38490 - 0.00000i$. The angular quantum number has the surprising effect of increasing real part of the quasinormal frequencies, but it almost does not affect imaginary part, especially for low overtones. In addition, the quasinormal frequencies also become evenly spaced for large overtone number and the spacing for imaginary part is $Im(\omega_{n+1}) - Im(\omega_n) \approx -i/4M$ which is same as that of the scalar, electromagnetic, and gravitational perturbations.

PACS numbers: 04.70.-s, 04.50.+h, 11.15.-q, 11.25.Hf

*Electronic address: jljing@hunnu.edu.cn

I. INTRODUCTION

Motivated by the beliefs that one can directly identify a black hole existence by comparing QNMs with the gravitational waves observed in the universe, and the study of QNMs may lead to a deeper understanding of the thermodynamic properties of black holes in loop quantum gravity [1] [2], as well as the QNMs of anti-de Sitter black holes have a direction interpretation in terms of the dual conformal field theory [3] [4] [5], the investigation of QNMs in black hole spacetimes becomes appealing recent years. Since Regge and Wheeler [6] first presented the idea of the QNMs and Chandrasekhar and Detweller [7] worked out the numerical calculation of the QNMs for the Schwarzschild black hole, a great deal of effort has been contributed to compute the QNMs of the black holes [10]-[24] for fields of integer spin, such as the scalar, electromagnetic and gravitational perturbations.

On the contrary, the study of the QNMs of the Dirac field is very limited [25][26][27]. Cho [25] studied the QNMs of the Dirac field of the Schwarzschild black hole with the help of the third-order WKB method, and Zhidenko [26] extended the study to the case of the Schwarzschild-de Sitter black hole by using the sixth-order WKB method. We [27] investigated the Dirac QNMs of the Reissner-Nordström de Sitter black hole using the Pöshl-Teller potential approximation. Nevertheless, in these papers the study of the Dirac QNMs were limited by using WKB or Pöshl-Teller potential approximative methods. The reason to obstruct study of the Dirac QNMs is that for static black holes the standard wave-equations

$$\left(\frac{d^2}{dr_*^2} + \omega^2\right) Z_{\pm} = V_{\pm} Z_{\pm} \quad (1.1)$$

possesses the special potentials

$$V_{\pm} = \lambda^2 \frac{\Delta}{r^4} \pm \lambda \frac{d}{dr_*} \frac{\sqrt{\Delta}}{r^2} \quad (1.2)$$

which are the functions of $\sqrt{\Delta}$ [25][27][28], where Δ is a function related to metric, say $\Delta = r^2 - 2Mr$ for the Schwarzschild black hole. So, it is very hard to calculate Dirac QNMs by using the known numerical methods (except WKB or Pöshl-Teller approximative methods) since we have to expand the potentials as a series at boundary of the field, such as at the event horizon. However, Castello-Branco *et al* [29] evaluated the Dirac QNMs of the Schwarzschild black hole by using the continued fraction approach in which they expanded the wave function as a series of $\sqrt{\Delta}/r$, and they found that the spacing for imaginary part is not $-i/4M$, as it takes place for scalar, electromagnetic, and gravitational perturbations, but $-i/8M$.

Recently, we [30][31] found that the wave functions and potentials of the Dirac field in the static spacetimes can be expressed as new forms, and the new wave functions are related to the standard wave functions Z_{\pm} in Eq. (1.1) in a simple way [30]. Starting from the new wave

functions and potentials, we [30][31] showed that the Dirac QNMs of the Schwarzschild anti-de Sitter and Reissner-Norström anti-de Sitter black holes can easily be worked out by using Horowitz-Hubeny approach [15]. We have checked that the new potentials also present correct quasinormal frequencies for the scalar and gravitational perturbations in the Schwarzschild black hole.

The main purpose of this paper is to study Dirac QNMs of the Schwarzschild black hole by using the new kinds of wave functions and potentials with the help of the continued fraction and Hill-determinant approaches. The results obtained here are different from Castello-Branco's one [29] (If taking $M = 1$, we can only obtain results $n = 0, 2, \dots$ in Ref. [29]. That is to say we do not find "specific modes"). We are sure that we do not miss any quasinormal frequency because the Hill-determinant approach gives all roots for $n \leq 15$.

The organization of this paper is as follows. In Sec. 2 the decoupled Dirac equations and corresponding wave equations in the Schwarzschild spacetime are obtained by using Newman-Penrose formalism. In Sec. 3 the numerical approach to computing the Dirac QNMs is introduced. In Sec. 4 the numerical results for the Dirac QNMs in the Schwarzschild black hole are presented. The last section is devoted to a summary.

II. DIRAC EQUATIONS IN THE SCHWARZSCHILD SPACETIME

The Dirac equations [32] are

$$\begin{aligned}\sqrt{2}\nabla_{BB'}P^B + i\mu\bar{Q}_{B'} &= 0, \\ \sqrt{2}\nabla_{BB'}Q^B + i\mu\bar{P}_{B'} &= 0,\end{aligned}\tag{2.1}$$

where $\nabla_{BB'}$ is covariant differentiation, P^B and Q^B are the two-component spinors representing the wave function, $\bar{P}_{B'}$ is the complex conjugate of P_B , and μ is the particle mass. In the Newman-Penrose formalism [33] the equations become

$$\begin{aligned}(D + \epsilon - \rho)P^0 + (\bar{\delta} + \pi - \alpha)P^1 &= 2^{-1/2}i\mu\bar{Q}^{1'}, \\ (\Delta + \mu - \gamma)P^1 + (\delta + \beta - \tau)P^0 &= -2^{-1/2}i\mu\bar{Q}^{0'}, \\ (D + \bar{\epsilon} - \bar{\rho})\bar{Q}^{0'} + (\delta + \bar{\pi} - \bar{\alpha})\bar{Q}^{1'} &= -2^{-1/2}i\mu P^1, \\ (\Delta + \bar{\mu} - \bar{\gamma})\bar{Q}^{1'} + (\bar{\delta} + \bar{\beta} - \bar{\tau})\bar{Q}^{0'} &= 2^{-1/2}i\mu P^0.\end{aligned}\tag{2.2}$$

For the Schwarzschild spacetime

$$ds^2 = \left(1 - \frac{2M}{r}\right) dt^2 - \frac{1}{1 - \frac{2M}{r}} dr^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2),\tag{2.3}$$

where M represents the black hole mass, the null tetrad can be taken as

$$\begin{aligned} l^\mu &= \left(\frac{r^2}{\Delta}, 1, 0, 0 \right), \\ n^\mu &= \frac{1}{2} \left(1, -\frac{\Delta}{r^2}, 0, 0 \right) \\ m^\mu &= \frac{1}{\sqrt{2}r} \left(0, 0, 1, \frac{i}{\sin\theta} \right), \end{aligned} \quad (2.4)$$

with

$$\Delta = r^2 - 2Mr. \quad (2.5)$$

Then, if we set

$$\begin{aligned} P^0 &= \frac{1}{r} \mathbb{R}_{-1/2}(r) S_{-1/2}(\theta) e^{-i(\omega t - \bar{m}\varphi)}, \\ P^1 &= \mathbb{R}_{+1/2}(r) S_{+1/2}(\theta) e^{-i(\omega t - \bar{m}\varphi)}, \\ \bar{Q}^{1'} &= \mathbb{R}_{+1/2}(r) S_{-1/2}(\theta) e^{-i(\omega t - \bar{m}\varphi)}, \\ \bar{Q}^{0'} &= -\frac{1}{r} \mathbb{R}_{-1/2}(r) S_{+1/2}(\theta) e^{-i(\omega t - \bar{m}\varphi)}, \end{aligned} \quad (2.6)$$

where ω and \bar{m} are the energy and angular momentum of the Dirac particles, after tedious calculation Eq. (2.2) can be simplified as

$$\sqrt{\Delta} \mathcal{D}_0 \mathbb{R}_{-1/2} = (\lambda + i\mu r) \sqrt{\Delta} \mathbb{R}_{+1/2}, \quad (2.7)$$

$$\sqrt{\Delta} \mathcal{D}_0^\dagger (\sqrt{\Delta} \mathbb{R}_{+1/2}) = (\lambda - i\mu r) \mathbb{R}_{-1/2}, \quad (2.8)$$

$$\mathcal{L}_{1/2} S_{+1/2} = -\lambda S_{-1/2}, \quad (2.9)$$

$$\mathcal{L}_{1/2}^\dagger S_{-1/2} = \lambda S_{+1/2}. \quad (2.10)$$

with

$$\begin{aligned} \mathcal{D}_n &= \frac{\partial}{\partial r} - \frac{iK}{\Delta} + \frac{n}{\Delta} \frac{d\Delta}{dr}, \\ \mathcal{D}_n^\dagger &= \frac{\partial}{\partial r} + \frac{iK}{\Delta} + \frac{n}{\Delta} \frac{d\Delta}{dr}, \\ \mathcal{L}_n &= \frac{\partial}{\partial \theta} + \frac{\bar{m}}{\sin\theta} + n \cot\theta, \\ \mathcal{L}_n^\dagger &= \frac{\partial}{\partial \theta} - \frac{\bar{m}}{\sin\theta} + n \cot\theta, \\ K &= r^2 \omega. \end{aligned} \quad (2.11)$$

We can eliminate $S_{+1/2}$ (or $S_{-1/2}$) from Eqs. (2.9) and (2.10) and obtain

$$\left[\frac{1}{\sin\theta} \frac{d}{d\theta} \left(\sin\theta \frac{d}{d\theta} \right) - \frac{\bar{m}^2 + 2\bar{m}s\cos\theta + s^2\cos^2\theta}{\sin^2\theta} + s + A_s \right] S_s = 0, \quad (2.12)$$

here and hereafter we take $s = +1/2$ for the case $S_{+1/2} (\mathbb{R}_{+1/2})$ and $s = -1/2$ for $S_{-1/2} (\mathbb{R}_{-1/2})$, and $A_{+1/2} = \lambda^2 - 2s$ and $A_{-1/2} = \lambda^2$. The angular equation (2.12) can be solved exactly and $A_s = (l - s)(l + s + 1)$, where l is the quantum number characterizing the angular distribution. So, for both cases $s = +1/2$ and $s = -1/2$ we have

$$\lambda^2 = \left(l + \frac{1}{2} \right)^2. \quad (2.13)$$

We will focus our attention on the massless case in this paper. Therefore, we can eliminate $\mathbb{R}_{-1/2}$ (or $\sqrt{\Delta}\mathbb{R}_{+1/2}$) from Eqs. (2.7) and (2.8) to obtain a radial decoupled Dirac equation for $\sqrt{\Delta}\mathbb{R}_{+1/2}$ (or $\mathbb{R}_{-1/2}$). Then, Introducing an usual tortoise coordinate

$$dr_* = \frac{r^2}{\Delta} dr \quad (2.14)$$

and resolving the equation in the form

$$\mathbb{R}_s = \frac{\Delta^{-s/2}}{r} \Psi_s, \quad (2.15)$$

we obtain the wave equation

$$\frac{d\Psi_s}{dr_*^2} + (\omega^2 - V_s)\Psi_s = 0, \quad (2.16)$$

where

$$V_s = -\frac{\Delta}{4r^2} \frac{d}{dr} \left[r^2 \frac{d}{dr} \left(\frac{\Delta}{r^4} \right) \right] + \frac{s^2 r^4}{4} \left[\frac{d}{dr} \left(\frac{\Delta}{r^4} \right) \right]^2 + is\omega r^2 \frac{d}{dr} \left(\frac{\Delta}{r^4} \right) + \frac{\lambda^2 \Delta}{r^4}. \quad (2.17)$$

Although the form of the potential is more complicated than standard wave-equations $\left(\frac{d^2}{dr_*^2} + \omega^2 \right) Z_{\pm} = V_{\pm} Z_{\pm}$ with the potentials $V_{\pm} = \lambda^2 \frac{\Delta}{r^4} \pm \lambda \frac{d}{dr_*} \frac{\sqrt{\Delta}}{r^2}$ [25][27][28], we will see that we can easily work out the Dirac quasinormal frequencies of the Schwarzschild black hole from Eqs. (2.16) and (2.17) by using continued fraction and Hill-determinant approaches.

III. NUMERICAL APPROACHES TO COMPUTING DIRAC QUASINORMAL MODES

For the Schwarzschild black hole, the QNMs are defined to be modes with purely ingoing waves at the event horizon and purely outgoing waves at infinity [7]. Then, the boundary conditions on wave function Ψ_s at the horizon ($r = r_+$) and infinity ($r \rightarrow +\infty$) can be expressed as

$$\Psi_s \sim \begin{cases} (r - r_+)^{-\frac{s}{2} - \frac{i\omega}{2\kappa}} & r \rightarrow r_+, \\ r^{-s+i\omega} e^{i\omega r} & r \rightarrow +\infty, \end{cases} \quad (3.1)$$

where κ is the surface gravity of the black hole.

A solution to Eq. (2.16) that has the desired behaviour at the boundary can be written in the form

$$\Psi_s = r^{-\frac{s}{2}+2i\omega}(r-r_+)^{-\frac{s}{2}-\frac{i\omega}{2\kappa}}e^{i\omega r}\sum_{m=0}^{\infty}a_m\left(\frac{r-r_+}{r}\right)^m \quad (3.2)$$

If we take $r_+ = 1$, the sequence of the expansion coefficient $\{a_m : m = 1, 2, \dots\}$ is determined by a three-term recurrence relation starting with $a_0 = 1$:

$$\begin{aligned} \alpha_0 a_1 + \beta_0 a_0 &= 0, \\ \alpha_m a_{m+1} + \beta_m a_m + \gamma_m a_{m-1} &= 0, \quad m = 1, 2, \dots \end{aligned} \quad (3.3)$$

The recurrence coefficient α_m , β_m , and γ_m are given in terms of m and the black hole parameters by

$$\begin{aligned} \alpha_m &= m^2 + (C_0 + 1)m + C_0, \\ \beta_m &= -2m^2 + (C_1 + 2)m + C_3, \\ \gamma_m &= m^2 + (C_2 - 3)m + C_4 - C_2 + 2, \end{aligned} \quad (3.4)$$

with

$$\begin{aligned} C_0 &= 1 - s - 2i\omega, \\ C_1 &= -4 + 8i\omega, \\ C_2 &= s + 3 - 4i\omega, \\ C_3 &= 8\omega^2 - s - 1 + 4i\omega - \lambda^2, \\ C_4 &= s + 1 - 4\omega^2 - (2s + 4)i\omega. \end{aligned} \quad (3.5)$$

The three terms recurrence relation (3.3) can be used to find quasinormal frequencies by using both the continued fraction and Hill-determinant methods which are introduced in the following.

[The continued fraction approach]: The ratio of the series coefficients $\{a_m : m = 1, 2, \dots\}$ is finite and can be determined in two ways:

$$\begin{aligned} \frac{a_{m+1}}{a_m} &= \frac{\gamma_m}{\alpha_m} \frac{\alpha_{m-1}}{\beta_{m-1-}} \frac{\alpha_{m-2}\gamma_{m-1}}{\beta_{m-2-}} \frac{\alpha_{m-3}\gamma_{m-2}}{\beta_{m-3-}} \dots - \frac{\beta_m}{\alpha_m} \\ &= \frac{-\gamma_{m+1}}{\beta_{m+1-}} \frac{\alpha_{m+1}\gamma_{m+2}}{\beta_{m+2-}} \frac{\alpha_{m+2}\gamma_{m+3}}{\beta_{m+3-}} \dots \end{aligned} \quad (3.6)$$

Using Eqs. (3.3) and (3.6) we can obtain an equality between two continued fractions, one of infinite length, and the other finite:

$$\left[\beta_m - \frac{\alpha_{m-1}\gamma_m}{\beta_{m-1-}} \frac{\alpha_{m-2}\gamma_{m-1}}{\beta_{m-2-}} \dots \frac{\alpha_0\gamma_1}{\beta_0} \right] = \left[\frac{\alpha_m\gamma_{m+1}}{\beta_{m+1-}} \frac{\alpha_{m+1}\gamma_{m+2}}{\beta_{m+2-}} \frac{\alpha_{m+2}\gamma_{m+3}}{\beta_{m+3-}} \dots \right], \quad (m = 1, 2, \dots). \quad (3.7)$$

at values of order N (for giving N , one require solving polynomials of the order of $2N + 2$ in ω) very much higher than the overtone number n of the quasinormal modes. Also, the method does not have any control over the stability of the root search algorithm.

We have checked that both the methods present the same results for $\lambda = 5$ to $\lambda = 10$ and $n = 0$ to $n = 15$.

IV. DIRAC QUASINORMAL MODES OF THE SCHWARZSCHILD BLACK HOLE

In this section we represent the numerical results of the Dirac quasinormal frequencies of the Schwarzschild black hole obtained by using the numerical approaches just outlined in the previous section. The results will be organized into two subsections: the fundamental quasinormal modes and high overtones.

A. Fundamental quasinormal modes

The fundamental quasinormal frequencies ($n=0$) for $\lambda = 1$ to $\lambda = 40$ are listed in the table (I). Fig. (1) shows that $\Delta\omega = \omega_{\lambda+1} - \omega_{\lambda}$ as a function of λ . From the table and figure we know that the fundamental quasinormal frequencies become evenly spaced for large λ and the spacing is given by

$$\Delta\omega = 0.38490 - 0.00000i. \quad (4.1)$$

B. High overtones

The Dirac quasinormal frequencies of the Schwarzschild black hole for $\lambda = 1$ to $\lambda = 5$ and $n = 1$ to $n = 20$ are given by table (II) and the first 21 Dirac quasi-normal frequencies for $\lambda = 1$ to $\lambda = 10$ are described by Fig. (2).

The dependence of the quasinormal frequencies on λ for the Schwarzschild black hole is described by Figs. (3) and (4). We learn from Figs. (3) and (4) that λ has the surprising effect of increasing real part of the quasinormal frequencies, but it almost does not affect imaginary part, especially for the lower overtones.

From table (II) and Fig. (5) we know that the Dirac QNMs for $\lambda = 1, 2$ demonstrate the following asymptotic behavior

$$Re(\omega_{n+1}) - Re(\omega_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (4.2)$$

$$Im(\omega_{n+1}) - Im(\omega_n) \approx -\frac{i}{4M} \quad \text{as } n \rightarrow \infty. \quad (4.3)$$

TABLE I: The fundamental Dirac quasinormal frequencies of the Schwarzschild black hole for $\lambda = 1$ to $\lambda = 40$.

λ	ω	λ	ω	λ	ω	λ	ω
1	0.365926-0.193965i	11	4.23212-0.192462i	21	8.08197-0.192453i	31	11.93128-0.192452i
2	0.760074-0.19281i	12	4.61717-0.192462i	22	8.46691-0.192453i	32	12.31620-0.192451i
3	1.14819-0.19261i	13	5.00219-0.192459i	23	8.85185-0.192453i	33	12.70112-0.192451i
4	1.53471-0.19254i	14	5.38720-0.192457i	24	9.23679-0.192453i	34	13.08604-0.192451i
5	1.92059-0.192507i	15	5.77220-0.192456i	25	9.62172-0.192452i	35	13.47096-0.192451i
6	2.30614-0.192490i	16	6.15718-0.192456i	26	10.00667-0.192452i	36	13.85588-0.192451i
7	2.69150-0.192479i	17	6.54215-0.192455i	27	10.39160-0.192452i	37	14.24080-0.192451i
8	3.07675-0.192473i	18	6.92711-0.192455i	28	10.77653-0.192452i	38	14.62572-0.192451i
9	3.46192-0.192468i	19	7.31207-0.192454i	29	11.16144-0.192452i	39	15.01064-0.192451i
10	3.84704-0.192464i	20	7.69702-0.192454i	30	11.54636-0.192452i	40	15.39554-0.192451i




FIG. 1: The spacing $\Delta\omega = \omega_{\lambda+1} - \omega_{\lambda}$ as the functions of λ for the fundamental quasinormal frequencies. The top figure is drawn for $\Delta\omega_{Re} = Re(\omega_{\lambda+1}) - Re(\omega_{\lambda})$ which shows that the spacing of the real part is 0.38490 for large λ . The bottom one is for $\Delta\omega_{Im} = Im(\omega_{\lambda+1}) - Im(\omega_{\lambda})$ which shows that the spacing of the imaginary part becomes zero for large λ .

TABLE II: Dirac quasinormal frequencies of the Schwarzschild black hole for $\lambda = 1$ to $\lambda = 5$ and $n = 1$ to $n = 20$.

n	ω ($\lambda = 1$)	ω ($\lambda = 2$)	ω ($\lambda = 3$)	ω ($\lambda = 4$)	ω ($\lambda = 5$)
1	0.295644-0.633857i	0.711666-0.594995i	1.11403-0.58543i	1.50860-0.581936i	1.89952-0.580296i
2	0.240740-1.128450i	0.638523-1.03691i	1.05321-0.99939i	1.45954-0.98382i	1.85899-0.976231i
3	0.208512-1.63397i	0.569867-1.5149i	0.97949-1.44176i	1.39383-1.40459i	1.80226-1.38504i
4	0.187528-2.14013i	0.516459-2.01144i	0.906879-1.91029i	1.3199-1.84711i	1.73409-1.81009i
5	0.17248-2.64556i	0.476221-2.51504i	0.843213-2.3969i	1.24587-2.3100i	1.66006-2.25269i
6	0.160962-3.15021i	0.445224-3.0209i	0.790087-2.89386i	1.17721-2.78892i	1.58544-2.71190i
7	0.151745-3.65419i	0.420575-3.52725i	0.746214-3.39616i	1.11627-3.27894i	1.51418-3.18513i
8	0.144128-4.15764i	0.400401-4.03343i	0.70972-3.90102i	1.06324-3.77605i	1.44849-3.66916i
9	0.13768-4.66067i	0.383487-4.53924i	0.678957-4.40701i	1.01733-4.27745i	1.38920-4.16097i
10	0.132118-5.16336i	0.369026-5.0446i	0.65265-4.91341i	0.97747-4.78135i	1.33621-4.65815i
11	0.12725-5.66576i	0.356461-5.54954i	0.629847-5.41982i	0.942637-5.28666i	1.28901-5.15892i
12	0.122937-6.16792i	0.345399-6.05409i	0.60984-5.92607i	0.911955-5.79270i	1.24690-5.66202i
13	0.119077-6.66989i	0.335551-6.55828i	0.59210-6.43207i	0.884707-6.29908i	1.20921-6.16662i
14	0.115592-7.17170i	0.326701-7.06215i	0.57622-6.93778i	0.860321-6.80556i	1.17532-6.67215i
15	0.112424-7.67335i	0.318685-7.56574i	0.56189-7.44319i	0.838336-7.31197i	1.14467-7.17822i
16	0.109525-8.17489i	0.311372-8.06907i	0.548867-7.94831i	0.818386-7.81825i	1.11682-7.68457i
17	0.106858-8.67631i	0.304662-8.57218i	0.536959-8.45316i	0.800173-8.32435i	1.09138-8.19105i
18	0.104393-9.17764i	0.298471-9.07509i	0.526008-8.95774i	0.783457-8.83023i	1.06803-8.69753i
19	0.102104-9.67888i	0.292733-9.57782i	0.515889-9.46207i	0.768038-9.33589i	1.04651-9.20394i
20	0.099979-10.1800i	0.287392-10.0804i	0.506497-9.96618i	0.753754-9.84132i	1.02661-9.71027i

It is shown that the spacing for imaginary part is $-i/4M$ which is same as that of the scalar, electromagnetic, and gravitational perturbations.

V. SUMMARY

The wave equations for the Dirac fields in the Schwarzschild black hole spacetime are obtained by means of the Newman-Penrose formulism. Then, the quasinormal frequencies cor-




FIG. 2: First 21 Dirac quasi-normal frequencies of the Schwarzschild black hole for $\lambda = 1$ to $\lambda = 10$.




FIG. 3: Dependence of the real parts of the quasinormal frequencies on λ for the Schwarzschild black hole, which shows that λ has the surprising effect of increasing $Re(\omega)$.

responding to the Dirac field perturbation in the black hole spacetime are evaluated by using continued fraction and Hill-determinant approaches. Three interesting results obtained in this paper are listed in the following: (i) The fundamental quasinormal frequencies become evenly spaced for large λ and the spacing is given by $\omega_{\lambda+1} - \omega_{\lambda} = 0.38490 - 0.00000i$. (ii) λ has the surprising effect of increasing real part of the quasinormal frequencies, but it almost does not affect imaginary part, especially for the low overtones. (iii) The quasinormal frequencies also become evenly spaced for large n and the spacing for imaginary part is $-i/4M$, which is same




FIG. 4: Dependence of the imaginary parts of the quasinormal frequencies on λ for the Schwarzschild black hole, which shows that λ almost does not affect imaginary part, $Im(\omega)$, especially for the lower overtones.




FIG. 5: Graphs of $\Delta\omega = \omega_{n+1} - \omega_n$ versus overtone number n for $\lambda = 1, 2$. The top figure is drawn for $\Delta\omega_{Re} = Re(\omega_{n+1}) - Re(\omega_n)$ which shows that $\Delta\omega_{Re}$ tends to zero for large n . The bottom one is for $\Delta\omega_{Im} = Im(\omega_{n+1}) - Im(\omega_n)$ which shows that $\Delta\omega_{Im} \approx -\frac{i}{4M}$ for large n .

as that of the scalar, electromagnetic, and gravitational perturbations.

In future papers in this series the new wave functions and potentials will be applied to study of Dirac QNMs in other black hole spacetimes, such as the charged Reissner-Norström black hole and the rotating Kerr black hole.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under Grant No. 10275024 and under Grant No. 10473004; the FANEDD under Grant No. 200317; and the SRFDP under Grant No. 20040542003.

-
- [1] S. Hod, Phys. Rev. Lett. **81**, 4293 (1998).
 - [2] O. Dreyer, Phys. Rev. Lett. **90**, 081301 (2003).
 - [3] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998).
 - [4] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998).
 - [5] S. Kalyana Rama and Sathiapalan, Mod Phys. Lett. A **14**, 2635 (1999).
 - [6] T. Regge and J. A. Wheeler, Phys. Rev. **108** 1063 (1957).
 - [7] S. Chandrasekhar and S. Detweiler, Proc. R. Soc. Lond. **A 344** 441 (1975).
 - [8] K. Kokkotas and B. Schmidt, Living Reviews Relativ. **2** 2 (1999).
 - [9] H. P. Nollert, Class. Quant. Grav. **16**, 159 (1999).
 - [10] J. S. F. Chan and R. B. Mann, Phys. Rev. D **55** 7546 (1997).
 - [11] D. Birmingham, I. Sachs, and S. N. Solodukhin, Phys. Rev. Lett. **88** 151301 (2002).
 - [12] V. Cardoso and J. P. S. Lemos, Phys. Rev. D **63** 124015 (2001).
 - [13] R. A. Konoplya, Phys. Rev. D **66** 084007 (2002).
 - [14] A. O. Starinets, Phys. Rev. D **66** 124013 (2002).
 - [15] G. T. Horowitz and V. E. Hubeny, Phys. Rev. D **62** 024027 (2000).
 - [16] M. R. Setare, Class. Quant. Grav. **21** 1453 (2003).
 - [17] M. R. Setare, Phys. Rev. D **69** 044016 (2004).
 - [18] J. Natario and R. Schiappa, hep-th/0411267.
 - [19] V. Cardoso, J. Natario, and R. Schiappa, hep-th/0403132.
 - [20] Y. Kurita and M. A. Sakagami, Phys. Rev. D **67** 024003 (2003).

- [21] V. Cardoso and J. P. S. Lemos, Phys. Rev. D **64** 084017 (2001).
- [22] V. Cardoso, R. Konoplya, and J. P. S. Lemos, gr-qc/0305037.
- [23] V. Cardoso and J. P. S. Lemos, Phys. Rev. D **63** 124015 (2001).
- [24] R. A. Konoplya, Phys. Rev. D **66** 044009 (2002).
- [25] H. T. Cho, Phys. Rev. D **68**, 024003 (2003).
- [26] A. Zhidenko, Class. Quant. Grav. **21** 273 (2004).
- [27] Jiliang Jing, Phys. Rev. D **69**, 084009 (2004).
- [28] Jiliang Jing, Phys. Rev. D **70**, 065004 (2004).
- [29] K. H. C. Castello-Branco, R. A. Konoplya, and A. Zhidenko, hep-th/0411055, accepted by Phys. Rev. D.
- [30] Jiliang Jing, gr-qc/0502010, Submitted to Phys. Rev. D.
- [31] Jiliang Jing, gr-qc/0502011, Submitted to Phys. Rev. D.
- [32] D. N. Page, Phys. Rev. D **14**, 1509 (1976).
- [33] E. Newman and R. Penrose, J. Math. Phys. (N. Y.) **3**, 566 (1962).
- [34] E. W. Leaver, Pro. R. Soc. Lond. **A** 402, 285 (1985); E. W. Leaver, Phys. Rev. D **34**, 384 (1986).
- [35] B. Majumdar and N. Panchapakesan, Phys. Rev. D **40**, 2568 (1989).
- [36] E. W. Leaver, Phys. Rev. D **41**, 2986 (1990).