

Testing frame-dragging with the Mars Global Surveyor spacecraft in the gravitational field of Mars

Lorenzo Iorio
*Viale Unità di Italia 68, 70125
Bari (BA), Italy
tel. 0039 328 6128815
e-mail: lorenzo.iorio@libero.it*

Abstract

In this paper we propose an interpretation of the time series of the RMS overlap differences of the out-of-plane part N of the orbit of the Martian polar orbiter Mars Global Surveyor (MGS) over a time span ΔP of about 5 years (14 November 1999-14 January 2005) in terms of the general relativistic gravitomagnetic Lense-Thirring effect. It turns out that the average of such a time series over ΔP , normalized to the predicted Lense-Thirring out-of-plane mean shift over the same time span, is $\mu = 1.0018 \pm 0.0053$. A linear fit of the considered time series over ΔP yields a normalized slope 1.1 ± 0.3 strengthening our interpretation of the determined MGS out-of-plane features of motion as due to the gravitomagnetic force of Mars.

Keywords: gravity tests; Lense-Thirring effect; Mars Global Surveyor

PACS: 04.80.-y, 04.80.Cc, 95.10.Ce, 95.55.Pe, 96.30.Gc, 96.12.Fe

1 The gravitomagnetic Lense-Thirring effect

In the framework of the weak-field and slow-motion linearized approximation of general relativity [31], the gravitomagnetic Lense-Thirring effect consists of tiny secular precessions of the longitude of the ascending node Ω and the argument of pericentre ω of the orbit of a test particle freely moving in the gravitational field of a slowly rotating body of mass M and angular momentum \mathbf{J} [27, 2, 9, 46, 1, 13]

$$\begin{cases} \dot{\Omega}_{\text{LT}} &= \frac{2GJ}{c^2 a^3 (1-e^2)^{3/2}}, \\ \dot{\omega}_{\text{LT}} &= -\frac{6GJ \cos i}{c^2 a^3 (1-e^2)^{3/2}}, \end{cases} \quad (1)$$

where G is the Newtonian gravitational constant, c is the speed of light in vacuum, a, e and i are the semimajor axis, the eccentricity and the inclination of the orbital plane to the central body's equator, respectively. It maybe of interest to recall that de Sitter in [10] worked out, for the first time, the gravitomagnetic pericentre precession in the case of equatorial orbits ($i = 0$ deg). In general, the gravitomagnetic force affects the in-plane transverse and the out-of-plane cross-track portions of the orbit of a test particle. Indeed, the shifts along the radial R , transverse T and normal¹ N directions can be, in general, expressed in terms of the Keplerian orbital elements as [3]

$$\begin{cases} \Delta R &= \sqrt{(\Delta a)^2 + \frac{[(e\Delta a + a\Delta e)^2 + (ae\Delta \mathcal{M})^2]}{2}}, \\ \Delta T &= a\sqrt{1 + \frac{e^2}{2}} \left[\Delta \mathcal{M} + \Delta \omega + \cos i \Delta \Omega + \sqrt{(\Delta e)^2 + (e\Delta \mathcal{M})^2} \right], \\ \Delta N &= a\sqrt{\left(1 + \frac{e^2}{2}\right) \left[\frac{(\Delta i)^2}{2} + (\sin i \Delta \Omega)^2 \right]}, \end{cases} \quad (2)$$

where \mathcal{M} is the mean anomaly. The Lense-Thirring shifts are, thus

$$\begin{cases} \Delta R_{\text{LT}} &= 0, \\ \Delta T_{\text{LT}} &= a\sqrt{1 + \frac{e^2}{2}} (\Delta \omega_{\text{LT}} + \cos i \Delta \Omega_{\text{LT}}), \\ \Delta N_{\text{LT}} &= a\sqrt{1 + \frac{e^2}{2}} \sin i \Delta \Omega_{\text{LT}}. \end{cases} \quad (3)$$

For an equatorial orbit there is only the transverse shift ΔT_{LT} , while for a polar ($i = 90$ deg) satellite only the out-of-plane shift ΔN_{LT} is present. This fact can be more intuitively grasped in the following way. In general, a time-variation of the argument of pericentre ω , which is an angle entirely residing in the orbital plane delimited by the line of the nodes and the apsidal line, can be obtained by subtracting the projection of the nodal motion to the orbital plane, i.e. $\cos i \dot{\Omega}$, from the precession of the longitude of the pericentre ϖ . For a polar orbital geometry the projection of the node's motion to the orbital plane is zero. According to the well-known Gauss variation equations [43], the remaining shift induced by a small perturbing acceleration \mathbf{A} is due to its in-plane radial A_R and transverse A_T components. The

¹It is customarily defined to be positive along the orbital angular momentum.

Lense-Thirring effect is due to a Lorentz-like acceleration [28]

$$\mathbf{A}_{\text{LT}} = -2 \left(\frac{\mathbf{v}}{c} \right) \times \mathbf{B}_g; \quad (4)$$

the gravitomagnetic field [28]

$$\mathbf{B}_g = \frac{G[3\mathbf{r}(\mathbf{r} \cdot \mathbf{J}) - r^2\mathbf{J}]}{cr^5} \quad (5)$$

for $i = 90$ deg lies in the test particle's orbital plane which contains also \mathbf{J} , so that the net gravitomagnetic force is entirely normal to the orbital plane.

Up to now, many attempts to test gravitomagnetism were implemented in the weak-field and slow-motion scenarios of the gravitational fields of some Solar System's bodies. Lense and Thirring considered the natural satellites of Jupiter and Saturn [27], but the possibility of using the jovian galilean moons is not yet viable [22]. In regard to the Sun and its rocky planets [20], the predictions of general relativity for the Lense-Thirring perihelia precessions were, in fact, found in agreement with the estimated corrections to the secular rates of the perihelia of the inner planets [38], but the errors are still large. As far as the Earth is concerned, a test of the Lense-Thirring effect on the orbit of a test particle was reported in [6]. Their authors analyzed the data of the laser-ranged LAGEOS and LAGEOS II satellites by using an observable independently proposed in [41, 42] and [21] involving the nodes of the LAGEOS satellites. The idea of using the LAGEOS satellite and, more generally, the Satellite Laser Ranging (SLR) technique to measure the Lense-Thirring effect in the terrestrial gravitational field with the existing artificial satellites was put forth for the first time by Cugusi and Proverbio in 1977-1978 [8, 9]; tests actually started in 1996 with LAGEOS and LAGEOS II [5]. Although the outcome of the test in [6] is in agreement with the predictions of general relativity, the realistic evaluation of the obtained total accuracy raised a debate [15, 16, 17, 19, 7, 29]. The GP-B spacecraft [11, 12] was launched in April 2004 with the aim of measuring another gravitomagnetic effect, i.e. the Schiff precession [39, 44] of the spins of four superconducting gyroscopes carried onboard. Preliminary results, released in April 2007, yield a measurement uncertainty of the order of 256 – 128%; the final results should be announced within the end of 2007, with a goal of an about 13% accuracy. (<http://einstein.stanford.edu/>). Recently, a preliminary 6% LT test on the orbit of the Mars Global Surveyor (MGS) spacecraft (see Table 1 for its orbital parameters) in the gravitational field of Mars was reported in [18]; indeed, the predictions of general relativity are able to accommodate, on average, about 94% of the Root-Mean-Square (RMS) overlap differences

Table 1: Orbital parameters of MGS [24]; a is the semimajor axis, i is the inclination to the Mars’equator and e is the eccentricity.

a (km)	i (deg)	e
3796	92.86	0.0085

in the out-of-plane part of the MGS orbit over 5 years. Finally, it must be mentioned that, according to Nordtvedt [35, 37], we would already have evidences of the existence of the gravitomagnetic interaction. Indeed, the multi-decade analysis of the Moon’orbit by means of the Lunar Laser Ranging (LLR) technique yields a comprehensive test of the various parts of order $\mathcal{O}(c^{-2})$ of the post-Newtonian equation of motion. The existence of gravitomagnetism as predicted by general relativity would, thus, be inferred with high precision from the reconstruction of certain periodic terms in the lunar motion. A 0.1% accuracy is claimed in [33], but, according to Kopeikin [25], LLR would not be capable to detect gravitomagnetic effects; Murphy et al. replied in [34]. Other comments on [33] are in [4]. Also the radial motion of the LAGEOS satellite would yield another confirmation of the existence of the gravitomagnetic interaction [36].

In this paper we perform a test of the Lense-Thirring effect by improving the accuracy of the previous one with MGS [18] by means of a different data set, a refinement of the overall calculation and a realistic and conservative assessment of the error estimate. The details of the adopted strategy can be found in [18, 24]; here we will briefly recall them.

2 Using the MGS data in the gravitational field of Mars

In [24] six years of MGS Doppler and range tracking data and three years of Mars Odyssey Doppler and range tracking data were analyzed in order to obtain information about several features of the Mars gravity field summarized in the global solution MGS95J. As a by-product, also the orbit of MGS was determined with great accuracy. In Figure 3 of [24] the time series of the RMS overlap differences of the radial, transverse and normal portions of its orbit over the entire mission (4 February 1999–14 January 2005) are

shown; their averages over such a time span amount to [24]

$$\left\{ \begin{array}{l} \langle \Delta R_{\text{diff}} \rangle = 15 \text{ cm}, \\ \langle \Delta T_{\text{diff}} \rangle = 1.5 \text{ m}, \\ \langle \Delta N_{\text{diff}} \rangle = 1.6 \text{ m}. \end{array} \right. \quad (6)$$

The RMS overlap differences are customarily used in satellite geodesy as indicators of the overall orbit accuracy [48], concisely accounting not only for the measurement errors, but also for all the systematic bias due to the mismodelled/unmodelled forces acting on a spacecraft [30]. Errors common to adjacent arcs are, by construction, cancelled; it is not so for effects growing in time like the one we are interested in here. Thus, any analytical calculation of possible systematic errors must ultimately cope with that is actually shown by the RMS overlap differences. It is important to note that in the suite of dynamical force models used to reduce the gathered observations general relativity is, in fact, present, but only with the gravitoelectric Schwarzschild-like part: the gravitomagnetic one is not included, so that, in general, it is fully accounted for by the RMS orbit overlap differences. From eq. (3) it turns out that only the out-of-plane portion of the orbit of a polar satellite like MGS is affected by gravitomagnetism via the Lense-Thirring effect: from eq. (1) and eq. (3) the mean gravitomagnetic normal shift, averaged over a temporal interval ΔP , is²

$$\langle \Delta N_{\text{LT}} \rangle = \frac{GJ\Delta P \sin i \sqrt{1 + \frac{e^2}{2}}}{c^2 a^2 (1 - e^2)^{3/2}}. \quad (7)$$

Moreover, no empirical out-of-plane accelerations were fitted in the data reduction process; this was only done for the along-track component of MGS orbit to account for other non-gravitational forces like thermal, solar radiation pressure and drag mismodelling. Thus, it is unlikely that the out-of-plane Lense-Thirring signal was removed from the data in the least-square procedure. Moreover, if the relativistic signature was removed so that the determined out-of-plane RMS overlap differences were only (or mainly) due to other causes like mismodelling or unmodelling in the non-gravitational forces, it is difficult to understand why the along-track RMS overlap differences (middle panel of Figure 3 of [24]) have almost the same magnitude,

²A 1/2 factor appears as a result of the time-average of a linear signal like the gravitomagnetic one over a time interval ΔP .

Table 2: Relevant parameters of Mars used in the text. The parameter $\alpha \equiv C/M\mathcal{R}^2$ is the normalized polar moment of inertia, \mathcal{R} is the reference Mars radius in MGS95J model, while \mathcal{P} is the Martian sidereal period of rotation. From such values and from $J = \alpha M\mathcal{R}^2 2\pi/\mathcal{P}$ it can be obtained $J = (1.92 \pm 0.01) \times 10^{32} \text{ kg m}^2 \text{ s}^{-1}$ for the Martian angular momentum.

GM ($10^9 \text{ m}^3 \text{ s}^{-2}$)	\mathcal{R} (km)	α	\mathcal{P} (d)
42828.37440 ± 0.00028 [24]	3396.0 [24]	0.3654 ± 0.0008 [24]	1.02595675 [45]

since the along-track component of the MGS orbit is much more affected by the non-gravitational accelerations (e.g. the atmospheric drag) than the out-of-plane one. Finally, the following consideration should further enforce our conclusion. The improved modelling of the non-gravitational forces acting on MGS introduced in [24] with respect to previous works [26, 51] in which the Lense-Thirring effect was not modelled as well, only affected in a relevant way the along-track RMS overlap differences (a factor 10 better than [26, 51]), not the normal ones (just a factor 2 better than [26, 51]).

By using a time span ΔP of slightly more than 5 years (14 November 1999–14 January 2005), so to remove the first months of the mission more affected by orbital maneuvers, high gain antenna deployment and non-gravitational perturbations, it is possible to obtain

$$\langle \Delta N_{\text{diff}} \rangle = 1.613 \text{ m.} \quad (8)$$

Let us, now, compare the result of eq. (8) with the mean gravitomagnetic shift predicted by eq. (7) over the same time span; according to our hypothesis, the Lense-Thirring force of Mars is responsible for the averaged normal signature of eq. (8). The figures of Table 2 for the Mars’ relevant parameters and those of Table 3 for the fundamental constants entering the calculation yield

$$\mu_{\text{meas}} \equiv \frac{\langle \Delta N_{\text{diff}} \rangle}{\langle \Delta N_{\text{LT}} \rangle} \Big|_{\Delta P} = 1.0018 \pm 0.0053; \quad (9)$$

according to our hypothesis, $\mu_{\text{expected}} = 1$. The error was conservatively evaluated as

$$\delta\mu \leq \delta\mu|_J + \delta\mu|_G + \delta\mu|_a, \quad (10)$$

Table 3: Values used for the defining, primary and derived constants (<http://ssd.jpl.nasa.gov/?constants#ref>).

constant	numerical value	units	reference
c	299792458	m s^{-1}	[32]
G	$(6.6742 \pm 0.0010) \times 10^{-11}$	$\text{kg}^{-1} \text{m}^3 \text{s}^{-2}$	[32]
1 mean sidereal day	86164.09054	s	[47]
1 mean sidereal year	365.25636	d	[47]

with

$$\begin{cases} \delta\mu|_J \leq \mu \left(\frac{\delta J}{J} \right) = 0.0052, \\ \delta\mu|_G \leq \mu \left(\frac{\delta G}{G} \right) = 0.0001, \\ \delta\mu|_a \leq 2\mu \left(\frac{\delta a}{a} \right) = \mathcal{O}(10^{-8}). \end{cases} \quad (11)$$

As can be noted, the major source of bias is represented by the angular momentum of Mars. Its uncertainty was evaluated as

$$\delta J \leq \delta J|_{\mathcal{R}} + \delta J|_{\alpha} + \delta J|_M, \quad (12)$$

with

$$\begin{cases} \delta J|_{\mathcal{R}} \leq 2J \left(\frac{\delta \mathcal{R}}{\mathcal{R}} \right) = 7 \times 10^{29} \text{ kg m}^2 \text{ s}^{-1}, \\ \delta J|_{\alpha} \leq J \left(\frac{\delta \alpha}{\alpha} \right) = 4 \times 10^{29} \text{ kg m}^2 \text{ s}^{-1}, \\ \delta J|_M \leq J \left[\frac{\delta(GM)}{GM} \right] = \mathcal{O}(10^{24}) \text{ kg m}^2 \text{ s}^{-1}. \end{cases} \quad (13)$$

We linearly added all the errors in a conservative way; in regard to α and GM this is particularly important because they are fit-for parameters of the global solution and, consequently, are certainly affected by correlations which were not explicitly released in [24]. It must also be noted that the uncertainties quoted in Table 2 are realistic, not the mere formal, statistical errors; indeed, the quoted uncertainty in α is, e.g., 15 times the formal one [24]. Also the formal error in GM was re-scaled [24]. We also considered an uncertainty in \mathcal{R} by assuming as a realistic measure of it the difference $\delta\mathcal{R} = 6080$ m between the reference MGS95J value for the equatorial radius quoted in Table 2 from [24] and the value 3389.92 km released in [50], instead of the error of 0.04 km quoted in [50]. In regard to the error in a , we assumed from eq. (6) $\delta a \approx 15$ cm in view of the very small eccentricity of the MGS orbit.

A further confirmation of the validity of our interpretation of the MGS out-of-plane motion in terms of the gravitomagnetic force of Mars comes from a linear³ fit of the time series of the RMS overlap differences over ΔP . Indeed, the estimated slope is $-0.69 \pm 0.20 \text{ m yr}^{-1}$, while general relativity predicts a Lense-Thirring out-of-plane rate of 0.62 m yr^{-1} , with the normal direction defined positive along the orbital angular momentum. The minus sign of the fit is due to the fact that in the analysis of [24] the MGS normal direction is, instead, defined positive in the opposite direction (Konopliv A, private communication 2007). Incidentally, let us note that should we assume such a fit as indicator of the existence of the Lense-Thirring effect, the predictions of general relativity would be confirmed within the experimental error which, however, would be larger than the one obtained by taking the mean of the time series of N over ΔP .

Another important remark concerning our measurement is the following one: the interpretation of the MGS orbital features presented in this paper and their production from the collected data were made by distinct and independent subjects in different times; moreover, the authors of [24] had not the Lense-Thirring effect in mind at all when they performed their analysis. Thus, the possibility of any sort of (un)conscious gimmicks, tricks or tweaking the data because the expected answer was known in advance [49] is a priori excluded in this test.

3 Conclusions

We conclude by summarizing our main result: the ratio μ of the mean of the time series of the cross-track N RMS overlap differences of MGS from 14 November 1999 to 14 January 2005 to the mean of the Lense-Thirring shift over the same time interval amounts to $\mu = 1.0018 \pm 0.0053$. Our interpretation of the features of the out-of-plane motion of MGS in terms of the gravitomagnetic force of Mars is enforced by a linear fit of the N time series, although the error would be, in this case, larger.

It must be noted that the present analysis is based on the so far processed data in [24]; other approaches which could fruitfully be used in the future may consist in re-processing the entire MGS data set including in the dynamical force models the gravitomagnetic force as well and checking if the average of the N time series gets smaller, and/or including a further solve-for parameter in the global least-squares solution accounting for the

³It may be just the case to note that, since the plots in Fig. 3 of [24] are semi-logarithmic, one should not visually look for a straight line in them.

Lense-Thirring effect itself.

Finally, let us note that, although we lost contacts with MGS since early November 2006, more independent tests could be, in principle, performed in the near future when a sufficient amount of data from the Odyssey and, especially, Mars Reconnaissance Orbiter (MRO) orbiters will be available and processed.

Acknowledgments

I gratefully thank A. Konopliv, NASA Jet Propulsion Laboratory (JPL), for having kindly provided me with the entire MGS data set. I am also grateful to the anonymous referees for their useful comments and suggestions. Finally, I am indebted to E. Lisi, P. Colangelo, G. Fogli, S. Stramaglia, and N. Cufaro-Petroni (INFN, Bari) for very stimulating discussions.

References

- [1] Ashby N and Allison T 1993 *Celest. Mech. Dyn. Astron.* **57** 537
- [2] Barker B M and O'Connell R F 1974 *Phys. Rev. D* **10** 1340
- [3] Christodoulidis D C, Smith D E, Williamson R G and Klosko S M 1988 *J. Geophys. Res.* **93**(B6) 6216
- [4] Ciufolini I 2007 arXiv:0704.3338v2
- [5] Ciufolini I, Lucchesi D M, Vespe F, Mandiello A 1996 *Il Nuovo Cim. A* **109** 575
- [6] Ciufolini I and Pavlis E C 2004 *Nature* **431** 958
- [7] Ciufolini I and Pavlis E C 2005 *New Astron.* **10** 636
- [8] Cugusi L and Proverbio E 1977 Paper presented at the International Symposium on Satellite Geodesy in Budapest from June 28 to July 1, 1977. *J. of Geodesy* **51** 249
- [9] Cugusi L and Proverbio E 1978 *Astron. Astrophys.* **69** 321
- [10] de Sitter W 1916 *Mon. Not. Roy. Astron. Soc.* **76** 699

- [11] Everitt C W F 1974 in: *Proc. Int. School Phys. "Enrico Fermi" Course LVI*. ed Bertotti B (New York: New Academic Press) pp. 331–360. Reprinted in: 2003 *Nonlinear Gravitodynamics* ed Ruffini R J and Sigismondi C (Singapore: World Scientific) pp 439–468.
- [12] Everitt C W F et al. 2001 in: *Gyros, Clocks, Interferometers...: Testing Relativistic Gravity in Space* ed Lämmerzahl C, Everitt C W F and Hehl F W (Berlin: Springer) pp 52–82.
- [13] Iorio L 2001 *Nuovo Cim. B* **116** 777 2001
- [14] Iorio L 2003 *Celest. Mech. Dyn. Astron.* **86** 277
- [15] Iorio L 2005a *New Astron.* **10** 603
- [16] Iorio L 2005b *New Astron.* **10** 616
- [17] Iorio L 2006a *J. of Geodesy* **80** 128
- [18] Iorio L 2006b *Class. Quantum Grav.* **23** 5451
- [19] Iorio L 2007a *Planet. Space Sci.* **55** 503
- [20] Iorio L 2007b *Planet. Space Sci.*, at press, arXiv:gr-qc/0507041v12
- [21] Iorio L and Morea A 2004 *Gen. Relativ. Gravit.* **36** 1321
- [22] Iorio L and Lainey V 2005 *Int. J. Mod. Phys. D* **14** 2039
- [23] Kaula W M 1966 *Theory of Satellite Geodesy: Applications of Satellites to Geodesy* (Waltham: Blaisdell Publishing Company)
- [24] Konopliv A S et al 2006 *Icarus* **182** 23
- [25] Kopeikin S M 2007 *Phys. Rev. Lett.*, at press, arXiv:gr-qc/0702120v2
- [26] Lemoine F G et al 2001 *J Geophys Res* **106** 23359
- [27] Lense J and Thirring H 1918 *Phys. Z.* **19** 156 translated and discussed by Mashhoon B, Hehl F W and Theiss D S 1984 *Gen. Relativ. Gravit.* **16** 711. Reprinted in: 2003 *Nonlinear Gravitodynamics* ed Ruffini R J and Sigismondi C (Singapore: World Scientific) pp 349–388
- [28] Lichtenegger H, Iorio L and Mashhoon B 2006 *Ann. Physik* **15** 868

- [29] Lucchesi D 2005 *Int. J. Mod. Phys. D* **14** 1989
- [30] Lucchesi D and Balmino G 2006 *Planet. Space Sci.* **54** 581
- [31] Mashhoon B, Gronwald F and Lichtenegger H 2001 in: *Gyros, Clocks, Interferometers...: Testing Relativistic Gravity in Space.* ed Lämmerzahl C, Everitt C W F and Hehl F W (Berlin: Springer) pp 83–108
- [32] Mohr P J and Taylor B N 2005 *Rev. Mod. Phys.* **77** 1
- [33] Murphy T, Nordtvedt K and Turyshev S 2007a *Phys. Rev. Lett.* **98** 071102
- [34] Murphy T, Nordtvedt K and Turyshev S 2007b *Phys. Rev. Lett.*, at press, arXiv:0705.0513v1
- [35] Nordtvedt K 1988a *Int. J. Theor. Phys.* **27** 1395
- [36] Nordtvedt K 1988b *Phys. Rev. Lett.* **61** 2647
- [37] Nordtvedt K 2003 in: *Nonlinear Gravitodynamics. The Lense–Thirring Effect* ed Ruffini R J and Sigismondi C (Singapore: World Scientific) pp 35–45
- [38] Pitjeva E V 2005 *Astron. Lett.* **31** 340
- [39] Pugh G E 1959 *WSEG, Research Memorandum No. 11*. Reprinted in: 2003 *Nonlinear Gravitodynamics. The Lense–Thirring Effect* ed Ruffini R J and Sigismondi C (Singapore: World Scientific) pp 414–426
- [40] Reigber Ch, et al., 2004 *J. of Geodesy*, accepted
- [41] Ries J C, Eanes R J and Tapley B D 2003a in: *Nonlinear Gravitodynamics. The Lense–Thirring Effect* ed Ruffini R J and Sigismondi C (Singapore: World Scientific) pp 201–211
- [42] Ries J C, Eanes R J, Tapley B D and Peterson G E 2003b in: *Proceedings of the 13th International Laser Ranging Workshop, NASA CP 2003-212248* ed Noomen R, Klosko S, Noll C and Pearlman M (Greenbelt: NASA Goddard) (Preprint http://cddisa.gsfc.nasa.gov/lw13/lw_proceedings.html#science)

- [43] Roy A E 2003 *Orbital Motion, Fourth edition* (Bristol: Institute of Physics Publishing)
- [44] Schiff L 1960 *Am. J. of Phys.* **28** 340
- [45] Seidelmann K P (ed) 1992 *Explanatory Supplement to the Astronomical Almanac* (Mill Valley: University Science Books)
- [46] Soffel M H 1989 *Relativity in Astrometry, Celestial Mechanics and Geodesy* (Berlin: Springer)
- [47] Standish E M 1995, in: Appenzeller I (ed), *Highlights of Astronomy*, Table 1, (Dordrecht: Kluwer Academic Publishers)
- [48] Tapley B D, Schutz B E, Born G H 2004 *Statistical Orbit Determination* (Elsevier Academic Press)
- [49] Weinberg S 1993 *Dreams of a final theory* (New York: Vintage Books)
- [50] Yoder C F 1995 in: *Global Earth Physics, A Handbook of Physical Constants, AGU Reference Shelf 1* (American Geophysical Union) Table 6
- [51] Yuan D N *et al* 2001 *J Geophys Res* **106** 23377