Reply to $\ Vorio's$ ⁸⁶high-precision m easurem ent⁰⁰ of fram e dragging with the M ars G lobal Surveyor", by K ris K rogh

Lorenzo Iorio Viale Unita di Italia 68, 70125 Bari (BA), Italy tel. 0039 328 6128815 e-mail: brenzo.iorio@ libero.it

A bstract

In this note we reply to the criticism s by K rogh concerning som e aspects of the recent fram e-dragging test perform ed by Iorio with the M ars G lobal Surveyor (MGS) spacecraft in the gravitational eld of M ars.

Keywords: gravity tests; Lense-Thirring e ect; Mars G lobal Surveyor

PACS: 04.80.-y, 04.80 C c, 91.10 Sp, 95.10 C e, 95.55 P e, 96.30 G c

1 Introduction

The remarks by K rogh (2007) to the analysis by Iorio (2007) concern with I) The observable used: Iorio (2007) would have m isinterpreted the M G S data II) The confrontation between the prediction of the gravitom agnetic Lense-Thirring shift and the data over the chosen time span P: Iorio (2007) would have incorrectly compared the 1.6 m value of the out-of-plane average orbit error released by K onoliv et al. (2006) for the entire M G S data set (4 February 1999-14 January 2005) to the Lense-T hirring shift calculated for a shorter time interval P (14 N ovem ber 1999-14 January 2005) III) The data set used: Iorio (2007) discarded som e of the initialm onths of the M G S data set IV) The im pact of the atm ospheric drag, neglected by Iorio (2007).

2 Our reply

I) The entire MGS data set was subdivided by Konopliv et al. (2006) in smaller time intervals of data called arcs, not to be confused with

three-dimensional spatial portions of trajectory, as it seems K rogh (2007) did. For MGS, the lengths of the arcs vary from 4 to 6 days, so to cover m any orbital revolutions (2 h). For each arc, am ong other things, the spacecraft position and velocity were estimated and used as starting point for a num erical propagation of the satellite motion by m eans of the dynam icalm odels which, in the case of MGS, did not include the general relativistic gravitom agnetic force. C ontiguous arcs were overlapped by an amount of 2 h, i.e. slightly more than one orbital revolution, and the RM S spacecraft position di erence am ong the predicted positions propagated from the estim ated ones in the previous arc and the estim ated positions of the subsequent arc was computed. Since the arc overlaps cover just one orbit, such RMS di erences, in fact, account for any un-m odelled/m is m odelled forces yielding secular, i.e. averaged over one orbital revolution, e ects, whatever their physical origin may be. Indeed, RMS of orbit solution overlaps are commonly used in satellite geodesy as useful and signi cant indicators of the overall orbit accuracy (Tapley et al. 2004). Conversely, they are also used to gain inform ation about system atic errors com ing from inaccurate m odelling of the forces acting on the spacecraft. For details see (Tapley et al. 2004). Of course, such a technique is insensitive to short-period e ects, i.e. having frequencies higher than the orbital one: only dynamical features of motion with timescales equal to, or larger than one orbital period can be sensed by such orbit overlap di erences. M oreover, the average orbit error h N di i of about 1.6 m does not refer to this or that particular arc overlap; instead, it com es from the m ean of the entire set of RMS orbit overlap di erences for the chosen time span P and is well representative of those un-modelled/m ism odelled forces yielding e ects which do not average out over P, as it is just the case of the Lense-Thirring signal. Tim e-varying patterns exhibiting well-de ned periodicities are, instead, mainly averaged out yielding little or no contribution to the average orbit error. Incidentally, from the above discussion about the meaning of the average orbit error, it should be apparent that it does not make sense to look for the error of the error, as, instead, seem ingly required by K rogh (2007) when he blam es Iorio (2007) for not having included the uncertainty in h N resi. Another criticism by K rogh (2007) is that the RMS overlap di erences would be unable to specify any orbital precession along with its sign. This is not correct; indeed, by prelim arily tting the time series used in Iorio (2007) over P with a straight line only one gets a slope of $0.69 \quad 0.20 \text{ m yr}^{-1}$, where the predicted

Lense-Thirring MGS out-ofplane rate (custom arily de ned positive along the MGS orbital angular momentum) amounts to 0.62 m yr 1 . The obtained m inus sign is due to the fact that K onopliv et al. (2006) de ned the norm all direction to be positive in the opposite direction of the MGS orbital angular momentum (K onopliv 2007, private communication). Incidentally, let us note that should such a linear the used as indicator of the existence of the Lense-Thirring e ect which is not an obliged choice its relativistic prediction would be fully commed within the experimental error which, however, would be larger than the 0.5% coming from the average of the signal over P. Instead, the hypothesis of a nulle ect would be rejected at more than 3 sigm a level.

- II) For the time span P chosen by Iorio (2007), the average orbit error, i.e. the mean of all the considered 348 RMS orbit overlap di erences from 14 November 1999 to 14 January 2005, amounts to 1.613 m, while the gravitom agnetic shift over the same time span amounts to 1.610 m. Thus, = 1:002. Note that here we are accurate to mm level; K onopliv et al. (2006), instead, did not go beyond the cm level (in the radial direction: $h R_{di} i = 0:15 m$); if we adopted such an approxim ation we would have had a 100% agreem ent.
- III) In regard to the choice of the data, we did not assembled an ad hoc patchwork of cut-and-paste data sets just to obtain a graceful result: our data set is continuous, sm ooth, without more or less ad hoc breaks. We simply decided to discard the rst portion of data because more a ected by various spurious elects (non-gravitational perturbations, orbitalm aneuvers, high gain antenna deployment) which would have, otherwise, introduced a 15% bias, i.e. we simply chosen the optim all experimental conditions to enhance the signal-to-noise ratio. We cannot see anything strange in that.
- IV) In regard to the assessment of the impact of the atmospheric drag on the cross-track portion of the orbit of MGS, let us note that it requires not only to consider the node , as apparently claimed by K rogh (2007), but also the inclination i according to (C hristodoulidis et al. 1988)

N = a
$$1 + \frac{e^2}{2} - \frac{(i)^2}{2} + (\sin i)^2$$
: (1)

A coording to, e.g., (M ilani et al. 1987), the perturbing acceleration A $_{\rm drag}$ due to the atm ospheric drag can be cast into the form

$$A_{drag} = \frac{1}{2} Z C_D \frac{S}{M} vv; \qquad (2)$$

where S=M is the spacecraft cross sectional area (perpendicular to the velocity) divided by its mass, C_D is the drag coe cient, is the atmospheric density (assumed to be constant over one orbital revolution), v is the satellite velocity in a planetocentric, non-rotating frame of reference and Z is a corrective coe cient accounting for the fact that the atm osphere is not at rest, but rotates with angular velocity !_A m ore or less rigidly with the planet; Z 1 for polar orbits (M ilani et al. 1987). W hile the secular, i.e. averaged over one orbital period T, drag shift on the node vanishes, it is not so for the inclination: indeed, it turns out (M ilani et al. 1987)

h ii _T
$$\frac{A_{drag}}{n^2 a} \frac{!_A}{n} + 0$$
 (e); (3)

where $n = {}^{p} \overline{GM = a^{3}}$ is the K eplerian m ean m otion. As a result, the orbital plane tends to approach the planet's equator; the terms in brackets is the ratio of the drag force to the N ew tonian m onopole. As usual in perturbation theory, a is m eant as evaluated on the unperturbed reference ellipse. Thus, the out-of-plane drag shift is from eq. (1)

h N _{drag}i a
$$\frac{h i_{drag}i}{\overline{2}}$$
: (4)

In the following we will assume that $!_{A}$ $!_{M ars} = 7:10$ 10 ⁵ s ¹. Let us see what happens if A_{drag} is of the same order of magnitude of the gravitom agnetic force exerted by M ars on MGS, i.e. 10 ¹¹ m s ². This approximation is reasonable and conservative not only because it de ness a natural threshold after which the Lense-Thirring force is swam ped by the drag, but also since the empirical cross-track accelerations (larger than the out-ofplane ones) tted by K onopliv et al. (2006) have just such a magnitude. It turns out that

$$h N_{drag}i_{T} = 1 = 10^{-5} m;$$
 (5)

a value far too sm all to be detected by the MGSRMS overlap di erences which, instead, are capable to appreciate a minimum of 10^{-2}

m (in the radial direction). Moreover, Konopliv et al. (2006) did model the atm ospheric drag. Last but not least, the along-track and radial portions of the MGS orbit are primarily a ected by the nongravitational forces, as clearly stated in (Lem oine et al. 2001; Konopliv et al. 2006); they exhibit tim e-varying patterns over 12 hr (Lem oine et al. 2001) which, m apped to the out-of-plane direction, are averaged out over our multi-year time span P.

A cknow ledgm ents

I gratefully thank A .K on pliv, NASA Jet P ropulsion Laboratory (JPL), for the explanation about the orientation adopted for the MGS norm all direction.

References

- [1] Christodoulidis D C, Sm ith D E, W illiam son R G and K losko S M 1988 J.Geophys. Res. 93 (B6) 6216
- [2] Iorio L 2007 gr-qc/0701042
- [3] Konopliv A S et al 2006 Icarus 182 23
- [4] K rogh K 2007 astro-ph/0701653
- [5] Lem oine F G et al 2001 J Geophys Res 106 23359
- [6] MilaniA, NobiliA M, Farinella P 1987 Non-gravitational perturbations and satellite geodesy (A dam Hilger, Bristol)
- [7] Tapley B D, Schutz B E, Born G H 2004 Statistical O rbit D eterm ination (E lsevier A cadem ic P ress)