

# MODELLING THE DECOHERENCE OF SPACETIME

J.T. WHELAN

*Department of Physics, 201 James Fletcher Building  
University of Utah, Salt Lake City, UT 84112*

The question of whether unobserved short-wavelength modes of the gravitational field can induce decoherence in the long-wavelength modes (“the decoherence of spacetime”) is addressed using a scalar field toy model with some features of perturbative general relativity. For some long-wavelength coarse grainings, the Feynman-Vernon influence phase is found to be effective at suppressing the off-diagonal elements of the decoherence functional. The requirement that the short-wavelength modes be in a sufficiently high-temperature state places limits on the applicability of this perturbative approach.

## 1 Environment-induced decoherence in generalized QM

To review briefly, the sum-over-histories implementation of generalized quantum mechanics, as formulated by Hartle<sup>1</sup>, assigns to each pair of field histories  $\{\varphi, \varphi'\}$  a *fine-grained decoherence functional*

$$D[\varphi, \varphi'] = \rho(\varphi_i, \varphi'_i) \delta(\varphi'_f - \varphi_f) e^{i(S[\varphi] - S[\varphi'])} \quad (1)$$

(in units where  $\hbar = 1 = c$ ), where  $\varphi_i = \varphi(t_i)$  and  $\varphi_f = \varphi(t_f)$  are the endpoints of the path  $\varphi$ , and similarly for  $\varphi'$ . One can *coarse grain* by partitioning the field histories into classes  $\{c_\alpha\}$  and summing the fine-grained decoherence functional (1) over all field histories lying in a given class:

$$D(\alpha, \alpha') = \int_{\alpha} \mathcal{D}\varphi \int_{\alpha'} \mathcal{D}\varphi' D[\varphi, \varphi']. \quad (2)$$

If this coarse-grained decoherence functional (which is a Hermitian matrix with indices  $\alpha$  and  $\alpha'$ ) is diagonal [ $D(\alpha, \alpha') \approx \delta_{\alpha\alpha'} p_\alpha$ ], then each diagonal element  $p_\alpha$  is the probability that the system follows a history in class  $c_\alpha$ . We say then that the coarse graining  $\{c_\alpha\}$  *decoheres*. When the alternatives do not decohere, quantum mechanical interference prevents the theory from assigning consistent probabilities (as defined by the classical probability sum rules).

Most decoherence in physical situations arises<sup>2</sup> when the variables  $\varphi$  can be split into “system” variables  $\Phi$  which can define coarse grainings of interest and “environment”  $\phi$  which is only relevant due to its interaction with the system. The action can then be divided:  $S[\varphi] = S_\Phi[\Phi] + S_E[\phi, \Phi]$ , into a piece  $S_\Phi$  which would describe the system in the absence of the environment and a

piece  $S_E$  which describes the environment and its coupling to the system. The decoherence functional  $D(\alpha, \alpha')$  for a coarse graining based solely on the system variables can be written in terms of  $D[\Phi, \Phi'] = \int \mathcal{D}\phi \mathcal{D}\phi' D[\varphi, \varphi']$ . Assuming that the density matrix describing the initial state factors:

$$\rho(\varphi_i, \varphi'_i) = \rho_\Phi(\Phi_i, \Phi'_i) \rho_\phi(\phi_i, \phi'_i), \quad (3)$$

then  $D[\Phi, \Phi']$  can be written

$$D[\Phi, \Phi'] = \rho_\Phi(\Phi_i, \Phi'_i) \delta(\Phi'_f - \Phi_f) e^{i(S_\Phi[\Phi] - S_\Phi[\Phi'] + W[\Phi, \Phi'])} \quad (4)$$

where

$$e^{iW[\Phi, \Phi']} = \int \mathcal{D}\phi \mathcal{D}\phi' \rho_\phi(\phi_i, \phi'_i) \delta(\phi'_f - \phi_f) e^{i(S_E[\phi, \Phi] - S_E[\phi', \Phi'])}. \quad (5)$$

$W[\Phi, \Phi']$  is called the Feynman-Vernon influence phase<sup>3</sup>; if the *influence functional*  $e^{iW}$  becomes small for  $\Phi \neq \Phi'$ , the “off-diagonal” parts of  $D[\Phi, \Phi']$  will be suppressed, causing alternatives defined in terms of  $\Phi$  to decohere.<sup>2</sup>

## 2 Toy Model

The model which we use to mimic general relativity is that of a scalar field  $\varphi$  on Minkowski space with action

$$S[\varphi] = -\frac{1}{2} \int d^4x (\partial_\mu \varphi)(\partial^\mu \varphi) [1 - (2\pi)^{3/2} \ell \varphi]. \quad (6)$$

This action has the same structure as the first few terms of a perturbative expansion of GR: a wave term followed by an interaction term containing two derivatives. In fact it can be obtained<sup>4</sup> from the Nordström-Einstein-Fokker theory<sup>5</sup> theory of a conformally flat metric  $g_{\mu\nu} = \Omega^2 \eta_{\mu\nu}$  by taking  $\varphi$  proportional to  $\Omega^4 - 1$  (motivated by the volume element  $\sqrt{|g|} = \Omega^4$ ) and  $\ell$  proportional to the Planck length, and expanding the action to the first interacting order in  $\ell$ .

## 3 On the applicability of perturbation theory

As described in section 1, decoherence is enforced if  $|e^{iW[\Phi, \Phi']}| \ll 1$  whenever  $\Phi - \Phi'$  is appreciable. However, since to zeroth order in perturbation theory, the system and environment are not coupled, a perturbative expansion will yield  $|e^{iW}| = 1 + \mathcal{O}(\ell)$ . In order to make  $|e^{iW}|$  small, the  $\mathcal{O}(\ell)$  term must be

comparable to unity, making this sort decoherence in some sense an inherently nonperturbative phenomenon.

It is still possible, however, to treat some aspects of the problem perturbatively if there is another small quantity  $\beta$ . In that case, while terms proportional to  $\ell$  can be small and be treated perturbatively, terms proportional to  $\frac{\ell}{\beta}$  can be large, producing the seemingly nonperturbative effect. In the present work, the small parameter is the inverse temperature of the thermal state assumed to describe the short-wavelength environment.

#### 4 Results

If the wavenumber  $k_c$  dividing the short- and long-wavelength regimes and the inverse temperature  $\beta$  of the environment obey  $\beta k_c \ll 1$ , one can calculate<sup>4</sup> the upper limit

$$e^{iW[\Phi, \Phi']} \leq \frac{1}{(1 + A^2[\Delta\Phi])^{1/4}}, \quad (7)$$

which is indeed small for large values of the suppression factor

$$A^2 \gtrsim \frac{\ell^2}{\beta^2} \int_{q < k_c} d^3q \int_{-q}^q d\omega |\Delta\Phi_{\mathbf{q}\omega}|^2 \frac{(2\pi)^2 (q^2 - \omega^2)^2}{32q|\omega|} \ln \left( 1 + \frac{|\omega|}{k_c} \right). \quad (8)$$

Considering, for example, a coarse graining by values of a “field average”  $\langle\Phi\rangle = \Delta\omega(\Delta q)^3 \Phi_{\mathbf{q}_0\omega_0}$ , the suppression factor is bounded (when  $q_0 \gg \omega_0$ ) by

$$A^2 \gtrsim \frac{\pi^2}{8} \frac{|\langle\ell\Delta\Phi\rangle|^2}{\beta^2 q_0^2} \frac{q_0^4}{\Delta\omega(\Delta q)^3}. \quad (9)$$

Since the factors  $\frac{1}{\beta q_0}$ ,  $\frac{q_0}{\Delta\omega}$ , and  $\frac{q_0}{\Delta q}$  are all large,  $A^2$  can become large and suppress the off-diagonal components of  $D[\Phi, \Phi']$  even when the difference  $\ell\Delta\Phi = \ell\Phi - \ell\Phi'$  is still small enough to be within the perturbative regime.

1. See, for example, J.B. Hartle, in *Gravitation and Quantizations*, proceedings of the 1992 Les Houches Summer School, ed. B. Julia and J. Zinn-Justin (North Holland, Amsterdam, 1995).
2. M. Gell-Mann and J.B. Hartle, *Phys. Rev. D* **47**, 3345 (1993).
3. R.P. Feynman and J.R. Vernon, *Ann. Phys. (N.Y.)* **24**, 118 (1963).
4. J.T. Whelan, preprint gr-qc/9612028
5. G. Nordström, *Phys. Z.* **13**, 1126 (1912), *Ann. d. Phys.* **40**, 856 (1913), **42**, 533 (1913), **43**, 1101 (1914); A. Einstein and A.D. Fokker, *Ann. d. Phys.* **44**, 321 (1914).