Cosmological perturbation with two scalar fields in reheating after inflation

A.Taruya [∗] and Y.Nambu †

Department of Physics, Nagoya University, Chikusa-ku, Nagoya 464-01, Japan

Abstract

We investigate the cosmological perturbation of two-scalar field model during the reheating phase after inflation. Using the exact solution of the perturbation in long-wavelength limit, which is expressed in terms of the background quantities, we analyze the behavior of the metric perturbation. The oscillating inflaton field gives rise to the parametric resonance of the massless scalar field and this leads to the amplification of the iso-curvature mode of the metric perturbations.

PACS: 98.80.-k; 98.80.Bp

Keywords: Cosmological perturbation; Reheating; Scalar fields; Parametric amplification

[∗] e-mail: ataruya@allegro.phys.nagoya-u.ac.jp

[†] e-mail: nambu@allegro.phys.nagoya-u.ac.jp

I. INTRODUCTION

Recently, a theory of reheating after inflation is developed and importance of the oscillating scalar fields is recognized [\[1](#page-11-0),[2](#page-11-0),[3,4\]](#page-11-0). The non-linear interaction between the scalar fields amplifies the fluctuations of the scalar fields by the effect of parametric resonance and the energy of the oscillating field is transfered to the fluctuation of the massless field by catastrophic particle production. A large amount of studies about the reheating process has been done [\[5\]](#page-11-0). From the viewpoint of the structure formation of our universe, one of the most important question is whether the growth of fluctuations of the matter field during reheating affects the large-scale inhomogeneities in the universe or not. To answer this question, we must study the evolution of the metric and the matter fluctuation in general relativistic treatment.

Concerning the works on the theory of cosmological perturbation, several authors studied a single scalar field model in the context of the reheating scenario and investigated a role of the coherent oscillating field on the evolution of large-scale structure in the universe[[6,7,8\]](#page-11-0). Hamazaki and Kodama analyze the metric perturbation in the model with two-component fluid and discuss the effect of parametric resonance[[9](#page-11-0)]. They evaluate the curvature perturbation ζ (Bardeen parameter) by replacing the scalar fields with the perfect fluids. The effect of parametric resonance is all included in the energy transfer term which should be given by phenomenological description. They conclude that ζ is well-conserved during reheating and the parametric resonant decay does not affect the scenario of the structure formation. Although their conclusion seems to be valid in the old version of the reheating scenario, which is dominated by the Born decay process, it is not clear that their analysis is appropriate when the non-linear evolution of the scalar fields may play an important role. We should keep in mind that the parametric resonance does not occur in the perfect fluid system.

In this letter, we analyze the cosmological perturbation of the scalar field model to clarify whether the dynamics of coherently oscillating scalar field affects the metric perturbation or not. The model we treat here contains two scalar fields ϕ , χ with the potential $V(\phi, \chi) =$ $m^2\phi^2/2 + g^2\phi^2\chi^2/2$ where ϕ is the inflaton and χ is the massless boson field.

II. BASIC EQUATIONS

We consider a homogeneous, isotropic and flat cosmological model with scalar type metric perturbations. In longitudinal (Newtonian) gauge, the metric is given by

$$
ds^{2} = -(1+2\Phi)dt^{2} + a^{2}(t)(1-2\Psi)\delta_{ij}dx^{i}dx^{j}.
$$
\n(1)

 Φ , Ψ correspond to the gauge-invariant potential, which characterize the physical mode of the metric perturbations. The Bardeen parameter which represents the spatial curvature perturbation on the uniform energy density slice in super-horizon scale is written by $[10,11,12,13]$

$$
\zeta = \Phi - \frac{H^2}{\dot{H}} (\Phi + H^{-1} \dot{\Phi}) ; \quad H \equiv \frac{\dot{a}}{a}.
$$
 (2)

For the later analyses, we use the following dimensionless variables:

$$
\tau = mt, \quad h = \frac{H}{m}, \quad \varphi_1 = \sqrt{\frac{4\pi}{3}} \frac{\phi}{m_{pl}}, \quad \varphi_2 = \sqrt{\frac{4\pi}{3}} \frac{\chi}{m_{pl}}, \quad \lambda = \frac{3}{4\pi} \left(\frac{gm_{pl}}{m}\right)^2. \tag{3}
$$

In terms of these variables, the background Einstein equations are given by

$$
h^{2} = \varphi_{1}^{\prime 2} + \varphi_{2}^{\prime 2} + 2U(\varphi_{1}, \varphi_{2}); \quad U = \frac{1}{2}\varphi_{1}^{2} + \frac{1}{2}\lambda\varphi_{1}^{2}\varphi_{2}^{2}, \tag{4}
$$

$$
\varphi_i'' + 3h\varphi_i' + \frac{\partial U}{\partial \varphi_i} = 0 \quad (i = 1, 2),\tag{5}
$$

where $(') = d/d\tau$. We use the following gauge-invariant variable to describe the perturbation:

$$
Q_i = \delta \varphi_i + \frac{\varphi_i'}{h} \Psi,\tag{6}
$$

where $\delta\varphi_i$ is perturbation of the scalar field φ_i . The variable Q_i was first introduced by Mukhanov [\[14\]](#page-11-0) and all the other gauge-invariant quantities can be expressed by these variables. Using the fact $\Phi = \Psi$ in the present model, the Bardeen parameter defined by (2) is written by

$$
\zeta = \frac{h}{\varphi_1'^2 + \varphi_2'^2} (\varphi_1' Q_1 + \varphi_2' Q_2). \tag{7}
$$

The evolution equation for Q_i is simply expressed by [\[15](#page-11-0)]

$$
Q''_i + 3hQ'_i + \left(\frac{k}{ma}\right)^2 Q_i + \sum_{j=1}^2 \left[\frac{\partial^2 U}{\partial \varphi_i \partial \varphi_j} - \frac{6}{a^3} \left(\frac{a^3}{h} \varphi'_i \varphi'_j\right)' \right] Q_j = 0.
$$
 (8)

This is our basic equation.

III. SOLUTION IN LONG WAVELENGTH LIMIT

As we are interested in the formation of the large-scale structure in our universe, it is important to investigate the metric fluctuation outside the scale of the Hubble horizon. Fortunately, we can obtain the exact solution of eq.(8) in the long wavelength limit $(k \to 0)$ and the result is expressed in terms of the background quantities.

Let us suppose the solution of the background field is written as $\varphi_i(\alpha, C)$, where $\alpha \equiv \log a$ and C is the parameter given by a suitable choice of the integration constants. α plays a role of the time parameter and we can trace the time evolution of the trajectory of the background solution in configuration space (φ_1, φ_2) using this parameter. C distinguishes trajectories in this space and remains constant on a each trajectory. Using the time parameter α , the field equation [\(5](#page-2-0)) is rewritten as

$$
\frac{d^2\varphi_i}{d\alpha^2} + \left(3 + h^{-1}\frac{dh}{d\alpha}\right)\frac{d\varphi_i}{d\alpha} + h^{-2}\frac{\partial U}{\partial \varphi_i} = 0.
$$
\n(9)

Differentiating (9) with respect to α , C, we can show that the equation of motion for the tangent vectors $d\varphi_i/d\alpha$, $d\varphi_i/dC$ becomes (8) in the limit $k \to 0$. Therefore these vectors are the independent solutions of (8) in long wavelength limit[[19\]](#page-11-0).

Since (8) are the second order coupled differential equations, we have four independent solutions. The remaining two solutions can be obtained as follows: define the component of a 2×2 matrix X as

$$
\mathbf{X}_{11} = \frac{d\varphi_1}{d\alpha}, \quad \mathbf{X}_{12} = \frac{d\varphi_1}{dC}, \quad \mathbf{X}_{21} = \frac{d\varphi_2}{d\alpha}, \quad \mathbf{X}_{22} = \frac{d\varphi_2}{dC}.
$$
 (10)

Eachrow vectors in X satisfy ([8\)](#page-3-0). Expressing the remaining solutions by the matrix Y and putting $\boldsymbol{Y} = \boldsymbol{X} \cdot \boldsymbol{P}$ yields

$$
P'' + (3h + 2X^{-1}X')P' = O.
$$
 (11)

We can obtain $\bm{P}' = \bm{X}^{-1} (\bm{X}^{-1})^T/a^3$ by using the fact that $\bm{X}' \cdot \bm{X}^{-1}$ is the symmetric matrix [\[19\]](#page-11-0). Thus the general solution of([8\)](#page-3-0) in long wavelength limit is given by

$$
Q_i = \sum_{j=1}^{2} \left(c_j^+ \mathbf{X}_{ij} + c_j^- \left[\mathbf{X} \int \frac{d\tau}{a^3} \mathbf{X}^{-1} (\mathbf{X}^{-1})^T \right]_{ij} \right), \tag{12}
$$

where (T) represents transpose of the matrix and $c_j^{\pm}(j=1,2)$ are arbitrary constants.

IV. BACKGROUND DYNAMICS IN REHEATING PHASE

We can know the behavior of the long wavelength perturbation from (12) if we obtainthe background solution. As we pay our attention to the reheating phase of the inflaton dominated universe, we assume the condition $\varphi_1, \varphi_2 \lesssim 1$ and $\lambda \varphi_2^2 \lesssim 1$. The former is the condition for the oscillation of the scalar fields. The latter condition implies that the potential energy is dominated by the massive inflaton.

We first solve the background equations in a naive treatment. Because $\lambda \varphi_2^2 \lesssim 1$, the evolutionequations $(4)(5)$ $(4)(5)$ $(4)(5)$ $(4)(5)$ can be solved separately and we get oscillatory behavior of the inflaton $\varphi_1 \simeq \bar{\varphi}(\tau) \cos \tau$, where $\bar{\varphi} \propto 1/\tau$. Substituting this into ([5\)](#page-2-0) again, we find that the equation for the field φ_2 becomes the same form as the Mathieu equation:

$$
(a^{3/2}\varphi_2)'' + [A + 2q\cos(2\tau)](a^{3/2}\varphi_2) = 0 \; ; \; a \propto \tau^{2/3}, \tag{13}
$$

where the coefficients A and q are time dependent functions given by $A = 2q = \lambda \bar{\varphi}^2(\tau)/2$. The stability/instability chart of the Mathieu equation says that φ_2 has unstable behavior at $1/3 \le q \le 1$, in the region of the first resonance band of the Mathieu function [[19](#page-11-0)].

To investigate the metric perturbation during the reheating, we must study the background evolution more precisely. We shall analyze the background dynamics by using the

renormalization group (RG) *method.* RG method is a technique of asymptotic analysis which improves the result of the naive perturbation and it provides an unified approach to solve differential equations including singular perturbation method [\[16](#page-11-0)] [\[17](#page-11-0)].

To apply the RG method, we identify the small expansion parameter ϵ as the amplitude of the scalar fields $\varphi_i \sim \mathcal{O}(\epsilon) < 1$. Since we want to take into account of the non-linear interaction, we assume $\tilde{\lambda} \equiv \epsilon \lambda \sim \mathcal{O}(1)$ which implies λ is not small parameter. Then the background quantities are expanded as follows:

$$
\varphi_i = \epsilon \varphi_{i(1)} + \epsilon^2 \varphi_{i(2)} + \cdots, \quad h = \epsilon h_{(1)} + \epsilon^2 h_{(2)} + \cdots.
$$
 (14)

The crucial treatment in the present model is to insert the mass term $\omega^2 \varphi_2$ in the equation of φ_2 [\(5](#page-2-0)). We put $\omega^2 = 1 + \epsilon \sigma$ and take the limit $\epsilon \sigma \to -1$ after the end of calculation. This treatment is necessary to extract the effect of parametric resonance in φ_2 field.

In appendix, we apply the RG method to the background equations $(4)(5)$ $(4)(5)$ $(4)(5)$. The final form of the solution up to $\mathcal{O}(\epsilon)$ becomes

$$
\varphi_1 \simeq h \cos \Theta \cos (\tau + \theta), \quad \varphi_2 \simeq h \sin \Theta \cos (\tau + \psi), \tag{15}
$$

where we replace $\epsilon h_{(1)}$ with h. The variables Θ , θ , ψ and h are time dependent and determinedby so-called RG equations $(A4)-(A7)$ $(A4)-(A7)$ $(A4)-(A7)$ $(A4)-(A7)$ which are the result of the renormalization of the initial amplitude.

We further impose the condition $\varphi_2 \ll \varphi_1 \ (\Theta \ll 1)$, which makes our analysis easier. We will restrict our analysis in the case of $\Theta \ll 1$ hereafter. [\(A6\)](#page-10-0) becomes $\theta' \simeq 0$ and we can set $\theta = 0$ without loss of generality. Eqs.([A4](#page-10-0))[\(A5](#page-10-0))[\(A7\)](#page-10-0) become

$$
\Theta' \simeq \frac{1}{2} q(\tau) \sin \gamma \cdot \Theta, \quad \gamma' \simeq -1 + q(\tau) (2 + \cos \gamma), \tag{16}
$$

$$
h' = -\frac{3}{2}h^2,\tag{17}
$$

where we define the time dependent parameter

$$
q(\tau) = \frac{1}{4}\lambda h^2. \tag{18}
$$

Since [\(17](#page-5-0)) yields $h \propto 1/\tau$, $q(\tau)$ is the decreasing function of time. Defining the new variables u, v by $u = \Theta \cos(\gamma/2)$, $v = \Theta \sin(\gamma/2)$, [\(16](#page-5-0)) is rewritten as follows:

$$
\begin{pmatrix} u' \\ v' \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 0 & 1-q \\ 3q-1 & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.
$$
 (19)

The eigenvalue of the matrix in the right hand side of (19) is real for $1/3 \le q \le 1$ and Θ can grow during this epoch. We see that the function q corresponds to the coefficient q in the Mathieu equation [\(13](#page-4-0)). Recall that we have $\varphi_1 \simeq h \cos \tau$, $\varphi_2 \simeq h \Theta \cos (\tau + \gamma/2)$ from [\(15](#page-5-0)), we recognize that the amplitude of φ_2 has the growing behavior as the result of the parametric resonance. Soon after q becomes smaller than $1/3$, Θ reaches a constant value. Fig.1 shows the evolution of the background fields obtained by numerical integration of [\(4](#page-2-0))[\(5](#page-2-0)). The field φ_2 (the solid line) is amplified during the short period denoted by the horizontal arrow, which corresponds to the resonance epoch $1/3 \le q \le 1$. We can evaluate howmuch φ_2 is amplified through the resonance epoch. From ([16](#page-5-0)), we have

$$
\Theta \simeq \Theta_0 \exp\left[\frac{1}{2} \int_{\tau_i}^{\tau} d\tau' q \sin \gamma\right].
$$
\n(20)

Using this expression, we get the upper bound

$$
\frac{\Theta(\tau_f)}{\Theta(\tau_i)} \lesssim \exp\left(c_*\sqrt{\lambda}\right); \quad c_* = \frac{1}{6} \left(1 - \frac{1}{\sqrt{3}}\right) \simeq 0.07,\tag{21}
$$

where we used the relation $q(\tau_f) = 1/3$, $q(\tau_i) = 1$. The above expression is valid if the condition $\Theta \ll 1$ is satisfied throughout the resonance epoch.

V. EVOLUTION OF THE BARDEEN PARAMETER

Behavior of the background solution obtained by RG method is translated to the metric perturbation ζ using the long wavelength solution [\(12](#page-4-0)). The reduced system ([16\)](#page-5-0) has the two integration constants: The initial values of γ , Θ . During the resonance epoch, γ approaches a constant value which does not depend on the initial value. We can regard Θ_0 (the initial value of Θ) as the parameter C described in Sec[.III.](#page-3-0) That is, Θ_0 is the parameter which distinguishes the trajectories in the reduced system (Θ, γ) . Substituting [\(15](#page-5-0)) into [\(12\)](#page-4-0) and evaluating the integral adiabatically, we find that the dominant contribution to the Bardeen parameter ζ comes from the solution $Q_i^{ad} \equiv d\varphi_i/d\alpha$ and

$$
Q_1^{iso} \equiv \frac{d\varphi_1}{dC} \simeq -h\Theta_{,C} \sin\Theta \cos\tau, \quad Q_2^{iso} \equiv \frac{d\varphi_2}{dC} \simeq h\Theta_{,C} \cos\Theta \cos(\tau + \frac{\gamma}{2}),\tag{22}
$$

where $\Theta_{,C} = \exp \left[\int d\tau' q/2 \cdot \sin \gamma \right]$. The other independent solutions are recognized as the decaying modes [\[19](#page-11-0)].

The solution Q_i^{ad} is referred to as the adiabatic growing mode [\[18\]](#page-11-0). Substituting Q_i^{ad} into [\(7](#page-3-0)), the Bardeen parameter remains constant in time. The constancy of ζ arises in the hydrodynamical perturbation without the entropy fluctuation [\[12](#page-11-0)]. For the solution Q_i^{iso} , ([7](#page-3-0)) gives

$$
\zeta^{iso} \simeq h\Theta_{,C} \sin \Theta \cos \Theta \cdot \frac{\sin \tau \cos \tau - \sin \left(\tau + \frac{\gamma}{2}\right) \cos \left(\tau + \frac{\gamma}{2}\right)}{\cos^2 \Theta \sin^2 \tau + \sin^2 \Theta \sin^2 \left(\tau + \frac{\gamma}{2}\right)}.
$$
\n(23)

We see that ζ^{iso} has the periodically sharp peaks around the zero points of φ_1' . This behavior also appears in the single field case [\[7](#page-11-0)]. Except for the short interval of the peaks, ζ^{iso} traces the amplitude of Q_1^{iso} . Therefore ζ^{iso} deviates from zero when $\Theta_{,C}$ grows due to the parametric resonance. Note that ζ^{iso} vanishes when the background field φ_2 is zero $(\Theta = 0)$. Because the initial amplitude of the curvature perturbation ζ^{iso} is vanishingly small, we can identify Q_i^{iso} is the contribution of the iso-curvature mode of the perturbation.

Weconfirm the behavior of ζ^{iso} by solving $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ numerically. In Fig.2, we show the time evolution of the iso-curvature mode. In contrast to the constancy of the adiabatic growing mode (the dashed line), we observe that ζ^{iso} (the solid line) is amplified during the resonance epoch when the amplitude of the background φ_2 field grows (see Fig.1). After the resonance epoch, ζ^{iso} approaches zero due to the Hubble damping which can be seen by the analyticresult (23). Using ([21\)](#page-6-0)(23), we can estimate the growth of ζ^{iso} except for the sharp peak:

$$
\left| \frac{\zeta^{iso}(\tau_f)}{\zeta^{iso}(\tau_i)} \right| \lesssim \frac{h(\tau_f)}{h(\tau_i)} \cdot \exp\left(2c_*\sqrt{\lambda}\right). \tag{24}
$$

VI. SUMMARY AND DISCUSSIONS

In this letter, we have investigated the cosmological perturbation of the model with two scalar fields in reheating phase after inflation. Using the background quantities, long wavelength solutions of the perturbation are obtained. Applying the RG method to the background dynamics in the coherent oscillating stage, we found that the massless field gets the effect of the parametric resonance. This leads to the amplification of the isocurvature metric perturbations. The result implies that the Bardeen parameter deviates from the initial amplitude in reheating phase even though the initial amplitude determined in the inflationary stage is dominated by the adiabatic mode. Therefore the analysis using the scalar field model is essential to study the metric perturbations when the non-linear dynamics of the scalar fields plays an important role.

The most important observed quantity is the power spectrum $\mathcal{P}_{\zeta}(k)$. Consider the evolution of the metric fluctuation $\zeta(k)$ whose initial amplitude is given by the quantum fluctuation of the scalar fields during inflation. As long as the wavelength of the fluctuation is outside the Hubble horizon, time evolution of $\zeta(k)$ is well-approximated by the solution for the long wavelength mode. The power spectrum for the present length-scale can be evaluated from the amplitude of $\zeta(k)$ when the fluctuations re-enter the Hubble horizon. Soon after the metric fluctuation $\zeta(k)$ suffers the parametric amplification during reheating, it approaches to the initial amplitude due to the Hubble damping. The horizon re-entry time becomes earlier as the wavelength of the fluctuation becomes shorter. Therefore, we can expect that the effect of parametric resonance on the power spectrum at the horizon re-entry appears at the small scale (large k). We will give a detail discussion about evolution of the power spectrum in the forthcoming paper [\[19\]](#page-11-0).

ACKNOWLEDGMENT

This work is partly supported by the Grand-In-Aid for Scientific Research of the Ministry of Education, Science, Sports and Culture of Japan (09740196).

APPENDIX: RG METHOD APPLYING TO THE BACKGROUND SYSTEM

In this appendix, we apply the RG method to the background system $(4)(5)$ $(4)(5)$ $(4)(5)$ $(4)(5)$. Following the treatment described in Sec.[IV,](#page-4-0)we substitute ([14\)](#page-5-0) to ([4\)](#page-2-0)([5\)](#page-2-0) and expand in powers of ϵ :

$$
\mathcal{O}(\epsilon^1): \varphi''_{i(1)} + \varphi_{i(1)} = 0, \quad (i = 1, 2)
$$

\n
$$
h_{(1)} = \sqrt{\varphi'^2_{1(1)} + \varphi^2_{2(1)} + \varphi^2_{2(1)}},
$$

\n
$$
\mathcal{O}(\epsilon^2): \varphi''_{1(2)} + \varphi_{1(2)} = -3h_{(1)}\varphi'_{1(1)} - \tilde{\lambda}\varphi^2_{2(1)}\varphi_{1(1)},
$$

\n
$$
\varphi''_{2(2)} + \varphi_{2(2)} = -3h_{(1)}\varphi'_{2(1)} - \sigma\varphi_{2(1)} - \tilde{\lambda}\varphi^2_{1(1)}\varphi_{2(1)}.
$$

We first solve the above equations order by order. The solutions up to $\mathcal{O}(\epsilon^2)$ become

$$
\varphi_1 = \epsilon \left[A_0 + \epsilon (\tau - \tau_0) \left\{ -\frac{3}{2} h_{(1)} A_0 + i \frac{\tilde{\lambda}}{2} \left(2|B_0|^2 A_0 + B_0^2 A_0^* \right) \right\} \right] e^{i\tau}
$$

$$
+ \epsilon^2 \frac{\tilde{\lambda}}{8} B_0^2 A_0 e^{i3\tau} + c.c. + \mathcal{O}(\epsilon^3)
$$

and

$$
\varphi_2 = \epsilon \left[B_0 + \epsilon (\tau - \tau_0) \left\{ -\frac{3}{2} h_{(1)} B_0 + i \frac{\tilde{\lambda}}{2} \left(2 |A_0|^2 B_0 + A_0^2 B_0^* \right) + \frac{i}{2} \sigma B_0 \right\} \right] e^{i\tau}
$$

$$
+ \epsilon^2 \frac{\tilde{\lambda}}{8} A_0^2 B_0 e^{i3\tau} + c.c. + \mathcal{O}(\epsilon^3),
$$

where *c.c.* means complex conjugate and $h_{(1)}$ is given by

$$
h_{(1)} = 2\sqrt{|A_0|^2 + |B_0|^2}.\tag{A1}
$$

 A_0 , B_0 are the amplitudes at the initial time $\tau = \tau_0$. The above solutions are correct within $\epsilon(\tau - \tau_0) \lesssim 1$ because the secular terms which grow as $(\tau - \tau_0)$ makes the perturbative

expansion break down. To improve the perturbation series, we introduce an arbitrary parameter μ . Splitting $\tau - \tau_0$ as $\tau - \mu + \mu - \tau_0$ and absorbing the terms containing $\mu - \tau_0$ into the renormalization constant Z_A , Z_B defined by $A_0 = A(\mu)Z_A(\tau_0, \mu)$, $B_0 = B(\mu)Z_B(\tau_0, \mu)$. This procedure is always possible for each order of expansion and we have the μ dependent solutions:

$$
\varphi_1 = \epsilon \left[A + \epsilon(\tau - \mu) \left\{ -\frac{3}{2} h_{(1)} A + i \frac{\tilde{\lambda}}{2} \left(2|B|^2 A + B^2 A^* \right) \right\} \right] e^{i\tau} \n+ \epsilon^2 \frac{\tilde{\lambda}}{8} B^2 A e^{i3\tau} + c.c. + \mathcal{O}(\epsilon^3),
$$
\n(A2)\n
$$
\varphi_2 = \epsilon \left[B + \epsilon(\tau - \mu) \left\{ -\frac{3}{2} h_{(1)} B + i \frac{\tilde{\lambda}}{2} \left(2|A|^2 B + A^2 B^* \right) + \frac{i}{2} \sigma B \right\} \right] e^{i\tau} \n+ \epsilon^2 \frac{\tilde{\lambda}}{8} A^2 B e^{i3\tau} + c.c. + \mathcal{O}(\epsilon^3).
$$
\n(A3)

Since μ does not appear in the original equations, the solution should not depend on μ . This requires $(\partial \varphi_i/\partial \mu)_\tau = 0$ for arbitrary τ and we obtain the RG equations:

$$
\begin{split} \frac{dA}{d\mu}&=\epsilon\left[-\frac{3}{2}h_{(1)}A+i\frac{\tilde{\lambda}}{2}\left(2|B|^{2}A+B^{2}A^{*}\right)\right]+\mathcal{O}(\epsilon^{2}),\\ \frac{dB}{d\mu}&=\epsilon\left[-\frac{3}{2}h_{(1)}B+\frac{i}{2}\sigma B+i\frac{\tilde{\lambda}}{2}\left(2|A|^{2}B+A^{2}B^{*}\right)\right]+\mathcal{O}(\epsilon^{2}). \end{split}
$$

By setting $\mu \equiv \tau$ in (A2)(A3), the secular terms are eliminated and we obtain the regular perturbation. Rewriting the variables A , B by

$$
A = \frac{1}{2}h_{(1)}\cos\Theta e^{i\theta}, \quad B = \frac{1}{2}h_{(1)}\sin\Theta e^{i\psi},
$$

and using([A1\)](#page-9-0), we get the solution [\(15\)](#page-5-0) and the above RG equations reduce to

$$
\Theta' = \epsilon \frac{\tilde{\lambda}}{16} h_{(1)}^2 \sin \gamma \sin 2\Theta,
$$
\n(A4)

$$
\gamma' = -1 + \epsilon \frac{\tilde{\lambda}}{4} h_{(1)}^2 \cos 2\Theta (2 + \cos \gamma), \tag{A5}
$$

$$
\theta' = \epsilon \frac{\tilde{\lambda}}{8} h_{(1)}^2 \cos \gamma \sin^2 \Theta, \qquad \gamma = 2(\psi - \theta), \tag{A6}
$$

$$
h'_{(1)} = -\epsilon \frac{3}{2} h_{(1)}^2,\tag{A7}
$$

where we set $\epsilon \sigma = -1$.

REFERENCES

- [1] A.Dolgov and D.Kirilova, Sov.J.Nucl.Phys.51 (1990) 172.
- [2] J.Traschen and R.Brandenberger, Phys.Rev.D 42 (1990) 2491.
- [3] L.Kofman, A.Linde and A.A.Starobinsky, Phys.Rev.Lett.73 (1994) 3195.
- [4] Y.Shtanov, J.Traschen and R.Brandenberger, Phys.Rev.D51 (1995) 5438.
- [5] e.g, L.Kofman, A.Linde and A.A.Starobinsky, Phys.Rev.D 56 (1997) 3258 and references there in.
- [6] Y.Nambu and A.Taruya, Prog.Theor.Phys.97 (1997) 83.
- [7] H.Kodama and T.Hamazaki, Prog.Theor.Phys.96 (1996) 949.
- [8] J.Hwang, Phys.Lett.B401 (1997) 241.
- [9] T.Hamazaki and H.Kodama, Prog.Theor.Phys.96 (1996) 1123.
- [10] D.Lyth, Phys.Rev.D **31** (1985) 1792.
- [11] D.S.Salopek, J.R.Bond and J.M.Bardeen, Phys.Rev.D 40 (1989) 1753.
- [12] V.F.Mukhanov, H.A.Feldman and R.H.Brandenberger, Phys.Rep. 215 (1992) 203.
- [13] J.G.Bellido and D.Wands, Phys.Rev.D 53 (1996) 5437.
- [14] V.F.Mukhanov, Sov.Phys.JETP 67 (1988) 1297.
- [15] J.Hwang, preprint [gr-qc/ 9608018.](http://arxiv.org/abs/gr-qc/9608018)
- [16] L.Y.Chen, N.Goldenfeld and Y.Oono, Phys.Rev.E 54 (1996) 376.
- [17] T.Kunihiro, Prog. Theor.Phys.94 (1995) 503.
- [18] D.Polarski and A.A.Starobinsky, Nucl.Phys.B385 (1992) 623; Phys.Rev.D 50 (1994) 6123.
- [19] A.Taruya and Y.Nambu, in preparation.

FIGURE CAPTIONS

- **Fig.1** The amplitude of the background fields φ_1 and φ_2 is shown as a function of time τ by solving([4\)](#page-2-0)([5\)](#page-2-0) numerically. The broken and the solid lines represent the inflaton(φ_1) and the massless field(φ_2) respectively. During the period denoted by the horizontal arrow, the field φ_2 is amplified by the effect of parametric resonance. We used the initialvalues which correspond to the expression ([15\)](#page-5-0) at $\tau = 0$ with the parameters $\lambda = 4000, h = 0.035, \Theta = 0.01, \gamma = 0.4, \theta = 0.0.$
- Fig.2The Bardeen parameter ζ^{iso} is plotted as a function of time τ by solving $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ $(4)(5)(8)$ numerically.We calculate $(4)(5)$ $(4)(5)$ $(4)(5)$ $(4)(5)$ with the same initial condition as described in Fig.1. As for the perturbations, we set the initial values corresponding to the expression ([22\)](#page-7-0) at $\tau = 0$. The figure shows that ζ^{iso} (the solid line) is amplified during the resonance epoch denoted by the horizontal arrow and approaches zero due to the Hubble damping. We also plotted the Bardeen parameter for the adiabatic growing mode (the dashed line) obtained by the substitution of Q_i^{ad} into [\(7](#page-3-0)).

Fig.1

Fig.2