

Einstein's equations in Ashtekar's variables constitute a symmetric hyperbolic system.

Mirta S. Iriondo* , Enzo O. Leguizamón †, and Oscar A. Reula ‡

FaMAF, Medina Allende y Haya de la Torre,
Ciudad Universitaria, 5000 Córdoba, Argentina

Fax nr: +54-51-334054

We show that the 3 + 1 vacuum Einstein field equations in Ashtekar's variables constitutes a first order symmetric hyperbolic system for arbitrary but fixed lapse and shift fields, by suitable adding to the system terms proportional to the constraint equations.

I. INTRODUCTION

In the mid-eighties a new representation for Hamiltonian General Relativity was introduced by Ashtekar [1]. The Ashtekar representation maybe understood as resulting from a complex canonical transformation that goes from the ADM gravitational phase space variables to a new set of variables. One of the major benefits of this representation is that the constraints take a simpler form in terms of these variables than in the ADM formulation. Another important aspect of this set of variables, surprisingly not yet exploited and not shared by the standard ADM representation, is that the evolution equation are a first order system of differential equations. Thus it is natural to consider the well-posedness of the classical initial value problem in this context.

Well posedness of Einstein equations in itself is a long ago closed issue. Since the initial data for the equations in Ashtekar's variables is in one to one, and continuous correspondence with the ADM variables, the result follows at once. Rather, the issue we pose here is whether Einstein's equations in Ashtekar's variables conform a good system to evolve arbitrary initial data using approximations schemes, be numerical or analytical. Bad choices of evolution gauges usually lead into spurious coordinate singularities which effectively destroy the smoothness of the map between variables. We are also concerned with consistency, convergence and stability properties of approximation methods which use these variables. These

issues have not been fully solved in the realm of the numerical analysis of non linear partial differential evolution equations, but all powerful results so far obtained make heavy use of the symmetric hyperbolic character of the equations. This is the main reason for checking this property here. Updated references to this problem are in [2], [3], [4], [5], and [6].

As discussed in the conclusions (section IV), one important property of the system we consider is that the lapse-shift pair can be given before hand on spacetime or can be obtained along evolution, either imposing evolution equations and so effectively enlarging the symmetric hyperbolic system, or imposing elliptic equations on each hypersurface $t = const.$

The plan of the paper is as follows. Section II is devoted to mathematical preliminaries where we recall some aspects of Ashtekar's Hamiltonian formulation. In section III we present the main result.

II. THE FIELD EQUATIONS IN ASHTEKAR'S VARIABLES

In this section we shall present the necessary definitions and a brief overview of the Hamiltonian formalism in spinorial variables in order to get Einstein field equations as a system of evolution and constraint equations (we follow [7], [8], and [9]).

Let M be a four dimensional manifold with a smooth metric g_{ab} of signature $(-+++)$. Furthermore, assume that M admits a time function t , a smooth scalar field on M , whose gradient is everywhere timelike and denote the time direction by t^a , which is a smooth vector field on M with affine parameter t , i.e. $t^a \nabla_a t = 1$. Then the one parameter family of surfaces Σ_t , defined by $t = const.$, are three dimensional spacelike surfaces. Finally we shall denote by n^a the future directed unit normal vector field to Σ_t , and the intrinsic and positive definite metric on Σ_t by q_{ab} . By projecting t^a into and orthogonally to Σ_t , we obtain the lapse and the shift fields N and N^a , thus $t^a = Nn^a + N^a$.

We shall also assume that our spacetime admits a spinorial structure (see [10], and references therein); i.e we shall assume that to each point $p \in M$ we can assign a complex two dimensional vector space W equipped with a nondegenerated symplectic form (skew 2-form). The elements of W will be denoted by ξ^A and the corresponding

*Supported by STINT, The Swedish Foundation for International Cooperation in Research and Higher Education.

†Fellow of Se.CyT-UNC.

‡Member of CONICET.

symplectic form by ϵ_{AB} (its inverse will be denoted by ϵ^{AB} , and is such that $\epsilon_{AC}\epsilon^{AB} = \delta_C^B$). The complex conjugate vector space associated with W will be denoted by \overline{W} , its elements by $\overline{\xi}^{A'}$ and the symplectic form by $\epsilon_{A'B'}{}^1$. Since the group that preserves the structure on W is $SL(2, C)$, ξ^A will be called a $SL(2, C)$ spinor at $p \in M$.

To relate tensors to spinors on M , one must fix a metric preserving isomorphism between the four dimensional real vector space $V = \{\xi^{AA'} \in W \otimes \overline{W} : \overline{\xi}^{AA'} = -\xi^{AA'}, g_{AA'BB'} = \epsilon_{AB}\epsilon_{A'B'}\}$ and the tangent space $T_p(M)$ with metric g_{ab} . This isomorphism is called an $SL(2, C)$ soldering form and is denoted by $\sigma^a{}_{AA'}$. Then

$$g^{ab} = \sigma^a{}_{AA'}\sigma^b{}_{BB'}\epsilon^{AB}\epsilon^{A'B'}.$$

The covariant derivative operator on tensors has a unique extension to spinors if we impose the condition

$$\nabla_a \sigma^b{}_{AA'} = 0.$$

In order to relate spinors on these surfaces to their intrinsic geometry, we shall need an additional structure on W which reduces the structure group $SL(2, C)$ to $SU(2)$. This additional structure on W is a positive definite hermitian inner product which is denoted by $G : W \times W \rightarrow C$, or

$$G_{AA'} : W \times \overline{W} \rightarrow C, \quad (\eta^A, \overline{\xi}^{A'}) \mapsto \overline{\xi}^{A'} G_{AA'} \eta^A.$$

In this notation hermiticity appears as $G_{A'A} = \overline{G}_{AA'}$. We will assume that (either ϵ_{AB} or) $G_{AA'}$ is so normalized that

$$\epsilon^{A'B'} G_{A'A} G_{B'B} = \epsilon_{AB}, \quad \text{or} \quad G_{A'A} G^{A'B} = \delta_A^B,$$

In the formalism 3+1 this additional structure is obtained by choosing

$$G_{AA'} := \sqrt{2} i n_{AA'} := \sqrt{2} i n^a \sigma_a{}_{AA'},$$

then as $\overline{\sigma}_a{}^{AA'} = -\sigma_a{}^{AA'}$ and n^a is timelike and future directed, we have the positive definite hermitian inner product that we were looking for. Via this inner product, we identify unprimed $SL(2, C)$ spinors on M with $SU(2)$ spinors on Σ_t . The relationship between the intrinsic geometry with $SU(2)$ spinors is given by:

$$\sigma_a{}^A{}_B := q_a{}^b \sigma_b{}^{AA'} G_{A'B}$$

that is the $SU(2)$ soldering form. Then

$$q_{ab} = -\sigma_a{}^A{}_B \sigma_b{}^B{}_A := -\text{tr}(\sigma_a \sigma_b).$$

Here a matrix notation is employed (and shall be used in the following) for unprimed spinor indices in which adjacent summed indices go from upper left to lower right, e.g., $(\sigma^a \sigma^b)_{A^B} = \sigma^a{}_{A^C} \sigma_C{}^b{}_{B^D}$.

There is a natural (canonical) spinorial connection associated with the metric such that $D_a \sigma^b = 0$. The pullback of the 4-dimensional canonical connection to the hypersurface is another connection on the 3-surface called *Sen Connection*, defined by

$$\mathcal{D}_a \alpha_A = q_a{}^b \nabla_b \alpha_A,$$

the curvature associated to this connection is denoted by $F_{abA}{}^B$ and is the pullback of the 4-dimensional curvature $F_{abA}{}^B = q_a{}^c q_b{}^d {}^4 R_{cdA}{}^B$. If we fix (for simplicity) a connection ∂ on the unprimed spinor indices, chosen to be real and flat, and replace ∇ by a $SL(2, C)$ Lie algebra-valued connection 1-form ${}^4 A_a{}^B$:

$$\nabla_a \lambda_A := \partial_a \lambda_A + {}^4 A_a{}^B \lambda_B,$$

the curvature tensor becomes

$${}^4 R_{ab} := 2\partial_{[a} {}^4 A_{b]} + [{}^4 A_a, {}^4 A_b].$$

Recall that ${}^4 R_{abA}{}^B$ is defined by

$$(\nabla_a \nabla_b - \nabla_b \nabla_a) \lambda_A = {}^4 R_{abA}{}^B \lambda_B,$$

thus we can write $F_{abC}{}^D$ in terms of a $SU(2)$ Lie algebra valued one form $A_a C^D$

$$F_{ab} := 2\partial_{[a} A_{b]} + [A_a, A_b].$$

Using these variables the action can be defined as:

$$S(\sigma, {}^4 A) = \int d^4 x ({}^4 \sigma) \sigma^a{}_{A^A'} \sigma^b{}_{B^A'} {}^4 R_{ab}{}^{AB},$$

where $({}^4 \sigma)$ denotes the square root of minus the determinant of the four metric, this action is presented in [9] and in [7]. Using the identity:

$$\begin{aligned} \sigma^a{}_{A^A'} \sigma^b{}_{B^A'} {}^4 R_{ab}{}^{AB} &= \text{tr} \left(\sqrt{2} i n^a \sigma^{b4} R_{ab} - \sigma^a \sigma^{b4} R_{ab} \right) \\ &= \text{tr} \left(-\sigma^a \sigma^{b4} R_{ab} + \sqrt{2} i N^{-1} \sigma^b \right. \\ &\quad \times (\mathcal{L}_t {}^4 A_b - \mathcal{D}_b ({}^4 A \cdot t)) \\ &\quad \left. - \sqrt{2} i N^{-1} N^a \sigma^{b4} R_{ab} \right), \end{aligned}$$

where ${}^4 A \cdot t = {}^4 A_a t^a$, we have

$$\begin{aligned} S &= \int dt \int d^3 x ({}^4 \sigma) \text{tr} \left(\sqrt{2} i N^{-1} \sigma^b (\mathcal{L}_t {}^4 A_b \right. \\ &\quad \left. - \mathcal{D}_b ({}^4 A \cdot t)) - \sigma^a \sigma^{b4} R_{ab} - \sqrt{2} i N^{-1} N^a \sigma^{b4} R_{ab} \right). \end{aligned}$$

Finally, since $\sigma^a{}_{AB}$ projects onto the 3-surface, and ${}^4 \sigma = N\sigma$, the action² becomes

¹For more definitions see [8].

²The surface terms are not included here.

$$S = \int dt \int d^3x \operatorname{tr} \left(\sqrt{2} i \tilde{\sigma}^b (\mathcal{L}_t A_b - \mathcal{D}_b({}^4 A \cdot t)) \right. \\ \left. - \mathbb{N} \tilde{\sigma}^a \tilde{\sigma}^b F_{ab} - \sqrt{2} i N^a \tilde{\sigma}^b F_{ab} \right).$$

with $\tilde{\sigma}^a{}_{AB} = (\sigma) \sigma^a{}_{AB}$ and $\mathbb{N} = (\sigma)^{-1} N$.

The action depends on five variables \mathbb{N} , N^a , ${}^4 A \cdot t$, $A_a{}^B$ and $\tilde{\sigma}^a{}_{AB}$. The first three variables play the role of the Lagrange multipliers, only the last two are dynamical variables. Varying the action with respect to the Lagrange multipliers we obtain the constraint equations:

$$\begin{aligned} C(\tilde{\sigma}, A) &:= \operatorname{tr}(\tilde{\sigma}^a \tilde{\sigma}^b F_{ab}) = 0, \\ C_a(\tilde{\sigma}, A) &:= \operatorname{tr}(\tilde{\sigma}^b F_{ab}) = 0, \\ \tilde{C}_A{}^B(\tilde{\sigma}, A) &:= \mathcal{D}_a \tilde{\sigma}^a{}^B = 0. \end{aligned} \quad (1)$$

varying with respect to the dynamical variables, yields the evolution equations:

$$\begin{aligned} \mathcal{L}_t \tilde{\sigma}^b &= [{}^4 A \cdot t, \tilde{\sigma}^b] + 2\mathcal{D}_a(N^{[a} \tilde{\sigma}^{b]}) \\ &\quad - \frac{1}{\sqrt{2}} i \mathcal{D}_a(\mathbb{N}[\tilde{\sigma}^a, \tilde{\sigma}^b]), \\ \mathcal{L}_t A_b &= \mathcal{D}_b({}^4 A \cdot t) + N^a F_{ab} + \frac{1}{\sqrt{2}} i \mathbb{N}[\tilde{\sigma}^a, F_{ba}]. \end{aligned} \quad (2)$$

III. SYMMETRIC HYPERBOLICITY

In order to obtain a symmetric hyperbolic system for any given lapse-shift pair, we suitably modify the field equations outside the constraint submanifold. Indeed, the modification only involves the addition of terms proportional to the constraints. Since we know that the evolution vector field is tangent to this manifold, the physical evolution, that is the dynamics inside the constraint manifold, remains unchanged. The modified evolution equations are

$$\begin{aligned} \mathcal{L}_t \tilde{\sigma}^b &= [{}^4 A \cdot t, \tilde{\sigma}^b] + 2\mathcal{D}_a(N^{[a} \tilde{\sigma}^{b]}) \\ &\quad - \frac{1}{\sqrt{2}} i \mathcal{D}_a(\mathbb{N}[\tilde{\sigma}^a, \tilde{\sigma}^b]) + \frac{1}{\sqrt{2}} i \mathbb{N}[\tilde{C}, \tilde{\sigma}^b] + N^b \tilde{C}, \\ \mathcal{L}_t A_b &= \mathcal{D}_b({}^4 A \cdot t) + N^a F_{ab} + \frac{1}{\sqrt{2}} i \mathbb{N}[\tilde{\sigma}^a, F_{ba}] \\ &\quad + \frac{i}{\sigma^2 \sqrt{2}} \mathbb{N} \tilde{\sigma}_b C + \frac{i}{\sigma^4} \mathbb{N} \tilde{\epsilon}_b{}^{dc} \tilde{\sigma}_c C_d. \end{aligned} \quad (3)$$

Since the Lagrange multiplier ${}^4 A_a t^a$ is an arbitrary function we can choose it as ${}^4 A_a t^a = A_a N^a$, i.e. we set its normal projection to zero. In this way, in the equation for the evolution of A_a , the principal part corresponding to the first two terms is replaced by the directional derivative of A_a .

To see that the system (3) is symmetric hyperbolic, we shall only need to consider its principal symbol. Recall that the principal symbol of a quasilinear evolution equation system

$$\dot{u}^l = B^l{}_j{}^a(u) \nabla_a u^j + M^l(u)$$

is given by $P(u, ik_a) = i B^l{}_j{}^a(u) k_a$. In our case u denotes $u = (\sigma^a, A_b)$.

Lemma III.1: *The equation system (3) for any fixed, but arbitrary lapse and shift fields is a symmetric hyperbolic system.*

Proof:

Since the vector u in our case is complex, symmetry of real systems corresponds to antihermiticity of the principal symbol with respect to some local hermitian inner product, i.e. an hermitian, positive definite bilinear form. In our case the natural one:

$$\begin{aligned} \langle u_2, u_1 \rangle &\equiv \langle (\tilde{\sigma}_2, A^2), (\tilde{\sigma}_1, A^1) \rangle \\ &\equiv \operatorname{tr}(\tilde{\sigma}_2^{\dagger a} \tilde{\sigma}_1^b) q_{ab} + \operatorname{tr}(A^2{}^{\dagger a} A^1{}_b) q^{ab}, \end{aligned}$$

is the one that works. Thus we need to prove that

$$P_{12} + P_{12}^{\dagger} \equiv \langle u_2, P u_1 \rangle + \langle u_2, P^{\dagger} u_1 \rangle = 0.$$

To obtain an expression for the symbol $P u_1$ we neglect all the terms which do not have derivatives of the dynamical variables $\tilde{\sigma}^b$ and A_b . We remark the *very important* fact that for both equations, (2) and (3), the principal part decouples into two blocks one acting on σ^b and the other acting on A_a , so we can write

$$P(\mathfrak{y}, ik_a) = P_{\sigma}(\mathfrak{y}, ik_a) \oplus P_A(\mathfrak{y}, ik_a), \quad (4)$$

in our case $\mathfrak{y} = (\tilde{\sigma}, A)$. The action of P on any solution u can be written as

$$\begin{aligned} P_{\sigma}(\tilde{\sigma}, ik_a) \tilde{\sigma}_1{}^b &= \frac{k_a}{\sqrt{2}} \mathbb{N}[\tilde{\sigma}^a, \tilde{\sigma}_1{}^b] + i k_a N^a \tilde{\sigma}_1{}^b, \\ P_A(\tilde{\sigma}, ik_c) A^1{}_a &= i N^b k_b A^1{}_a - \frac{1}{\sqrt{2}} \mathbb{N}[\tilde{\sigma}^b, k_a A^1{}_b] \\ &\quad + \frac{1}{\sqrt{2}} \mathbb{N}[\tilde{\sigma}^b, k_b A^1{}_a] \\ &\quad - \frac{1}{\sqrt{2}} \mathbb{N}[\tilde{\sigma}_a \operatorname{tr}([\tilde{\sigma}^d, \tilde{\sigma}^b] k_d A^1{}_b)] \\ &\quad - \mathbb{N}[\tilde{\epsilon}_a{}^{bc} \tilde{\sigma}_c \operatorname{tr}(\tilde{\sigma}^d k_b A^1{}_d)] \\ &\quad + \mathbb{N}[\tilde{\epsilon}_a{}^{bc} \tilde{\sigma}_c \operatorname{tr}(\tilde{\sigma}^d k_d A^1{}_b)]. \end{aligned} \quad (5)$$

Adding the expressions for $P_{\sigma 12}$ and $P_{\sigma 12}^{\dagger}$ we obtain

$$\begin{aligned} P_{\sigma 12} + P_{\sigma 12}^{\dagger} &\equiv \langle \tilde{\sigma}_2, P_{\sigma} \tilde{\sigma}_1 \rangle + \langle \tilde{\sigma}_2, P_{\sigma}^{\dagger} \tilde{\sigma}_1 \rangle \\ &= -\frac{k_a}{\sqrt{2}} \mathbb{N} \operatorname{tr}(\tilde{\sigma}^a [\tilde{\sigma}_1{}^b, \tilde{\sigma}_2{}_b]) \\ &\quad + \frac{k_a}{\sqrt{2}} \mathbb{N} \operatorname{tr}(\tilde{\sigma}^a [\tilde{\sigma}_1{}_b, \tilde{\sigma}_2{}^b]) \\ &\quad - i k_a N^a \operatorname{tr}(\tilde{\sigma}_2{}_b \tilde{\sigma}_1{}^b) + i k_a N^a \operatorname{tr}(\tilde{\sigma}_2{}^b \tilde{\sigma}_1{}_b) \\ &= 0. \end{aligned} \quad (6)$$

here we have used that $\tilde{\sigma}^{a\dagger} = -\tilde{\sigma}^a$. It only remains to prove a similar result for P_{A12} , using in the expression for $P_A A^1_a$ that

$$A^1_a = -\frac{1}{\sigma^2} \text{tr}(A^1_a \tilde{\sigma}_e) \tilde{\sigma}^e$$

and

$$[\tilde{\sigma}^a, \tilde{\sigma}^b]_{A^B} = \frac{\sqrt{2}}{\sigma^2} \tilde{\epsilon}^{ab} \tilde{\sigma}^c A^B_c,$$

we obtain

$$\begin{aligned} \frac{\sigma^4}{\mathbb{N}} P_A A^1_a &= \tilde{\epsilon}^{bem} \tilde{\sigma}_m k_a \text{tr}(A^1_b \tilde{\sigma}_e) - \frac{\sigma^2 i}{\mathbb{N}} N^b k_b \text{tr}(A^1_a \tilde{\sigma}_e) \tilde{\sigma}^e \\ &\quad - \tilde{\epsilon}^{bem} \tilde{\sigma}_m k_b \text{tr}(A^1_a \tilde{\sigma}_e) - \tilde{\epsilon}^{dbe} \tilde{\sigma}_a k_d \text{tr}(A^1_b \tilde{\sigma}_e) \\ &\quad - \tilde{\epsilon}_a^{bc} \tilde{\sigma}_c k_b \text{tr}(A^1_d \tilde{\sigma}^d) + \tilde{\epsilon}_a^{bc} \tilde{\sigma}_c k_d \text{tr}(A^1_b \tilde{\sigma}^d). \end{aligned}$$

Thus we have

$$\begin{aligned} \frac{\sigma^4}{\mathbb{N}} (P_{A12} + P^\dagger_{A12}) &= -\frac{\sigma^2 i}{\mathbb{N}} N^b k_b \text{tr}(A^{2\dagger a} \tilde{\sigma}^e) \text{tr}(A^1_a \tilde{\sigma}_e) \\ &\quad + \frac{\sigma^2 i}{\mathbb{N}} N^b k_b \text{tr}(A^{2\dagger a} \tilde{\sigma}^e) \text{tr}(A^1_a \tilde{\sigma}_e) \\ &\quad + \tilde{\epsilon}^{bem} \text{tr}(A^{2\dagger a} \tilde{\sigma}_m) k_a \text{tr}(A^1_b \tilde{\sigma}_e) \\ &\quad - \tilde{\epsilon}^{bem} \text{tr}(A^{2\dagger a} \tilde{\sigma}_m) k_b \text{tr}(A^1_a \tilde{\sigma}_e) \\ &\quad - \tilde{\epsilon}^{dbe} \text{tr}(A^{2\dagger a} \tilde{\sigma}_a) k_d \text{tr}(A^1_b \tilde{\sigma}_e) \\ &\quad - \tilde{\epsilon}_a^{bc} \text{tr}(A^{2\dagger a} \tilde{\sigma}_c) k_b \text{tr}(A^1_d \tilde{\sigma}^d) \\ &\quad + \tilde{\epsilon}_a^{bc} \text{tr}(A^{2\dagger a} \tilde{\sigma}_c) k_d \text{tr}(A^1_b \tilde{\sigma}^d) \quad (7) \\ &\quad + \tilde{\epsilon}^{bem} \text{tr}(A^{2\dagger b} \tilde{\sigma}_e) k_a \text{tr}(A^1_a \tilde{\sigma}_m) \\ &\quad - \tilde{\epsilon}^{bem} \text{tr}(A^{2\dagger a} \tilde{\sigma}_e) k_b \text{tr}(A^1_a \tilde{\sigma}_m) \\ &\quad - \tilde{\epsilon}^{dbe} \text{tr}(A^{2\dagger b} \tilde{\sigma}_e) k_d \text{tr}(A^1_a \tilde{\sigma}_a) \\ &\quad - \tilde{\epsilon}_a^{bc} \text{tr}(A^{2\dagger d} \tilde{\sigma}_d) k_b \text{tr}(A^1_a \tilde{\sigma}_c) \\ &\quad + \tilde{\epsilon}_a^{bc} \text{tr}(A^{2\dagger b} \tilde{\sigma}^d) k_d \text{tr}(A^1_a \tilde{\sigma}_c). \end{aligned}$$

Rearranging the indices and using the antisymmetry of the Levi-Civita tensor density we get that the first term cancels out the second, the 4th cancels out the 9th, and the 5th and 6th cancel 10th and 11th.

Finally, using

$$2A^{[a} \tilde{\sigma}^{b]} = \frac{1}{\sigma^2} \tilde{\epsilon}^{abe} \tilde{\epsilon}_{dce} A^d \tilde{\sigma}^c,$$

the remaining terms vanish, yielding the antihermiticity of the principal symbol. This concludes the proof of the Lemma. ♠

IV. CONCLUDING REMARKS

The result obtained here assumes, for simplicity, a given and fixed, lapse-shift pair. This assumption is not

very practical, for one would like, for instance, to choose the pair along the evolution in order to avoid the occurrence of spurious coordinate singularities. There are at least two ways to loosen the assumption made. The first is to incorporate the lapse-shift pair into the evolution equations by choosing appropriate equations for them in such a way as to obtain a bigger symmetric hyperbolic system. The second is to impose some elliptic equations on the pair in such a way as to be able to estimate the L^2 norm of the derivatives of it -for they appear on the equations for $(\tilde{\sigma}, A)$ - in terms of the L^2 norms of $(\tilde{\sigma}, A)$. This would be the case if one were to use Witten's equation to determine the pair. Besides these two ways of fixing the lapse-shift pairs along evolution there does not seem to be any other obvious candidate.

-
- [1] *New Hamiltonian formulation of general relativity*, A. Ashtekar. Physical Review D, **36**, 6, 1587, 1987.
 - [2] *Hyperbolic reductions for Einstein's equations*, H. Friedrich, Class. Quantum Grav., **13**, 1451-1469, 1996.
 - [3] *Geometrical Hyperbolic systems for General relativity and Gauge theories*, A. Abraham-A. Anderson-Y. Choquet Bruhat-J. York, Class. Quantum Grav., **14**, A9-A22, 1997.
 - [4] *New Formalism for Numerical Relativity*, C. Bona-J. Massó-E. Seidel-J. Stela, Physical Review Letters, **75**, 4, 600-603, 1995.
 - [5] *Horizon Boundary condition for black hole spacetime*, P. Anninos-G. Daues-J. Massó-E. Seidel-Wai-Mo Suen, Physical Review D, **51**, 10, 5562-5578, 1995.
 - [6] *First-Order Symmetric Hyperbolic Einstein's Equations with Arbitrary Fixed Gauge*, S. Frittelli and O. Reula, Physic. Rev. Letters, **76**, 4667-4670, 1996.
 - [7] *New Perspectives in Canonical Gravity*, A. Ashtekar, edited by Bibliopolis, 1988.
 - [8] *Quantum Theory of Spin-3/2 Fields in Einstein Spaces*, A. Sen. International Journal of Theoretical Physics, **21**, 1, 1982.
 - [9] *Covariant Action for Ashtekar's Form of Canonical Gravity*, T. Jacobson and L. Smolin, Class. Quantum Grav., **5**, 583-594, 1988.
 - [10] *Spinors and space-time*, R. Penrose and W. Rindler, edited by Cambridge University Press, **1**, 1984.