THE 'UPS' AND 'DOWNS' OF A SPINNING BLACK HOLE

C.M. CHAMBERS, W.A. HISCOCK, B.E. TAYLOR

Department of Physics, Montana State University, Bozeman, MT 59717-3840, USA

We report and comment upon the principal results of an investigation into the evolution of rotating black holes emitting massless scalar radiation via the Hawking process. It is demonstrated that a Kerr black hole evaporating by the emission of scalar radiation will evolve towards a state with $a \approx 0.555M$. If the initial specific angular momentum is larger than this value the hole will spin down to this value; if it is less it will spin up to this value. The addition of higher spin fields to the picture strongly suggests the final asymptotic state of a realistic evaporation process will be characterized by an a/M = 0.

1 Introduction

It is generally accepted that an isolated black hole evaporating via the Hawking radiation mechanism will, regardless of its initial state, approach a final asymptotic state described by the Schwarzschild solution. Indeed, the evolution of rotating black holes radiating fields of spin s has been studied in some detail by Page.¹ For a Kerr black hole, emitting a fixed set of fields with s > 0, Page has shown that the angular momentum loss proceeds more quickly than the mass loss, resulting in a final evolutionary state with $a_* \equiv a/M = 0$, which is indeed characterized by the Schwarzschild solution. However, Page's work does suggest that a black hole emitting scalar fields (s = 0) might evolve differently and approach a state with $a_* > 0$. For s > 0, it was found that the dominant emission modes at low a_* were those with $\ell = m = s$.^a If the same holds true for scalar fields then the dominant mode will be $\ell = 0$, which carries off energy but no angular momentum. In that case, one can envisage a situation in which the mass loss occurs much more rapidly than the angular momentum loss and leads to a final state with a_* nonzero.

Here, we summarize the main results and conclusions of a numerical study into the evolutionary path of a rotating black hole emitting massless scalar fields via the Hawking process.² For a Kerr black hole emitting only scalar radiation, we find the rate at which mass and angular momentum are lost leads to a final evolutionary state characterized by an $a_* \approx 0.555$. Further to this we determine the number of scalar fields required to cause a black hole, emitting a fixed set of higher spin fields, to evolve to a state of nonzero a_* .

2 Results

The evolution of an isolated Kerr black hole, evaporating via the Hawking mechanism, is represented by the rates at which its mass (M) and angular momentum (J) decrease. Following Page we express these rates in terms of the scale invariant quantities $f(a_*) \equiv -M^2 dM/dt$ and $g(a_*) \equiv -Ma_*^{-1} dJ/dt$, where t is the usual Boyer-Lindquist time coordinate for the Kerr spacetime.^b Figure 1 shows the scale

^{*a*}Here ℓ and *m* are the usual angular eigenvalues associated with orbital angular momentum.

 $[^]b\mathrm{Expressions}$ for f and g, and the numerical procedures used to calculate them, can be found in the work of Taylor et~al. 3




Figure 1: The scale invariant mass and angular momentum loss rates, f and g, versus a_* due to a single scalar field. The behavior of these functions is explained in the text.

invariant mass loss rate f and the scale invariant angular momentum loss rate g of a Kerr black hole evaporating via purely scalar radiation. Unlike the higher spin case,¹ where both f and q are monotonically increasing functions of a_* , the mass loss rate due to the emission of massless scalar radiation alone, can clearly be seen to possess a minimum at $a_* \approx 0.574$. This exceptional behavior can be ascribed to the fact that a scalar field, unlike higher spin fields, is able to radiate in an $\ell = 0$ mode, which is not superradiant. Our numerical study shows that the mass loss rate, due solely to the $\ell = 0$ mode, is a monotonically decreasing function of a_* and that at low a_* this mode dominates the mass loss experienced by the black hole. As a_* , and hence superradiance, increases the contribution of higher ℓ modes to the mass loss becomes significant and dominates at high a_* . The combined effect at intermediate a_* is the observed minimum. The angular momentum loss rate q can be seen to be a monotonically increasing function of a_* . Since the $\ell = 0$ mode cannot carry off angular momentum, the behavior of g can be understood purely in terms of superradiance. As a_* and the effects of superradiance increase more angular momentum is efficiently extracted from the hole.

In order to gauge the relative loss rate of mass to angular momentum, we define the scale invariant quantity, $h(a_*)$, by

$$h(a_*) \equiv \frac{d\ln a_*}{d\ln M} = \frac{g(a_*)}{f(a_*)} - 2.$$
 (1)

Figure 2 shows $h(a_*)$ versus a_* for a Kerr black hole emitting purely scalar radiation. The important feature of this plot is the existence of a zero in h at a finite, but nonzero, value of a_* , namely $a_* \approx 0.555$. The rate of change of a_* with respect to time is given by

$$\frac{da_*}{dt} = -\frac{a_*hf}{M^3} \,. \tag{2}$$

Since the black hole is in isolation it can gain neither mass nor angular momentum and so f and g, by definition, must be positive definite functions of a_* – as demonstrated in Figure 1. A black hole with $0.555 < a_* < 1$ will have a positive value of h




Figure 2: The quantity h is plotted versus a_* . The zero of h indicates that a black hole emitting purely massless scalar radiation will evolve towards a state described by a value of $a_* \approx 0.555$.

and thus Eq. 2 implies that a_* will decrease over time until it reaches an asymptotic state characterized by an $a_* \approx 0.555$. Similarly, a black hole with $0 < a_* < 0.555$ has a negative value of h and so a_* will increase until the black hole achieves a state with $a_* \approx 0.555$. Therefore, the existence of a zero in h at $a_* = 0.555$ implies that a Kerr black hole will evolve to and remain at this value of a_* .

It is interesting to ask to what extent the inclusion of scalar field emission in the evaporation processes studied by Page affects the evolution. Utilizing the results for higher spin fields,¹ we have calculated the number of scalar fields in conjunction with a fixed set of higher spin fields that results in a final state with $a_* > 0$. For the set consisting of 1 spin-2, 1 spin-1 and 3 spin-1/2 fields, we find that 32 scalar fields are required. As no massless scalar fields have yet been discovered in Nature it appears that the conventional view of black hole evaporation will continue to hold in any realistic evaporation process.

Acknowledgments

CMC is a fellow of The Royal Commission For The Exhibition Of 1851 and gratefully acknowledges their financial support. Travel to Israel for CMC was supported by NSF Grant Nos. PHY-9722529 to North Carolina State University, and PHY-9511794 to Montana State University. The work of WAH and BT was supported in part by NSF Grant No. PHY-9511794.

References

- 1. D.N. Page, *Phys. Rev.* D 14, 3260 (1976).
- C.M. Chambers, W.A. Hiscock and B. Taylor, *Phys. Rev. Lett.* 78, 3249 (1997).
- 3. B.E. Taylor, C.M. Chambers and W.A. Hiscock, in preparation.