P. A. Hogan

Mathematical Physics Department, University College Dublin, Belfield, Dublin 4, Ireland

and

C. Barrabès and G. F. Bressange Laboratoire de Mathématiques et Physique théorique UPRES-A 6083, CNRS

Université de Tours, 37200 France.

PACS numbers: 04.30.+x, 04.20Jb

Recently [1] a simple solution of the vacuum Einstein–Maxwell field equations was given describing a plane electromagnetic shock wave sharing its wave front with a plane gravitational impulse wave. We present here an exact solution of the vacuum Einstein–Maxwell field equations describing the head–on collision of such a wave with a plane gravitational impulse wave. The solution has the Penrose–Khan solution and a solution obtained by Griffiths as separate limiting cases. In a recent paper[1] a construction is given of a solution of the vacuum Einstein–Maxwell field equations describing a plane electromagnetic shock wave sharing its wave front with a plane gravitational impulse wave. The wave is simpler than previous examples of such objects (see, for example [2], which is discussed in [1]) in that the space–time on one side of the null hypersurface history of the wave front is conformally flat (and is a special case of a Bertotti– Robinson[3] space–time) and on the other side is flat.The homogeneous special case of this wave has line–element which can be put in the form

$$ds^{2} = (1 + b^{2}v_{+}^{2})^{-1} \left\{ 2du \, dv - (1 + lv_{+})^{2}dx^{2} - (1 - lv_{+})^{2}dy^{2} \right\} , \qquad (1)$$

where b, l are constants,  $v_+ = v \theta(v)$  with  $\theta(v)$  the Heaviside step function (equal to 1 for v > 0 and equal to zero for v < 0). For this space-time the only non-vanishing Newman-Penrose component of the Maxwell field is

$$\phi_0 = \frac{b\,\theta(v)}{1+b^2v_+^2} \,\,, \tag{2}$$

and the only non-vanishing Newman-Penrose component of the Weyl tensor is

$$\Psi_0 = -l\,\delta(v) \ . \tag{3}$$

Here  $\delta(v)$  is the Dirac delta function. Thus both the Maxwell and Weyl tensors are type N in the Petrov classification with  $\partial/\partial u$  as degenerate principal null direction. The null hypersurface v = 0 is a null hyper*plane* and is the history of a plane electromagnetic shock wave on account of (2) and of a plane gravitational impulse wave on account of (3). We can remove the shock by putting b = 0 and we can remove the gravitational impulse wave by putting l = 0. We consider now the head—on collision of a wave of the type described by (1) with a plane gravitational impulsive wave. This latter will be described by the space-time with line-element

$$ds^{2} = 2du \, dv - (1 + ku_{+})^{2} dx^{2} - (1 - ku_{+})^{2} dy^{2} , \qquad (4)$$

with k a constant and  $u_{+} = u \theta(u)$ . Following the usual procedure in setting up such a collision problem (see [4]) we consider the space-time to have lineelement (1) in the region u < 0 and have line-element (4) for v < 0 (the two line-elements coincide in the overlapping region u < 0, v < 0). The line-element in the region u > 0, v > 0 (after the collision) has the Rosen-Szekeres form [4]

$$ds^{2} = 2e^{-M} du \, dv - e^{-U} \left( e^{V} dx^{2} + e^{-V} dy^{2} \right) , \qquad (5)$$

where M, U, V are each functions of (u, v) satisfying the O'Brien–Synge [5] junction conditions: When v = 0

$$e^{V} = \frac{1+ku}{1-ku}$$
,  $e^{M} = 1$ ,  $e^{-U} = 1-k^{2}u^{2}$ , (6)

and when u = 0

$$e^{V} = \frac{1+lv}{1-lv}$$
,  $e^{M} = 1+b^{2}v^{2}$ ,  $e^{-U} = \frac{1-l^{2}v^{2}}{1+b^{2}v^{2}}$ . (7)

In addition the Maxwell field in the region u > 0, v > 0 has two non-vanishing Newman-Penrose components [4]  $\phi_0, \phi_2$  which are both functions of (u, v) and satisfy the boundary conditions: when  $v = 0, \phi_2 = 0$  and when  $u = 0, \phi_0 = b(1 + b^2v^2)^{-1}$ . It is now a matter of solving the vacuum Einstein-Maxwell equations in the region u > 0, v > 0 (these can be found in [4] for example) for the unknown functions  $U, V, M, \phi_2, \phi_0$  subject to the above boundary conditions. We find the following expressions for these functions:

$$e^{-U} = \frac{F}{1 + b^2 v^2} , \qquad (8)$$

$$e^{V} = \frac{1 + ku\sqrt{1 - l^{2}v^{2}} + lv\sqrt{1 - k^{2}u^{2}}}{1 - ku\sqrt{1 - l^{2}v^{2}} - lv\sqrt{1 - k^{2}u^{2}}} , \qquad (9)$$

$$e^{-M} = \frac{H^2}{(1+b^2v^2)\left[(1-k^2u^2)(1-l^2v^2)F\right]^{1/2}},$$
(10)

$$\phi_2 = \frac{-kbv\sqrt{1-l^2v^2}}{\left[(1-k^2u^2)F\right]^{1/2}H} , \qquad (11)$$

and

$$\phi_0 = \frac{b\left\{(l^2 + b^2)lkuv^3 + \sqrt{1 - k^2u^2}(1 - l^2v^2)^{3/2}\right\}}{(1 + b^2v^2)\left[(1 - l^2v^2)F\right]^{1/2}H} , \qquad (12)$$

where

$$F = 1 - k^2 u^2 - l^2 v^2 - k^2 b^2 u^2 v^2 , \qquad (13)$$

and

$$H = \sqrt{1 - k^2 u^2} \sqrt{1 - l^2 v^2} - k l u v .$$
 (14)

A calculation of the Weyl tensor components reveals the expected curvature singularity at F = 0 for u > 0, v > 0. There are two important special limiting cases: (1) if b = 0 the solution above becomes the Penrose–Khan [6] solution describing the space–time following the collision of two plane impulsive gravitational waves and (2) if l = 0 the solution becomes the Griffiths [4, 7] solution describing the space–time following the collision of a plane gravitational impulse wave and a plane electromagnetic shock wave. Clearly further collisions involving the type of plane wave described here by (1) can be envisaged. We thank Professor Werner Israel for helpful and encouraging discussions.

## **References:**

- Hogan, P. A. 1997 "Plane Gravitational Waves and Bertotti–Robinson Space–Times", University College Dublin, preprint.
- [2] Chandrasekhar, S. and Xanthopoulos, B. C. 1985 Proc. R. Soc. A398, 223.
- [3] Kramer, D., Stephani, H., Mac Callum, M. A. H. and Herlt, E. Exact Solutions of Einstein's Equations (VEB Deutscher Verlag der Wissenschaften, Berlin 1980).
- [4] Griffiths, J. B. Colliding Plane Waves in General Relativity (Clarendon Press, Oxford 1991).
- [5] O'Brien, S. and Synge, J. L. 1952 Commun. Dublin Inst. Adv. Stud. A, no.9.
- [6] Penrose, R. and Khan, K. A. 1971 Nature 229, 185.
- [7] Griffiths, J. B. 1975 Phys. Lett., A54, 269.