Critical Collapse of Skyrmions

Piotr Bizoń*

Department of Mathematics, University of Michigan, Ann Arbor, Michigan

Tadeusz Chmaj Institute of Nuclear Physics, Cracow, Poland (July 8, 2021)

We study first order phase transitions in the gravitational collapse of spherically symmetric skyrmions. Static sphaleron solutions are shown to play the role of critical solutions separating black-hole spacetimes from no-black-hole spacetimes. In particular, we find a new type of first order phase transition where subcritical data do not disperse but evolve towards a static regular stable solution. We also demonstrate explicitly that the near-critical solutions depart from the intermediate asymptotic regime along the unstable manifold of the critical solution.

04.25.Dm, 04.40.-b, 04.70.Bw

The understanding of the dynamics of gravitational collapse is a major theoretical challange in classical general relativity. A few years ago a new twist into this problem was provided by Choptuik's numerical discovery of critical behaviour in the gravitational collapse of spherically symmetric massless scalar field [1]. Analysing the evolution of one-parameter families of regular initial data, Choptuik showed that the transition between collapse and dispersion has a universal character which is reminiscent of a second order phase transition with the continuous change of black-hole mass. Subsequently, similar critical behaviour has been observed in several other models of gravitational collapse, and significant progress has been made towards the analytical understanding of the numerical phenomenology (see [2] for a recent review and bibliography).

Recently, a qualitatively new type of critical collapse was found in the Einstein-Yang-Mills system [3], where, for some initial data, the fundamental Bartnik-McKinnon sphaleron (static soliton with one unstable mode) plays the role of a critical solution. In this case there is a gap in the spectrum of black-hole masses which is reminiscent of the first order phase transition.

Surprisingly, none of the models studied so far had a *stable* static solution (a "star") and consequently the possible endpoints of evolution were basically reduced to the collapse/dispersion alternative. As far as we know, the only exception is the collapse/oscillaton alternative recently found in the collapse of massive scalar field [4]. If a model has a star-like solution, then there will be an open set of initial data which evolve towards this solution (basin of attraction) and it is natural to expect the occurence of critical behaviour at the boundary of this set.

The Einstein-Skyrme model which possesses the whole zoo of static regular solutions, both stable and unstable, seems to be a suitable testing ground to address this issue. In this letter we focus our attention on a new type of first order phase transition we found in this model for baryon-number-one data. We also briefly discuss a novel feature of mass gap in the baryon-number-zero sector. A detailed description of both first and second order phase transitions will be published elsewhere [5].

We consider the Einstein-Skyrme system, so the matter in our model is an SU(2)-valued scalar function U(x)(called a chiral field) with dynamics given by the Lagrangian

$$L = \frac{f^2}{4} Tr(\nabla_a \nabla^a U^{-1}) + \frac{1}{32e^2} Tr[(\nabla_a U)U^{-1}, (\nabla_b U)U^{-1}]^2, \quad (1)$$

where ∇_a is the covariant derivative with respect to the spacetime metric. The two coupling constants f and e are hereafter set to one, which amounts to using f/e and 1/ef as the units of mass and length, respectively.

We specialize to spherical symmetry. For the metric we use the polar time slicing and the areal radial coordinate

$$ds^{2} = -e^{-2\delta(r,t)}N(r,t)dt^{2} + N^{-1}(r,t)dr^{2} + r^{2}d\Omega^{2}.$$
 (2)

For the chiral field we assume the hedgehog ansatz $U = exp(i\vec{\sigma} \cdot \hat{r}F(r,t))$, where $\vec{\sigma}$ is the vector of Pauli matrices. Using overdots and primes to denote $\partial/\partial t$ and $\partial/\partial r$ respectively, we introduce an auxilary variable $P = ue^{\delta}N^{-1}\dot{F}$, where $u = r^2 + 2\sin^2 F$. Then, the Einstein-Skyrme equations reduce to

$$\dot{F} = e^{-\delta} N \frac{P}{u},\tag{3}$$

$$\dot{P} = (e^{-\delta} N u F')' + \sin(2F) e^{-\delta} \left(N (\frac{P^2}{u^2} - F'^2) - \frac{u}{2r^2} - \frac{1}{2} \right), \quad (4)$$

$$\dot{N} = -\frac{2\alpha}{r}e^{-\delta}N^2 P F', \qquad (5)$$

$$\delta' = -\frac{\alpha u}{r} \left(\frac{P^2}{u^2} + F'^2\right),\tag{6}$$

$$N' = \frac{1-N}{r} - \frac{\alpha}{r} \left(2\sin^2 F + \frac{\sin^4 F}{r^2} + uN(\frac{P^2}{u^2} + F'^2) \right).$$
(7)

Here $\alpha = 4\pi G f^2$ is the dimensionless coupling constant. In order to solve the initial value problem numerically we have implemented a free evolution scheme using a second order finite difference method on uniform grid. The conservation of the hamiltonian constraint (7) was used only to check the accuracy of the code. To ensure regularity at the center we impose the boundary condition F(r,t) = O(r) for $r \to 0$. To minimize reflections from the outer boundary of the grid we impose a sort of outgoing wave condition there. Asymptotic flatness requires that $F(r,0) = B\pi + O(1/r^2)$ at infinity, where the integer B, called the baryon number, is equal to the topological degree of the chiral field. As long as no horizon forms, the baryon number of initial data is automatically preserved during the evolution.

Below we describe our results for regular initial data with baryon number one and zero.

B = 1: Let us recall [6] that in this case, for $\alpha < \alpha_1 \simeq 0.040378$, the Eqs.(3-7) have two regular static solutions (solitons) which we denote below by X^s and X^u with X standing for (F, P, N, δ) . The solution X^s is linearly stable - it is the skyrmion distorted by gravity. The second solution X^u has one linearly unstable mode. These two solutions coalesce at the bifurcation point $\alpha = \alpha_1$, and disappear for $\alpha > \alpha_1$. For given α the mass m_u of the unstable soliton is larger than the mass m_s of the stable soliton (for example, for $\alpha = 0.02$ we have $m_s \simeq 68.05$ and $m_u \simeq 80.70$). As $\alpha \to 0$, m_s tends (from below) to the mass of the flat-space skyrmion ($\simeq 72.92$), whereas $m_u \to \infty$, which is the manifestation of a nonperturbative character of the solution X^u .

Since for B = 1 the dispersion of the chiral field to infinity is forbidden, generic initial data end up either as black holes or as the stable soliton X^s . The respective basins of attraction of these two final states depend on the coupling constant α . This is illustrated in Fig. 1 for a typical time-symmetric kink-type initial data.



FIG. 1. Basins of attraction of the one-parameter family of B = 1 initial data $F_p(r, 0) = \pi \tanh(r/p)$ as a function of the coupling constant α . Critical initial data are represented by the solid line curve. Initial data lying in the region enclosed by the critical curve and the *p*-axis evolve towards the skyrmion X^s , while initial data lying outside that region form a black hole. As $\alpha \to 0$ (not shown in the figure) the basin of attraction of X^s tends to the whole positive *p*-axis.

Notice that for $\alpha < \alpha_1$ there are two critical configurations along the one-parameter family F_p (the points A and B on the dashed line segment on Fig. 1). Since the parameter p (the width of a kink) is inversely proportional to the "condensation" of initial mass, the transition at the point A could be interpreted as a "weak/strong" field transition. However at the point Bit is the "weaker" configuration that collapses, simply because it is unable to get rid of the excess mass by the time it shrinks to size of the stable soliton. This behaviour indicates that it is difficult to formulate a criterion for collapse in terms of initial data alone.

The critical initial data lying on the boundary of the basin of attraction of X^s asymptote to the unstable solution X^u . In other words, the codimension one stable manifold of the solution X^u divides (locally) the phase space into collapsing and non-collapsing data. Near-critical data approach X^u , stay in its vicinity for some time, and eventually collapse to a black hole or decay into X^s . This is shown in Fig. 2 for a marginally subcritical data (lying slightly below the point *B* on Fig. 1).



FIG. 2. A sequence of profiles of F(r, t) (solid lines) versus $\ln(r)$ from a marginally subcritical evolution of the B = 1 initial data. The stable and unstable static solutions are represented by the dotted and dashed lines, respectively. Here t is the proper time at spatial infinity (we use the normalization $\delta(\infty, t) = 0$).

During the final stage of the evolution of a marginally supercritical solution (i.e. when the solution runs away from the vicinity of the critical solution and forms a black hole) almost no energy is radiated away to infinity. Therefore the mass gap in this first order phase transition equals m_u (up to 0.1%).

For subcritical data the process of settling down to the stable soliton seems to have a universal late-time behaviour which is dominated by a fundamental quasinormal mode. The parameters of this mode depend strongly on α . Having convinced ourselves that these parameters are indeed universal, that is do not depend on the excitation, we have determined them from the evolution of finite perturbations of X^s (see Fig. 3). The details of such perturbation are washed out very rapidly and a characteristic exponentially damped "ringing" dominates the late time evolution.



FIG. 3. Quasinormal ringing about the stable soliton for $\alpha = 0.03$. We plot $dN = |N(r_0, t) - N^s(r_0)|$ from the evolution of initial data $F(r, 0) = \pi \tanh(r)$ (dots). Here $r_0 = 0.8$. The least squares fit for the times t > 10 (represented by the solid line) gives the frequency $\omega \simeq 0.494$ and the damping rate $\tau \simeq 11.69$.

The above results are in perfect agreement with linear [7,6] and nonlinear [8] stability analysis of gravitating skyrmions. In the intermediate asymptotics the evolution of near-critical data is well approximated by the linearization about X^u

$$X_p(r,t) \approx X^u(r) + C(p-p^*)e^{\lambda t}\delta X^u(r) + \frac{\text{decaying}}{\text{modes}}, \quad (8)$$

where $\delta X^u(r)$ is the single unstable eigenmode associated with a positive eigenvalue λ . Depending on whether $p > p^*$ or $p < p^*$, the solution X_p eventually collapses to a black hole or decays to X^s . The "lifetime" T of a near-critical solution staying in the vicinity of X^u is determined by the time in which the amplitude of the unstable mode grows to a finite size: $|p - p^*|e^{\lambda T} \sim O(1)$, which gives $T \sim -\lambda^{-1} \ln |p - p^*|$. Thus, the larger λ , the better fine-tuning is required to see the solution X^u clearly pronounced as the intermediate attractor.

We have verified the formula (8) by comparing the unstable eigenmode δX^u with the snapshots of a nearcritical solution departing from the vicinity of X^u . This is shown in Fig. 4 for the same initial data as in Fig. 2. For t = 24 the profile of $\dot{F}(r, 24)$ is practically indistinguishable from the (suitably normalized) unstable eigenmode $\delta F^u(r)$. A slight deviation of $\dot{F}(r, 23)$ from $\delta F^u(r)$ for large r is due to the fact that by t = 23 the decaying modes in (8) have not yet died away completely. On the other hand, for t = 25 the deviation for small r signals the onset of nonlinear regime.



FIG. 4. The snapshots of $\dot{F}(r,t)$ (solid lines) departing from the vicinity of the critical solution X^u . On each snapshot we superimpose (dashed lines) the unstable eigenmode $\delta F^u(r)$ computed via linear perturbation analysis. For t = 24the amplitude of $\delta F^u(r)$ is normalized to the maximum of $\dot{F}(r, 24)$. The corresponding amplitudes for t = 23 and t = 25are rescaled by the factors $e^{-\lambda}$ and e^{λ} , respectively.

Using (8), we have also computed the eigenvalue λ directly from the nonlinear evolution by monitoring $F_p(r, 24)$ at several discrete radii r_i , i = 1, ..., n, and evaluating the averaged quantity $\frac{1}{n\Delta t} \sum \ln \left(\dot{F}_p(r_i, 24 + \Delta t) / \dot{F}_p(r_i, 24) \right)$ for some small Δt . The result agrees up to three decimal places with the eigenvalue obtained via linear stability analysis.

B = 0: In this topological sector, the Eqs.(3-7) have a pair of static regular solutions for $\alpha < \alpha_0 \simeq 0.00147$ [6]. One of them has two unstable modes, hence it plays no role in the evolution of generic initial data. The second solution, call it Y^u , has one unstable mode, so it is a candidate for the critical solution. Indeed, analysing the ingoing "generalized Gaussian" profile F(r, 0) = $Ar^3 e^{-(r-r_0)^4/\Delta^4}$, we have found Y^u as the intermediate attractor at the border between collapse and dispersion. Since in this case the critical behaviour is analogous to the first order phase transition in the EYM model [3], here we discuss only the issue of mass gap in this transition. Analysing near-supercritical solutions we found that, in contrast to the previously studied cases, the mass gap is *not* equal to the mass of the sphaleron. In order to determine the mass gap more precisely we adopted the following strategy. Rather than improving the accuracy of fine-tuning (which is computationally time-consuming because the eigenvalue of the unstable eigenmode is rather large), we began the evolution with specially prepared initial data of the form of the static solution Y^u plus the unstable eigenmode with a small positive amplitude: $Y(r, 0) = Y^u(r) + \epsilon \delta Y^u(r)$. In other words, we pushed the static solution along its unstable manifold and let it collapse. It turned out that the mass

gap is considerably smaller than the mass of the critical solution (which of course sets the upper bound for a mass gap). This is illustrated in Fig. 5, where one can clearly see how a substantial amount of the initial mass is being radiated away to infinity. The resulting black hole has the mass $\simeq 253.1$, which is 75.6% of the mass of Y^u .



ln(r)

FIG. 5. The mass aspect $\mu(r,t)$ (defined by $N = 1 - 2\mu/r$) from the evolution of the static solution Y^u perturbed along the unstable direction (for $\alpha = 0.00145$). The mass inside radius r at time t measured in units of f/e is given by $m(r,t) = \frac{4\pi}{\alpha}\mu(r,t)$. The plateau of $\mu(r,t)$ determines the mass gap.

Concluding, we believe that the results presented here have clarified certain aspects of first order phase transitions in gravitational collapse and, in combination with the results of [3] and [4], they help to understand which features of this phenomenon are generic.

Acknowledgments. This research was supported in part by the KBN grant PB750/P3/94/06.

- * On leave of absence from Institute of Physics, Jagiellonian University, Cracow, Poland.
- [1] M. W. Choptuik, Phys. Rev. Lett. 70, 9-12 (1993).
- [2] C. Gundlach, preprint gr-qc/9712084.
- [3] M. W. Choptuik, T. Chmaj, and P. Bizoń, Phys. Rev. Lett. 77, 424-427 (1996).
- [4] P. R. Brady, C. M. Chambers, and S. M. C. V. Gonçalves, Phys. Rev. D56, 6057-6061 (1997).
- [5] P. Bizoń, T. Chmaj, and Z. Tabor, in preparation.
- [6] P. Bizoń and T. Chmaj, Phys. Lett. B 297, 55-61 (1992).
- [7] M. Heusler, S. Droz, and N. Straumann, Phys. Lett. B 271, 61-67 (1991).
- [8] M. Heusler, N. Straumann, and Z. Zhou, Helv. Phys. Acta 66, 615-631 (1993).