DISCRETE AND FINITE GENERAL RELATIVITY

Manoelito M de Souza and Robson N. Silveira

Universidade Federal do Espírito Santo - Departamento de Física

29065.900 - Vitória-ES-Brasil

(December 2, 2024)

We develop the General Theory of Relativity in a formalism with extended causality that describes physical interaction through discrete, transversal and localized pointlike fields. The homogeneous field equations are then solved for a finite, singularity-free, point-like field that we associate to a "classical graviton". The standard Einstein's continuous formalism is retrieved by means of an averaging process, and its continuous solutions are determined by the chosen imposed symetry. The Schwarzschild metric is obtained by the imposition of spherical symmetry on the averaged field.

PACS numbers: $04.20.Cv \quad 04.30. + x \quad 04.60. + n$

I. INTRODUCTION

If we can assume that the gravitational interaction between two masses, as any other elementary interaction, is fundamentally of a quantum nature, that is, mediated by a discrete and localized agent (the graviton), then we must conclude that the General Theory of Relativity (GTR) is a wonderful average geometric description of such phenomenon: it replaces this intermediating agent by the metric of a curved spacetime, smoothing and hiding the discreteness, the localized and anisotropic aspects of the quantum interaction; it is a wonderful description because it fits [1] our observational data, what can be seen then as a consequence of the minuteness of the action of a single graviton.

If this can be accepted as a possibly correct description then GTR just describes an effective average continuous interaction; then it would not be correct to try to reobtain the fundamental quantum picture from this effective description, and besides, this would be an impossible task as they have support on distinct and not-compatible topologies. The concept of a corpuscle (a quantum) is associated to the geometry of a line (its support manifold, free of singularity) while the field interaction of GTR is associated to the geometry of a lightcone, which is singular at its vertex. This may explain the singularities of GTR and the difficulties with its quantization [2,3].

Another procedure [4], that in our opinion is the most appropriate would be, back to the origins, start with a classical description that contemplates the discrete, localized and anisotropic nature of the fundamental interactions, but that reproduces GTR, as a continuous and distributed-field theory, upon an averaging procedure. This, actually, should apply for all standard field theories. We have obtained very positive results with a such approach on Electrodynamics [5] and now we describe our first similar experiences with General Relativity.

In Section II, for the sake of completeness, we reproduce a brief review [4,5] on extended causality and its applications to field theory. In Section III we show how the discrete fundamental field can be seen as an elementary part of the standard continuous field. The continuous field is not necessary for a definition of the discrete field; this is just an heuristic view. The theory must be defined in terms of the discrete field; the continuous field and its standard formalism are retrieved in terms of effective averages of the discrete field. The General Theory of Relativity is described in terms of discrete fields in Section IV, and in Section V the homogeneous field equations is solved for a discrete solution. The Schwarzschild metric is recovered with the assumption of spherical symmetry. Finally we conclude in Section VI discussing on its physical meaning and consequences.

II. CAUSALITY IN FIELD THEORY

In standard field theory formalism the evolution of a field is constrained by

$$\Delta \tau^2 = -\Delta x^2,\tag{1}$$

where Δx is a four-vector in a Minkowski spacetime of metric $\eta = diag(1, 1, 1, -1)$. In our notation we omit the spacetime indices when this does not compromise the text comprehension. So, x stands for x^{μ} , ∂ for ∂_{μ} , and $A(x, \tau)$ for a vector field $A^{\mu}(x, \tau)$, for example. τ is a real-valued parameter. So, (1) just expresses that Δx cannot be spacelike. This is the *local causality*, and (1) defines the change of propertime $\Delta \tau$ associated to Δx . Geometrically it is the definition of a three-dimensional double cone; Δx is the four-vector separation between a generic event

 $x^{\mu} \equiv (\vec{x}, t)$ and the cone vertex. See the Figure 1. This conic hypersurface is the support manifold for the field definition: a free field cannot be inside nor outside but only on the cone. The cone-aperture angle θ is given by

$$\tan \theta = \frac{|\Delta \vec{x}|}{|\Delta t|}, \qquad c = 1, \tag{2}$$

or $\Delta \tau^2 = (\Delta t)^2 (1 - \tan^2 \theta)$. A change of the supporting cone corresponds to a change of speed of propagation and is an indication of interaction. Special Relativity restricts θ to the range $0 \le \theta \le \frac{\pi}{4}$, which corresponds to a restriction on $\Delta \tau$:

$$0 \le |\Delta \tau| \le |\Delta t|. \tag{3}$$



Fig. 1. The relation $\Delta \tau^2 = -\Delta x^2$, a causality constraint, is seen as a restriction of access to regions of spacetime. It defines a three-dimension cone which is the spacetime available to a point physical object at the cone vertex.

Eq. (1) is a constraint on a free physical object, a particle or a field, to remain on the cone. The concept of extended causality corresponds to a more restrictive constraint; it requires that (1) be continuous and simultaneously applied to the neighbourhood of x: $(\Delta \tau + d\tau)^2 = -(\Delta x + dx)^2$, which with (1) becomes $\Delta \tau d\tau + \Delta x. dx = 0$. This is equivalent to the imposition of a second constraint, besides the first one (1):

$$d\tau + f.dx = 0. \tag{4}$$

f is a constant spacelike $(f^2 = -1)$ four-vector tangent to the cone, a cone generator, and it is defined by $f^{\mu} = \frac{\Delta x^{\mu}}{\Delta \tau}$, if $\Delta \tau \neq 0$; it is lightlike $(f^2 = 0)$ in the limiting case when $\Delta \tau = 0$.

The equation (4) can be obtained from direct differentiation of (1), and geometrically it defines a hyperplane tangent to the cone (1). We have from (4) that

$$f_{\mu} = -\frac{\partial \tau}{\partial x^{\mu}},$$

when τ is seen as a function of x, a solution of (1): $\tau = \tau_0 \pm \sqrt{-(\Delta x)^2}$. For $\Delta \tau = 0$, f_{μ} is orthogonal to the hyperplane (4), but it is also a lightcone generator.

Imposing in field theory the two constraints, (1) and (4), instead of just (1), as it is usually done, corresponds to knowing the initial position and velocity in point-particle dynamics.

Together, the constraints (1), in the form $d\tau^2 + dx^2 = 0$, and (4) are equivalent to the single condition $(dx)^2 + (f.dx)^2 = 0$, that may be put as

$$dx.\Lambda^f.dx = 0, (5)$$

with $\Lambda^f_{\mu\nu} = \eta_{\mu\nu} + f_{\mu}f_{\nu}$, $(f_{\mu} = \eta_{\mu\nu}f^{\nu})$, which is a projector orthogonal to f^{μ} , $f.\Lambda.f = 0$. Therefore the constraint (5) allows only displacements dx^{μ} parallel to f^{μ} . A fixed four-vector f at a point represents a fiber in the spacetime, a straight line tangent to f^{μ} , the f-generator of a local cone (1). For a massless field, (1) defines a lightcone, (4) defines a hyperplane tangent to this lightcone, and f^{μ} is the lightcone generator tangent to this hyperplane.

One can summarise it by saying that while the local causality restricts the available space-time of a free physical object to a conic three-dimensional hypersurface, the extended causality restricts it to just a straight line, a cone

III. FIELDS AND FIELD EQUATIONS

 $A_f(x)$ is a f-field, that is, a field defined on a fiber f. It is distinct of the field A(x) of the standard formalism, which is defined on the cone. $A_f(x)$ may be seen as the restriction of A(x) to a fiber f, i.e.

$$A(x,\tau)_f = A(x,\tau) \Big|_{dx.\Lambda^f.dx=0} \tag{6}$$

It is a point-like field, the intersection of the wave-front $A(x,\tau)$ with the fiber f. See the Figure 2. This definition (6) would not make any sense if the point character (discrete and localized) of A_f could not be sustained during its time evolution governed by its wave equation. It is remarkable that it remains as a pointlike field [5] as it propagates. Conversely, we have that

$$A(x,\tau) = \frac{1}{4\pi} \int d^2 \Omega_f A_f(x,\tau),\tag{7}$$

where the integral represents the sum over all f directions on the cone (1). 4π is a normalization factor, $4\pi = \int d^2\Omega_f$. The physical interpretation associates $A_f(x)$, a point-perturbation propagating along the lightcone generator f, with a physical photon - we call it a classical photon - and A(x), the standard continuous field, to the effect of the classical photon smeared on the lightcone spacetime. It is worthwhile to underline here a remark about the physical distinction between $A_f(x,\tau)$ and $A(x,\tau)$. They do not represent equivalent physical descriptions. $A_f(x,\tau)$ corresponds to a single real physical photon with f being its four-vector velocity and with transverse electromagnetic fields, while $A(x,\tau)$, due to the smearing process (7), corresponds to a continuous distribution of fictitious unphysical photons with longitudinal electromagnetic field. This explains all the relative difficulties we have on quantizing the Maxwell field $A(x,\tau)$. Another remarkable distinction is that $A_f(x,\tau)$ is a finite pointwise field while $A(x,\tau)$ has a singularity introduced [5] by the smearing process (7). Figure 2 shows the relationship between the fields A_f and A for a process involving the emission of a single physical photon $A_f(x)$; A here is its space average.

The derivatives of $A_f(x)$, allowed by the constraint (5), are the directional derivatives along f, which with the use of (4) or of (5) we write as

$$\partial_{\mu}A_{f} = \left(\frac{\partial}{\partial x^{\mu}} + \frac{\partial\tau}{\partial x^{\mu}}\frac{\partial}{\partial\tau}\right)A(x,\tau)\Big|_{dx,\Lambda^{f},dx=0} = \left(\frac{\partial}{\partial x^{\mu}} - f_{\mu}\frac{\partial}{\partial\tau}\right)A_{f} \equiv \nabla_{\mu}A_{f}.$$
(8)

With ∇ replacing ∂ for taking care of the constraint (5), the propertime τ is treated as a fifth independent coordinate.

Fig. 2. The relationship between the fields A_f and A. The three doted circles represent, at three instants of time, the field A as an spherically symmetric signal emitted by a charge at the point P. The straight line PQRS... is the fiber f, a light-cone generator tangent to f^{μ} . The points Q, R, and S are a classical photon A_f at three instants of time.

The field equation for a massless field defined on a lightcone generator f is, consequently,

$$\eta^{\mu\nu}\nabla_{\mu}\nabla_{\nu}A_f(x,\tau) = J(x,\tau),\tag{9}$$

or, explicitly

$$(\eta^{\mu\nu}\partial_{\mu}\partial_{\nu} - 2f^{\mu}\partial_{\mu}\partial_{\tau})A_f(x,\tau) = J(x,\tau), \tag{10}$$

as $f^2 = 0$. J is its source four-vector current.

An integration over the f degrees of freedom in (9) reproduces with the use of (7) the usual wave equation of the



$$\eta^{\mu\nu}\partial_{\mu}\partial_{\nu}A(x) = J(x),\tag{11}$$

as $\int d^2 \Omega_f f^{\mu} \partial_{\mu} \partial_{\tau} A_f(x) = 0$ because, as can be shown [4], $A_f(x) = A_{-f}(x)$. So, the standard formalism is retrieved from this *f*-formalism with the A(x) as the average of $A_f(x)$, in the sense of (7).

IV. GENERAL RELATIVITY WITH EXTENDED CAUSALITY.

We want to apply here this theory for the gravitational field in the Einstein's General Relativity with

$$g^f_{\mu\nu} = g_{\mu\nu} \Big|_{dx.\Lambda_f.dx=0}$$

The simplest way is just to write

$$g^{f}_{\alpha\beta} = \eta_{\alpha\beta} + H_{f}(x,\tau)f_{\alpha}f_{\beta} \tag{12}$$

where $H_f(x,\tau)$ represents a local spacetime deformation produced by the presence of a single graviton propagating along a straight line tangent to f, $f^2 = 0$. The four vector parameter f, we remind, is a constant, which is an expression of the graviton freedom that freely propagates up to the point where it suffers an interaction (it is absorbed). In extended causality all interactions are discrete and localized in a point and there is no place for self-interactions. This is just a consequence of f being constant! From $g_f^{\alpha\beta}g_{\beta\mu}^f = \delta_{\mu}^{\alpha}$ we have $g_f^{\alpha\beta} = \eta^{\alpha\beta} - H_f(x,\tau) \eta^{\alpha\mu}f_{\mu} \eta^{\beta\nu}f_{\nu}$. As $f^{\mu} =: g_f^{\mu\nu}f_{\nu} = \eta^{\mu\nu}f_{\nu}$, because of $f^2 = 0$, we can write $g_f^{\alpha\beta} = \eta^{\alpha\beta} - H_f(x,\tau)f^{\alpha}f^{\beta}$. Observe that $g_f^{\alpha\beta}$ and $g_{\alpha\beta}^f$ are both bi-linear on f. This enormous simplification -the absence of non-linearity- is exclusively a consequence of (12) and of f being lightlike. They are justified with the classical vision of $g_f - \eta$ as a point-field describing a graviton, freely propagating with the velocity of light. It is important to remark that (12) does not imply any kind of weak field approximation: $H_f(x,\tau)f_{\alpha}f_{\beta}$ is equal to $g_{\alpha\beta}^f - \eta_{\alpha\beta}$ whichever be $g_{\alpha\beta}^f$. An immediate consequence of $f^2 = 0$ is that $\det g_{\alpha\beta}^f = \det \eta_{\alpha\beta}^f = -1$, which indicates that (12) describes a singularity-free field. Let us write $(\Gamma_{\beta\gamma}^{\lambda})^f$ as the Christoffel symbols restricted to the line f, defined by $(\Gamma_{\beta\gamma}^{\lambda})^f = \frac{1}{2}g_f^{\lambda\sigma}(\nabla_{\gamma}g_{\beta\sigma}^f + \nabla_{\beta}g_{\sigma\gamma}^f - \nabla_{\sigma}g_{\beta\gamma}^f)$. For notation simplicity we will write $(\Gamma_{\beta\gamma}^{\lambda})^f$ as just $\Gamma_{\beta\gamma}^{\lambda}$ without the index f, and $\nabla_{\alpha}H_f \equiv H_{\alpha}$ and $\nabla_{\alpha}\nabla_{\beta}H_f \equiv H_{\alpha\beta}$. Then,

$$2\Gamma^{\mu}_{\alpha\beta} = f_{\alpha}f^{\mu}H_{\beta} + f^{\mu}f_{\beta}H_{\alpha} - f_{\beta}f_{\alpha}\eta^{\mu\nu}H_{\nu} + f_{\alpha}f_{\beta}Hf.H, \qquad (13)$$

and $2\Gamma^{\lambda} \equiv 2g_{f}^{\beta\gamma}\Gamma^{\lambda}_{\beta\gamma} = f^{\lambda}f^{\alpha}H_{\alpha} \equiv f^{\lambda}f.H$ So, the harmonic coordinate conditions ($\Gamma^{\lambda} = 0$) imply on

$$f.H = 0. \tag{14}$$

The physical meaning of this coordinate condition is best exposed in the case of solutions to the inhomogeneous field equations, which will be discussed in a subsequent paper, where it reveals to be exactly like the gauge condition [5] of discrete electrodynamics, a constraint between the direction of emission (absorption) of a graviton and the consequent changes in the state of motion of its source (sink).

As a consequence we have also $\Gamma^{\lambda}_{\beta\lambda} = 0$ and

$$R^{f}_{\rho\alpha\beta\sigma} = \frac{1}{2} (f_{\rho}f_{\beta}H_{\sigma\alpha} - f_{\alpha}f_{\beta}H_{\sigma\rho} - f_{\rho}f_{\sigma}H_{\beta\alpha} + f_{\alpha}f_{\sigma}H_{\beta\rho}), \tag{15}$$

$$R^{f}_{\rho\beta} = \frac{1}{2} f_{\rho} f_{\beta} \eta^{\alpha\sigma} H_{\sigma\alpha} \equiv \frac{1}{2} f_{\rho} f_{\beta} \Box_{f} H, \qquad (16)$$

and

$$R^f = g_f^{\rho\beta} R^f_{\rho\beta} \equiv 0. \tag{17}$$

For reasons of clarity and simplicity we will consider here, on this first work on this subject, just solutions to the homogeneous Einstein field equations, which, with (16) and (17) are then reduced to

$$\Box H(m-) = 0 \tag{19}$$

A very simple equation indeed, a consequence of (12) and of $f^2 = 0$. The light-like f in (12) eliminates all the intrinsic non-linearity of General Relativity. But one should be warned that only the inhomogeneous equations are really meaningful in extended causality because the physical properties of the emitted (absorbed) field reflects the changes its emission (absorption) caused on the state of motion of its source (sink). The changes in the sources provide valuable informations about the field, like its angular momentum and its state of polarization. We will not discuss any further these shortcomings as they are just consequences of a solution to an (homogeneous) equation without a source term. They are not present when, in a subsequent work, we consider a solution to the inhomogeneous equations. This extremely simple, information depleted system is, nonetheless, reach enough to justify its presentation as a first introduction to the subject. It will enlighten the physical significance of continuous solutions of the standard formalism, like the Schwarzschild metric for example.

V. DISCRETE SOLUTION

The most general solution to this equation can be obtained, for example, from a Fourier expansion

$$H(x,\tau) = \int d^5 p H(p) \ e^{i(p_{\mu}x^{\mu} + p_5\tau)},\tag{19}$$

with x and τ treated as five independent variables. The simplest solution to (18) and (19) with $f^2 = 0$, is $H(p) = \delta[(p_{\mu} - f_{\mu}p_5)^2]$ or, for mathematical convenience, $H(p) = \chi \frac{|p \cdot f|}{p \cdot f} \delta[(p_{\mu} - f_{\mu}p_5)^2] = \frac{\chi}{2p \cdot f} \delta(p_5 - \frac{p^2}{2p \cdot f})$, where χ is a constant. Then we have that

$$H(x,\tau) = \chi \int d^4p \frac{e^{i(p_\mu x^\mu + \frac{p^2}{2p.f}\tau)}}{2p.f}.$$
(20)

It is crucial in this expression that one has p.f in the integrand denominator instead of the p^2 that one would have in the usual local-causality formalism, which would give origin to a $\frac{1}{r}$ -dependence and, therefore, a metric with a singularity on r = 0. The extended causality with its anisotropy determined by the existence of a graviton allows the replacement of p^2 by p.f. This radically changes the nature and characteristics of the theory.

We observe that the integrand of (20) has a singularity at $p \cdot f = 0$ and that the exponent in the integrand is also only defined at this point if

$$p^2\Big|_{p.f=0} = 0.$$
(21)

This implies on a system of two simultaneous equations

$$p^2 = p_{\rm T}^2 + p_{\rm L}^2 - p_4^2 = 0, \tag{22}$$

and

$$p.f = p_{\rm L} |\vec{f}| - p_4 f_4 = 0, \tag{23}$$

where the subindices L and T stand, respectively, for longitudinal and transversal with respect to the space part \hat{f} of f.

As
$$f^2 = 0$$
 we may write $\epsilon = \frac{|f|}{f_4} = \pm 1$, so that (23) becomes $p_4 = \epsilon p_L$. Equation (22) is, consequently, equivalent to

$$p_{\rm T} = 0. \tag{24}$$

The conditions (22) and (24) are full of physical significance: the first one requires a massless field and the second one implies that $\Delta x_{\rm T} = 0$; only the $x_{\rm L}$, that is the longitudinal coordinate, participates in the system evolution [4]. So, we can see, in anticipation, that the field H_f only propagates along the fiber f. The equation (20), as a consequence of (24), is reduced to

$$H_f(x,\tau) = \chi \int \frac{dp_{\rm L} dp_4}{2f_4(p_4 - \epsilon p_{\rm L})} e^{i(p_{\rm L}x_{\rm L} + p_4t + \frac{p_4 + \epsilon p_{\rm L}}{2f_4}\tau)},\tag{25}$$

$$\frac{p_{\rm L}^2 - (p_4)^2}{2p.f} = \frac{(\epsilon p_{\rm L} + p_4)(\epsilon p_{\rm L} - p_4)}{2(\epsilon p_{\rm L} - p_4)f_4} = \frac{\epsilon p_{\rm L} + p_4}{2f_4}.$$
(26)

We now make explicit the integration on the coordinate p_4 ,

$$H_f(x,\tau) = \lim_{\varepsilon \to 0} \chi \int dp_{\rm L} dp_4 \frac{e^{i(p_{\rm L}x_{\rm L} + p_4t + \frac{p_4 + \epsilon p_{\rm L}}{2f_4}\tau)}}{2f_4(p_4 - \epsilon p_{\rm L} \pm i\varepsilon)},\tag{27}$$

which produces

$$H(x,\tau)_f = 2\pi i a \chi \theta[a(t+\frac{\tau}{2f_4})] \int \frac{dp_{\rm L}}{2f_4} e^{ip_{\rm L}(x_{\rm L}+\epsilon t+\frac{\epsilon}{f_4}\tau)},\tag{28}$$

where a stands for ± 1 , a sign that comes from the choice of the contour in a Cauchy integral, i.e. the sign of $\pm i\varepsilon$. The signs of $a = \pm 1$ are connected [4] respectively to the creation and annihilation of $g_f(x, \tau)$; and $\theta(t)$ is the step function ($\theta(t \ge 0) = 1$; $\theta(t < 0) = 0$). On the other hand

$$\frac{1}{2\pi} \int \frac{dp_{\rm L}}{f_4} e^{ip_{\rm L}(x_{\rm L}+\epsilon t+\frac{\epsilon\tau}{f_4})} = \frac{1}{2\pi\epsilon} \int \frac{\epsilon dp_{\rm L}}{f_4} e^{i\frac{\epsilon p_{\rm L}}{f_4}(f_{\rm L}x_{\rm L}+f_4t+\tau)} = \frac{1}{\epsilon} \delta(\tau+f.x), \tag{29}$$

and therefore, we have for (28)

$$H(x,\tau)_f = \frac{ia(2\pi)^2}{\epsilon} \chi \theta[a(t+\frac{\tau}{2f_4})]\delta(\tau+f.x).$$
(30)

So, after some simple algebra and a redefinition of the constant χ , we may write

$$H(x,\tau)_f = \chi \theta(at) \delta(\tau + f \cdot x). \tag{31}$$

Therefore, with $\tau = 0$ accounting for the massless field, and t > 0, because we are considering only the emitted field (a = +1) and not the absorbed one, the metric (12) becomes

$$g_{\alpha\beta}^{\ f} = \eta_{\alpha\beta} + \chi \delta(f \cdot \Delta x) f_{\alpha} f_{\beta}. \tag{32}$$

In spherical coordinates we have

$$g^{f}_{\alpha\beta}(t,r,\theta,\varphi) = \begin{cases} \eta_{\alpha\beta}, & \text{for } \theta \neq \theta_{f}, \ \varphi \neq \varphi_{f}; \\ \eta_{\alpha\beta} + Hf_{\alpha}f_{\beta} & \text{for } \theta = \theta_{f}, \ \varphi = \varphi_{f}, \end{cases}$$
(33)

or explicitly

$$g_{\alpha\beta}^{f}(t,r,\theta,\varphi) = \begin{pmatrix} -1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & r^{2} & 0\\ 0 & 0 & 0 & r^{2}sen^{2}\theta \end{pmatrix} + \chi\delta(f.\Delta x) \begin{pmatrix} f_{0}^{2} & f_{0}f_{r} & 0 & 0\\ f_{r}f_{0} & f_{r}^{2} & 0 & 0\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \text{for} \quad \theta = \theta_{f}, \ \varphi = \varphi_{f}$$
(34)

with $f_{\mu} = (f_0, f_r, 0, 0)$, and with θ_f and φ_f , defining the space direction \vec{f} of f. The metric $g^f_{\alpha\beta}$ represents a single, let's say, "classical quantum" of gravity propagating along a line f and observed as an event $(t, r, \theta_f, \varphi_f)$ at the probe mass. Let us, in an abuse of language, call it the graviton on the fiber f, for shortness.

VI. RETRIEVING THE SCHWARZSCHILD FIELD

The presence of a "graviton" on the fiber f breaks the otherwise spherical symmetry in (33,34). It is not, of course, a static solution. There is no static solution in an extended causality formalism. As we will see, the observed (electromagnetic and gravitational) static fields are just average fields, apparently static as a consequence of the large number of quanta exchanged and of the inertial limitations of our measuring apparatus. From this discrete, localized and singularity-free solution $g^f_{\alpha\beta}$ on the lightcone-generator f we can recover the standard continuous and distributed solutions $a_{\alpha\beta} a$ with just an integration over the f-parameter. Like in the case of the electromagnetic field, a continuous

(non-physical) gravitons $g_{\mu\nu}^{f'}$, each one still propagating on the same fiber f. What distinguishes the physical graviton from the fictitious ones is that only f corresponds to the field four-velocity, and so only \vec{f} is collinear to the \vec{x} , the direction of propagation of the gravitons. This is the exact analogous to what happens in the discretization of the electromagnetic field [5]. The continuous solution so obtained is determined by its chosen symmetry. Let us chose an spherically symmetric distribution of $g_{\mu\nu}^{f'}$.

Then, from

$$g_{\alpha\beta}(x) = \frac{1}{4\pi} \int d^2 \Omega_{f'} \ g_{\alpha\beta}^{f'}(x), \tag{35}$$

and with $-f_0 = f_r = 1$, we have for $\theta = \theta_f$, $\varphi = \varphi_f$

$$g_{\alpha\beta} = \eta_{\alpha\beta} + \frac{\chi}{2\pi} \begin{pmatrix} 1 & -1 & 0 & 0\\ -1 & 1 & 0 & 0\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0 \end{pmatrix} \int d^2 \Omega_{f'} \ \delta(f' \cdot x).$$
(36)

Writing $f'.x = f'^0t - \vec{f'}.\vec{r} = f'^0(t - r\cos\theta_{f'})$, the integration $\int d^2\Omega'_f \,\delta(f'\cdot x)$ may be written as

$$\int d\varphi'_f \, sen\theta'_f \, d\theta'_f \, \delta(t - |\vec{r}| \cos\theta'_f) = \frac{2\pi}{r} \int_{-1}^1 d\cos\theta'_f \, \delta(\cos\theta'_f - \frac{t}{r}) = \begin{cases} \frac{2\pi}{r}, & \text{for } t \in [0, r];\\ 0, & \text{for } t \notin [0, r]. \end{cases}$$
(37)

This condition on t means that the deformation on the flat spacetime that we are associating to a graviton is not null as far as t is smaller, or at least, equal to $\frac{r}{c}$, (the time that the graviton, after being emitted by the source at the origin, takes to reach the probe mass at (t, r, θ_f, ϕ_f) , where it is absorbed. This process is continued for $t > \frac{r}{c}$ by other gravitons subsequently emitted [6]. So, the large number of gravitons emitted (and absorbed) in any realistic experiment transmit the idea of continuity and of a static field.

Then we can write for $\theta = \theta_f$ and $\varphi = \varphi_f$ that

$$ds^{2}(t, r, \theta_{f}, \varphi_{f}) = -(1 - \frac{\chi}{r}) dt^{2} + (1 + \frac{\chi}{r}) dr^{2} - \frac{2\chi}{r} dt dr + r^{2} d^{2}\Omega,$$
(38)

where $d^2\Omega = d\theta^2 + sen^2\theta \, d\varphi^2$, which actually is null for fixed θ and φ . A well-known [7] simple coordinate transformation puts (38) in the standard diagonal form

$$ds^{2}(\theta_{f},\varphi_{f}) = -(1-\frac{\chi}{r})dt^{2} + (1-\frac{\chi}{r})^{-1}dr^{2} + r^{2}d^{2}\Omega,$$
(39)

and

$$ds^{2}(\theta,\varphi) = -dt^{2} + dr^{2} + r^{2}d^{2}\Omega, \quad \text{for} \quad \theta \neq \theta_{f}, \ \varphi \neq \varphi_{f}.$$

$$\tag{40}$$

We recognize (39) as the Schwarzschild metric. It is a consequence of the assumed isotropy in the distribution of ficticious gravitons; other distinct symmetries, of course, generate other distinct metrics. The probe mass detects the Schwarzschild field, although the spacetime remains flat outside the segment on the line \vec{f} that connects the two interacting points (the probe mass and the source on the origin). This is consistent with the extended causality concept: there is no field where there is no interaction! See the Figure 3.

Fig. 3. A probe mass m, wherever be it placed, detects a Schwarzschild metric on the spacetime around the mass M on the origin. The observer concludes then that the M is the symmetry center of a Schwarzschild spacetime, but actually the spacetime is flat except on the straight-line segment connecting m to M; it is completely flat, except at the origin (M), in the absence of m.

In General Relativity, like in Electrodynamics [5], the standard continuous field can be retrieved from the discrete one



these unphysical fields are the responsible for the complications on an otherwise simple quantization process; one may assume that in General Relativity they make this quantization impossible. In both theories singularities are just consequences of the averaging process, of using these averages as if they were the actual fundamental fields.

VII. CONCLUSIONS

We have shown, in previous works [2,3], that the problems of both classical and quantum field theories with singularities, divergencies, and difficulties of quantization (for the classical ones) are consequences of being defined with support on the lightcone; then their fields are not the real fundamental ones but just their effective averages. The actually fundamental fields must be defined with support on the lightcone generators. This corresponds to adopting extended, instead of local, causality. The Maxwell theory of electromagnetism has been shown [5] to be free of these problems when formulated on the lightcone generator, that is, in terms of finite and discrete pointlike fields (photons). The standard formalism with its all known problems is recuperated when the photon fields are replaced by continuous fields defined by the photons effective averages on the lightcone. An important message then is that electromagnetic field singularities are not real physical objects but just artificial consequences of using an inappropriate formalism. The remarkable in the present work is that all these considerations on the electromagnetic field are now repeated for the gravitational field of the General Theory of Relativity. Although these two fields have many similarities, as they are both massless gauge fields, their respective theories have many disparing properties as the non-linearity of the theory of gravity and its geometric association to the space-time metric.

We have shown here that the Schwarzschild metric, a static spherically symmetric field, can be seen as the average effect of the flux of discrete pointlike fields exchanged between two point masses. Its time independence and its singularity, a really not physical object, are consequences of taking an averaging by the fundamental field. And more, there is field only where and when there is a probe mass! A mass alone (as any single source) has no field around it! This is quite a change and, certainly, of no easy acceptation as it goes against the prevalent trend of seeing the field singularities as real physical objects, and the continuous metric field as a true physical representation of the world geometry (not just an approximation), nonetheless the unsourmountable difficulties that this implies on having a gravity quantum theory.

Acknowledgements

R. N. Silveira acknowledges his grant from CAPES for writing his M.Sc. dissertation. The authors thank Marcelo Sarandy for a critical reading of the manuscript.

- [1] C.M. Will, Int. J. Mod. Phys. D 1, 13(1992).
- [2] M. M. de Souza, "Classical Electrodynamics and the Quantum Nature of Light. J. of Phys. A: Math. Gen. 30 (1997)6565-6585.
- [3] M. M. de Souza, "Dynamics and causality constraint on the lightcone." hep-th/9708096. To be published.
- [4] M. M. de Souza, "Discrete Fields on the Lightcone." hep-th/9708066. To be published.
- [5] M. M. de Souza, J.V. Costa, A.S. Pedra, "Finite Discrete Electromagnetic Field Theory." hep-th/9712069. To be published.
- [6] M. M. de Souza, "Classical Fields and the Quantum concept." hep-th/9610145.
- [7] S. Weinberg, Gravitation and Cosmology, p. 176. John Wiley, New York, 1971.