COSMIC STRING CREATED FROM VACUUM FLUCTUATION

Arkadii Popov *
Department of Geometry, Kazan State Pedagogical University,
Mezhlauk str., 1, Kazan 420021, RUSSIA

Abstract

The possibility of the cosmic string creation by the vacuum fluctuations of quantum fields in the self-consistent semiclassical theory of gravity is discussed. We use the approximate method for obtaining vacuum expectation value of the renormalized stress-energy tensor of conformally invariant quantum fields in static cylindrically symmetric spacetimes. We have obtained the particular solutions of Einstein equations for the different boundary conditions at the cylinder symmetry axis.

There is very interesting question: can the vacuum fluctuation of quantum fields to be the single source of curvature or nontrivial topological structure of spacetime? One may find the solution of this problem in the theory of quantum gravity. But a fully satisfactory theory of quantum gravity does not yet exist. As a first step on the way of solution of this problem one may to use the self-consistent semiclassical theory of gravity. This theory sets the classical Einstein tensor equal to the expectation value of the stress-energy tensor operator of the quantized matter fields present (throughout we use units such that $c = \hbar = G = 1$)

$$G_{\mu\nu} = 8\pi \langle T_{\mu\nu} \rangle. \tag{1}$$

The primary difficulty in the theory of semiclassical gravity is that the effects of the quantized gravitational field are ignored. A secondary difficulty of semiclassical gravity is that $\langle T_{\mu\nu} \rangle$ depends strongly on the metric and is generally difficult to calculate. Most calculations of $\langle T_{\mu\nu} \rangle$ have been made on fixed classical background. These results can not be used in selfconsistent semiclassical theory. There have been approximate calculations made for conformally coupled fields in any static spherically symmetric Einstein spacetime [1, 2]. The analytical approximations to $\langle T_{\mu\nu} \rangle$ have been obtained by Anderson, Hiscock and Samuel [3] for massless or massive scalar fields with an arbitrary coupling ξ to scalar curvature in static spherically symmetric spacetimes. In this work we use the Killing ansatz giving approximate vacuum expectation value of the stress-energy tensor $\langle T_{\mu\nu} \rangle$ for conformally coupled massless field in static spacetimes which was obtained by Frolov and Zelnikov [4]. The Killing ansatz is constructing as an approximate expression for the vacuum expectation value $\langle T_{\mu\nu} \rangle$ using the Riemann tensor, the Killing vector and their covariant derivatives. The tensor $\langle T_{\mu\nu} \rangle$ is constructed so that the conservation law $\langle T^{\nu}_{\mu} \rangle_{;\nu} = 0$ and the conformal trace anomaly are fulfilled. In addition a correct scaling transformation behaviour of $\langle T_{\mu\nu} \rangle$ is necessary. Then the following expression can be obtained for $\langle T_{\mu\nu} \rangle$:

$$\langle T_{\mu\nu}\rangle = \langle T_{\mu\nu}(g_{\mu\nu}, \xi^{\mu}, q_1, q_2)\rangle, \tag{2}$$

^{*}E-mail: popov@kspu.ksu.ras.ru

where $g_{\mu\nu}$ is the metric tensor, ξ^{μ} is the Killing vector of static spacetime, q_1 and q_2 are the arbitrary constants of ansatz.

The metric for the general static cylindrically symmetric spacetime is

$$ds^{2} = -e^{2F}dt^{2} + b^{2}dz^{2} + dr^{2} + c^{2}d\varphi^{2},$$
(3)

where F, b and c are functions of the proper distance r.

We restrict our consideration to the special case

$$F = const. (4)$$

From this assumption it is easy to obtained the following expression for $\langle T_{\mu\nu} \rangle$ [5]:

$$T_{\mu\nu} = 2\left[\alpha(F - F_0) + q_1\right] \left[\frac{2}{3}R_{;\mu\nu} - 2R_{\mu\nu;\sigma}^{;\sigma} - \frac{14}{3}RR_{\mu\nu} + 8R_{\mu\sigma}R_{\nu}^{\sigma} + g_{\mu\nu}\left(\frac{1}{3}R_{;\sigma}^{;\sigma} - 3R_{\alpha\beta}R^{\alpha\beta} + \frac{5}{3}R^2\right) + \xi_{\mu\nu}\left(4R_{\alpha\beta}R^{\alpha\beta} - 2R^2\right)\right] + \alpha\xi_{\mu\nu}\left(\frac{2}{3}R_{;\sigma}^{;\sigma} + 2R_{\alpha\beta}R^{\alpha\beta} - \frac{2}{3}R^2\right) + q_2\left[\frac{1}{3}R_{;\mu\nu} - RR_{\mu\nu} + g_{\mu\nu}\left(-\frac{1}{3}R_{;\sigma}^{;\sigma} + \frac{1}{12}R^2\right) + \xi_{\mu\nu}R_{;\sigma}^{;\sigma}\right],$$
 (5)

where the notations are used:

$$\xi^{\mu}(e^{-F_0}, 0, 0, 0), \quad \xi^2 = \xi^{\sigma}\xi_{\sigma}, \quad \xi_{\mu\nu} = \xi_{\mu}\xi_{\nu}/\xi^2,$$
 (6)

$$\alpha = \frac{1}{2^9 \cdot 45\pi^2} \left[12h(0) + 18h\left(\frac{1}{2}\right) + 72h(1) \right],\tag{7}$$

 $h\left(s\right)$ is the number of helisity states for the conformal massless field of a spin s.

Four nontrivial Einstein equations for this case are given in Appendix. This equation set are compatible only in the case

$$q_2 = 0, \quad 2[\alpha(F - F_0) + q_1] = -\alpha$$
 (8)

and independent equations of this set can be written in form

$$\frac{b''}{2b} + \frac{c''}{c} + \frac{b'c'}{2bc} = 8\pi\alpha \left(-\frac{b''''}{b} + \frac{5b'''b'}{3b^2} - 2\frac{b'''c'}{bc} - \frac{c'''b'}{3bc} + \frac{5b''^2}{3b^2} - \frac{4b''c''}{3bc} \right) - \frac{5b''b'^2}{3b^3} + \frac{b''b'c'}{b^2c} + \frac{b''c'^2}{bc^2} - \frac{c''b'^2}{3b^2c} + 2\frac{c''b'c'}{bc^2} + \frac{b'^3c}{3b^3c} - \frac{b'c'^3}{bc^3} \right) \tag{9}$$

$$\frac{b''}{b} + \frac{c''}{2c} + \frac{b'c'}{2bc} = 8\pi\alpha \left(-\frac{c''''}{c} - \frac{b'''c'}{3bc} - 2\frac{c'''b'}{bc} + \frac{5c'''c'}{3c^2} - \frac{4b''c''}{3bc} + \frac{5c''^2}{3c^2} + \frac{5c''^2}{3c^2} + \frac{5c'''c'}{3bc} - \frac{5c''c'^2}{3bc} - \frac{5c''c'^2}{3c^2} - \frac{b'b'c'}{bc^3} \right) \tag{10}$$

The solution without source at the symmetry axis must satisfy the boundary conditions:

$$b(r) = b_0 + \frac{b_0'''}{2!}r^2 + \frac{b_0''''}{4!}r^4 + 0(r^5) \quad \text{for } r \to 0,$$
 (11)


$$c(r) = c_0' r + \frac{c_0'''}{3!} r^3 + 0(r^5) \quad \text{for } r \to 0,$$
 (12)


where

$$c_0' = 1, \ b_0'''' = b_0 \left[\frac{b_0''^2}{b_0^2} - \frac{3}{8(8\pi\alpha)} \left(\frac{b_0''}{b_0} + \frac{c_0'''}{c_0'} \right) \right], \tag{13}$$

 b_0, b_0'', c_0''' are arbitrary constants. These expressions can be obtained from equations (32) -(35) by substitution the functions b(r) and c(r) power series of r.

The equations (9) and (10) were solved by numerical methods for the scalar field $\left(\alpha = \frac{12}{2^9 \cdot 45\pi^2}\right)$. The results are demonstrated in figures:





Three cases which correspond to different boundary conditions are represented on these figures:

1)
$$c'_0 = 1$$
; $c'''_0 = 2$; $b_0 = 1$; $b''_0 \approx 1.64218$

2)
$$c_0' = 1$$
; $c_0''' = 1$; $b_0 = 1$; $b_0'' \approx 0.69542$

1)
$$c_0' = 1$$
; $c_0''' = 2$; $b_0 = 1$; $b_0'' \approx 1.64218$
2) $c_0' = 1$; $c_0''' = 1$; $b_0 = 1$; $b_0'' \approx 0.69542$
3) $c_0' = 1$; $c_0''' = 0.5$; $b_0 = 1$; $b_0'' \approx 0.29978$

The asymptotic behaviour of b(r) and c(r) can be investigated analytically. From the numerical calculations, we see that for sufficiently large r, both the functions can be represented as a sum of two distinct components

$$b(r) = b_{mon}(r) + \delta b(r); \quad c(r) = c_{mon}(r) + \delta c(r), \tag{14}$$

where $b_{mon}(r)$ and $c_{mon}(r)$ are strictly monotone increasing, $\delta b(r)$ and $\delta c(r)$ are bounded oscillating functions. The magnitudes of these components are connected as follows:

$$\frac{\delta b(r)}{b_{mon}(r)} \sim \frac{\delta c(r)}{c_{mon}(r)} \sim \epsilon \ll 1. \tag{15}$$

Moreover, the magnitudes of the amplitudes and frequencies of $\delta b(r)$ and $\delta c(r)$ are of order unity:

$$\delta b^{(n)}(r) \sim \delta c^{(n)}(r) \sim \epsilon^0, \tag{16}$$

while

$$b_{mon}(r) \sim c_{mon}(r) \sim \epsilon^{-1}. \tag{17}$$

In fact, one supposes that the function $b_{mon}(r)$ and $c_{mon}(r)$ increases like r The numerical calculations also allow us to estimate the magnitudes of the derivatives of the monotonic components:

$$b_{mon}^{(n)}(r) \sim c_{mon}^{(n)}(r) \sim \epsilon^{(n-1)}$$
. (18)

Having these basic estimates we can expand the equations (9,10) keeping the leading $O(\epsilon)$ terms

$$\frac{\delta b''''}{b_{mon}} + \frac{1}{8\pi\alpha} \left(\frac{\delta b''}{2b_{mon}} + \frac{\delta c''}{c_{mon}} \right) = 0, \tag{19}$$

$$\frac{\delta c''''}{c_{mon}} + \frac{1}{8\pi\alpha} \left(\frac{\delta b''}{2b_{mon}} + \frac{\delta c''}{c_{mon}} \right) = 0.$$
 (20)

The solutions of these equations are

$$\frac{\delta b}{b_{mon}} = \frac{D_1}{\omega^2} e^{i\omega r} + \frac{D_2}{\omega^2} e^{-i\omega r} + \frac{D_3}{(\omega^2/3)} e^{(\omega/\sqrt{3})r} + \frac{D_4}{(\omega^2/3)} e^{-(\omega/\sqrt{3})r} + D_5 r + D_6, \tag{21}$$

$$\frac{\delta c}{c_{mon}} = \frac{D_1}{\omega^2} e^{i\omega r} + \frac{D_2}{\omega^2} e^{-i\omega r} - \frac{D_3}{(\omega^2/3)} e^{(\omega/\sqrt{3})r} - \frac{D_4}{(\omega^2/3)} e^{-(\omega/\sqrt{3})r} + D_7 r + D_8, \tag{22}$$

where

$$\omega^2 = \frac{3}{16\pi\alpha} \tag{23}$$

and D_i are arbitrary constants (D_1 and D_2 are complex conjugate, $D_3, D_4, D_5, D_6, D_7, D_8$ are real). Note that the parameters D_i are constants only in the framework of the $O(\epsilon^2)$ approximation. In fact, they may be slowly varying functions of r such that their derivatives are of $O(\epsilon^2)$. Note also that the boundary conditions were chosen so that $D_3 = 0$.

In order to obtain the approximate but analytic expressions for the monotonic components $b_{mon}(r)$ and $c_{mon}(r)$ more work is necessary. For this we must expand equations (32-35) up to order $O(\epsilon^2)$ and substitute the solutions (21, 22) into these equations. Take into account that these equations contain terms of three different types: the terms which consist only of the monotonic functions $b_{mon}(r)$ and $c_{mon}(r)$; the terms which contain one oscillating function $\delta b(r)$ or $\delta c(l)$ (or its derivatives); and the terms which contain a product of two oscillating functions $\delta b(r)$ or $\delta c(l)$ (or its derivatives). After substituting the solutions (21,22) into (32-35) one can

check that the these equations consist of two independent parts: the monotonic one and the oscillating one. Both these parts must be equated to zero separately. Equating the monotonic part to zero we obtain the equations for $b_{mon}(r)$ and $c_{mon}(r)$:

the 00 component

$$\frac{b''_{mon}}{b_{mon}} + \frac{c''_{mon}}{c_{mon}} + \frac{b'_{mon}c'_{mon}}{b_{mon}c_{mon}} = 8\pi\alpha \frac{8}{3}D_1D_2,\tag{24}$$

the zz component

$$\frac{c_{mon}^{\prime\prime}}{c_{mon}} = 0,\tag{25}$$

the rr component

$$\frac{b'_{mon}c'_{mon}}{b_{mon}c_{mon}} = 8\pi\alpha \frac{8}{3}D_1D_2,\tag{26}$$

the $\varphi\varphi$ component

$$\frac{b_{mon}^{"}}{b_{mon}} = 0. (27)$$

And the oscillating part of these equations gives the correction of the order ϵ^2 for $\delta b(r)$ and $\delta c(r)$. The solutions of equations (24-27) are

$$b = k_1 \frac{r}{\sqrt{8\pi\alpha}} + k_2 + 0(\epsilon), \tag{28}$$

$$c = k_3 \frac{r}{\sqrt{8\pi\alpha}} + k_4 + 0(\epsilon),\tag{29}$$

$$D_1 D_2 = \frac{3}{(8\pi\alpha)8r^2} + 0(\epsilon^3),\tag{30}$$

where k_i are constants.

Finally, the asymptotic form of metric is

$$ds^{2} = -dt^{2} + (k_{1} \frac{r}{\sqrt{8\pi\alpha}} + k_{2})^{2} \left[1 + \frac{\sqrt{16\pi\alpha/3}}{r} \sin\left(\frac{r}{\sqrt{16\pi\alpha/3}} + r_{0}\right) \right]^{2} dz^{2}$$

$$+ dr^{2} + (k_{3} \frac{r}{\sqrt{8\pi\alpha}} + k_{4})^{2} \left[1 + \frac{\sqrt{16\pi\alpha/3}}{r} \sin\left(\frac{r}{\sqrt{16\pi\alpha/3}} + r_{0}\right) \right]^{2} d\varphi^{2}, \tag{31}$$
where $\sin(r_{0}) = \frac{ReD_{1}}{\sqrt{D_{1}D_{2}}}, \cos(r_{0}) = \frac{-ReD_{1}}{\sqrt{D_{1}D_{2}}}.$

The asymptotic form of the angular part of these solutions coincides with the angular part of cosmic string. However, in the case of string the conic part of spacetime is 2-dimensional Minkowski spacetime minus a wedge and in the considered cases the conic part of spacetime is 2-dimensional Minkowski spacetime plus a wedge. The value of the cone excess angle is of order unit and it is determined by the boundary conditions at the symmetry axis.

The work was supported by Russian Foundation for Basic Researches grant No. 96-02-17366a

Appendix

The equation $G_0^0 = 8\pi \langle T_0^0 \rangle$ give

$$\frac{b''}{b} + \frac{c''}{c} + \frac{b'c'}{bc} = 8\pi\alpha \left(-\frac{2b''''}{3b} - \frac{2c''''}{3c} + \frac{2b'''b'}{3b^2} - \frac{4b'''c'}{3bc} - \frac{4c'''b'}{3bc} \right)
+ \frac{2c'''c'}{3c^2} + \frac{4b''^2}{3b^2} - 2\frac{b''c''}{bc} + \frac{4c''^2}{3c^2} - \frac{2b''b'^2}{3b^3} + \frac{4b''b'c'}{3b^2c} + \frac{2b''c'^2}{3bc^2} + \frac{2c''b'^2}{3b^2c}
+ \frac{4c''b'c'}{3bc^2} - \frac{2c''c'^2}{3c^3} - \frac{2b'^3c'}{3b^3c} + \frac{2b'^2c'^2}{3b^2c^2} - \frac{2b'c'^3}{3bc^3} \right).$$
(32)

The equation $G_z^z = 8\pi \langle T_z^z \rangle$ give

$$\frac{c''}{c} = 8\pi\alpha \left(-\frac{4b''''}{3b} + \frac{2c''''}{3c} + \frac{8b'''b'}{3b^2} - \frac{8b'''c'}{3bc} + \frac{2c'''b'}{3bc} - \frac{2c'''c'}{3c^2} + 2\frac{b''^2}{b^2} - \frac{2b''c''}{3bc} - \frac{4c''}{3c^2} + \frac{2b'''c'}{3bc} - \frac{2b''c''}{3c^2} + \frac{2b'''c''}{3bc} - \frac{2b''c''}{3bc} - \frac{2b''c''}{3bc} - \frac{2b'''c''}{3bc} - \frac{2b'''c$$

$$-\frac{8b''b'^2}{3b^3} + \frac{2b''b'c'}{3b^2c} + \frac{4b''c'^2}{3bc^2} - \frac{4c''b'^2}{3b^2c} + \frac{8c''b'c'}{3bc^2} + \frac{2c''c'^2}{3c^3} + \frac{4b'^3c'}{3b^3c} - \frac{2b'^2c'^2}{3b^2c^2} - \frac{4b'c'^3}{3bc^3}\right). \tag{33}$$

The equation $G_r^r = 8\pi \langle T_r^r \rangle$ give

$$\frac{b'c'}{bc} = 8\pi\alpha \left(-\frac{4b'''b'}{3b^2} + \frac{2b'''c'}{3bc} + \frac{2c'''b'}{3bc} - \frac{4c'''c'}{3c^2} + \frac{2b'''^2}{3b^2} - \frac{2b''c''}{3bc} + \frac{2c''^2}{3c^2} + \frac{4b''b'^2}{3b^3} \right) - 2\frac{b''b'c'}{b^2c} + \frac{2b''c'^2}{3bc^2} + \frac{2c''b'c'}{3b^2c} - 2\frac{c''b'c'}{3bc^2} + \frac{4c''c'^2}{3c^3} - \frac{2b'^3c}{3b^3c} + \frac{2b'^2c'^2}{3b^2c^2} - \frac{2b'c'^3}{3bc^3} \right).$$
(34)

The equation $G_{\varphi}^{\varphi} = 8\pi \langle T_{\varphi}^{\varphi} \rangle$ give

$$\frac{b''}{b} = 8\pi\alpha \left(\frac{2b''''}{3b} - \frac{4c''''}{3c} - \frac{2b'''b'}{3b^2} + \frac{2b'''c'}{3bc} - \frac{8c'''b'}{3bc} + \frac{8c'''c'}{3c^2} - \frac{4b''^2}{3b^2} - \frac{2b''c''}{3bc} + 2\frac{c''}{c^2}\right)$$

$$+\frac{2b''b'^2}{3b^3} + \frac{8b''b'c'}{3b^2c} - \frac{4b''c'^2}{3bc^2} + \frac{4c''b'^2}{3b^2c} + \frac{2c''b'c'}{3bc^2} - \frac{8c''c'^2}{3c^3} - \frac{4b'^3c'}{3b^3c} - \frac{2b'^2c'^2}{3b^2c^2} + \frac{4b'c'^3}{3bc^3}\right). \tag{35}$$

References

- [1] D.N. Page, Phys.Rev. D **25**, 1499 (1982)
- [2] M.R. Brown, A.C. Ottewill, D.N. Page, Phys.Rev. D 33,2840 (1986)
- [3] P.R. Anderson, W.A. Hiscock, D.A. Samuel, Phys.Rev. D 51, 4237 (1995)
- [4] V.P. Frolov and A.I. Zelnikov, Phys.Rev. D 35, 3031 (1987)
- [5] S.V. Sushkov, Phys. Lett. A 164, 33 (1992)