

Gravitational Analog of the Electromagnetic Poynting Vector

L.M. de Menezes ¹

*Dept. of Physics and Astronomy, University of Victoria, Victoria, B.C. Canada
V8W 3P6*

Abstract

The gravitational analog of the electromagnetic Poynting vector is constructed using the field equations of general relativity in the Hilbert gauge. It is found that when the gravitational Poynting vector is applied to the solution of the linear mass quadrupole oscillator, the correct gravitational quadrupole radiation flux is obtained. Further to this, the Maxwell-like gravitational Poynting vector gives rise to Einstein's quadrupole radiation formula. The gravitational energy-momentum (pseudo) tensor obtained is symmetric and traceless. The former property allows the definition of angular momentum for the free gravitational field.

PACS 04.30.+x

1 Introduction

Despite the rather different frame works from which they arise, the equations of general relativity have many similarities with those of the electromagnetic field. These similarities are seen in the weak field limit of general relativity where the geodesic equations of motion and the field equations have essentially the same structure as the equations of electromagnetic theory. In this work, general relativity and electromagnetism are brought one step closer. It is shown that in the Hilbert gauge, the field equations of general relativity can be generated from a Lagrangian density similar in structure to that of electrodynamics. The structure of the energy-momentum (pseudo) tensor obtained from this Lagrangian density coincides with that of the free electromagnetic field: it is symmetric, traceless and can be constructed from the electromagnetic energy-momentum tensor by replacing the Maxwell tensor $F^{\mu\nu}$ by a suitable antisymmetric gravito-electromagnetic (pseudo) tensor $\Pi^{\mu\nu}$. It is remarkable that the simple Maxwell-like gravitational (pseudo) tensor gives rise

¹ E-mail: demenezes@uvphys.phys.uvic.ca

to Einstein's quadrupole radiation formula. To the author's knowledge this work represents a novel approach to the study of gravitational fields in the weak field limit.

2 Electromagnetism and gravity

The electromagnetic action [1]:

$$S = - \sum \int m \, ds - \int A_\mu J^\mu \, d\Omega - \frac{1}{16\pi} \int F_{\mu\nu} F^{\mu\nu} \, d\Omega \quad (1)$$

gives rise to the first and second pair of Maxwell's equations

$$F_{\mu\nu,\alpha} + F_{\alpha\mu,\nu} + F_{\nu\alpha,\mu} = 0, \quad F_{,\nu}^{\mu\nu} = -4\pi J^\mu, \quad (2)$$

where

$$F_{\mu\nu} = A_{\nu,\mu} - A_{\mu,\nu}, \quad J^\mu = (\rho, \rho \mathbf{v}). \quad (3)$$

In (3), A_ν is a suitable vector potential and J^μ is the source of the electromagnetic field, namely charges ρ and currents $\rho\mathbf{v}$. In the Lorentz gauge $A_{,\mu}^\mu = 0$, the second pair of Maxwell's equations becomes the wave equation with sources

$$\square A^\mu = -4\pi J^\mu, \quad \square \equiv \nabla^2 - \frac{\partial^2}{\partial t^2}. \quad (4)$$

The equations of motion of a charge in an electromagnetic field are

$$m \frac{dU^\mu}{ds} = e F^{\mu\nu} U_\nu. \quad (5)$$

It turns out that in linearized general relativity, the field equations in the Hilbert gauge $\psi_{,\nu}^{\mu\nu} = 0$ reduce to

$$\square \psi^{\mu\nu} = -16\pi \mathfrak{S}^{\mu\nu}, \quad \psi^{\mu\nu} \equiv h^{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} h, \quad h = \eta_{\mu\nu} h^{\mu\nu} \quad (6)$$

and the equations of motion of the particle (for time-independent gravitational fields) become

$$\frac{dU_\mu}{ds} = (h_{0j,\mu} - h_{0\mu,j}) U^j + \frac{1}{2} h_{00,\mu}. \quad (7)$$

From the above, the linearized equations of general relativity resemble those of electromagnetism with the gravito-electromagnetic ‘tensor’ defined as

$$f_{\mu\nu} \equiv h_{0\nu,\mu} - h_{0\mu,\nu}. \quad (8)$$

The first and second pair of the gravitational field equations (in the Hilbert gauge) can be defined respectively as

$$\begin{aligned} \Pi_{\mu\nu,\alpha} + \Pi_{\alpha\mu,\nu} + \Pi_{\nu\alpha,\mu} &= 0, \quad \Pi_{\mu\nu} \equiv (\psi_{0\nu,\mu} - \psi_{0\mu,\nu}), \\ \square\psi^{\mu\nu} &= -16\pi\mathfrak{S}^{\mu\nu}. \end{aligned} \quad (9)$$

It appears that it may be possible to construct a gravitational energy-momentum (pseudo) tensor with properties similar to those of the electromagnetic energy-momentum tensor. However, it is worth noting that the standard approach identifies $\mathfrak{S}^{\mu\nu}$ with the total energy-momentum pseudo-tensor via the gauge condition $\psi^{\mu\nu}{}_{,\mu} = 0$. The gravitational analog of the electromagnetic Poynting vector is concocted in the following section.

3 The gravitational Poynting vector

We start by noting that the time-time and space-time components of the second pair of the gravitational field equations (9) are given by

$$\Pi_{,\nu}^{\mu\nu} = -16\pi\mathfrak{S}^{0\mu}. \quad (10)$$

The antisymmetry of $\Pi^{\mu\nu}$ yields the conserved quantity (energy)

$$E = \int \mathfrak{S}^{00} d^3x. \quad (11)$$

The Lagrangian density that generates equations (10) is

$$L = -\frac{1}{64\pi} \Pi_{\mu\nu}\Pi^{\mu\nu} - \rho_m U_\alpha B^\alpha, \quad B^\alpha \equiv \psi^{0\alpha}. \quad (12)$$

In the derivation of the field equations (10), B_α is treated as a vector field and not as the component of a two index object. The energy-momentum (pseudo) tensor Θ_ν^μ for the free gravitational field ($\rho_m = 0$) obtained upon symmetrization of

$$\Theta_\nu^\mu = \frac{\partial B_\sigma}{\partial x^\nu} \frac{\partial L}{\partial \left(\frac{\partial B_\sigma}{\partial x^\mu} \right)} - \delta_\nu^\mu L, \quad (13)$$

is essentially that of the free electromagnetic field

$$\Theta_{\nu}^{\mu} = \frac{1}{16\pi} \left(\Pi^{\sigma\mu} \Pi_{\nu\sigma} + \frac{1}{4} \delta_{\nu}^{\mu} \Pi_{\alpha\sigma} \Pi^{\alpha\sigma} \right). \quad (14)$$

It is shown below that the source free gravitational field equations $\Pi^{\mu\nu}{}_{,\nu} = -16\pi \mathfrak{S}^{0\mu}$ and the cyclic property (9) of $\Pi_{\mu\nu,\alpha}$ give the continuity equation $\Theta_{\nu,\mu}^{\mu} = 0$ up to third order in the metric perturbation:

$$\begin{aligned} \Theta_{\nu,\mu}^{\mu} &= \frac{1}{16\pi} \left(\Pi^{\sigma\mu}{}_{,\mu} \Pi_{\nu\sigma} + \Pi^{\sigma\mu} \Pi_{\nu\sigma,\mu} + \frac{1}{2} \Pi_{\alpha\sigma,\nu} \Pi^{\alpha\sigma} \right) \\ \Pi^{\sigma\mu} \Pi_{\nu\sigma,\mu} &= \frac{1}{2} \Pi^{\sigma\mu} (\Pi_{\nu\sigma,\mu} + \Pi_{\mu\nu,\sigma}) \\ \Rightarrow \Pi^{\sigma\mu} \Pi_{\nu\sigma,\mu} + \frac{1}{2} \Pi_{\alpha\sigma,\nu} \Pi^{\alpha\sigma} &= \frac{1}{2} \Pi^{\sigma\mu} (\Pi_{\nu\sigma,\mu} + \Pi_{\mu\nu,\sigma} + \Pi_{\sigma\mu,\nu}) = 0 \\ \Rightarrow \Theta_{\nu,\mu}^{\mu} &= \frac{1}{16\pi} \Pi^{\sigma\mu}{}_{,\mu} \Pi_{\nu\sigma} = -\mathfrak{S}^{0\sigma} \Pi_{\nu\sigma}. \end{aligned} \quad (15)$$

In vacuum $\mathfrak{S}^{0\sigma}$ contains only products of the derivatives of the metric perturbation. Therefore, the right-hand-side of (15) is third order in $\psi^{\mu\nu}$ and can be neglected. This yields $\Theta_{\nu,\mu}^{\mu} = 0$ to third order in the metric perturbation. It will be shown in the following sections that the simple Maxwell-like gravitational Poynting vector is consistent with the radiation flux calculated with the Landau-Lifshitz pseudo-tensor. The gravitational Poynting vector of (14) gives rise to Einstein's quadrupole radiation formula.

4 Linear mass quadrupole oscillator

In this section we calculate the gravitational energy flux of a linear mass quadrupole oscillator using the Maxwell-like gravitational energy-momentum (pseudo) tensor (14). The solution for the linear mass quadrupole oscillator is well known [2,3]. The non-vanishing components of the metric are given by

$$\begin{aligned} \psi^{33} &= A \cos \omega \xi, \quad \psi^{03} = \cos \theta \psi^{33}, \quad \psi^{00} = \cos^2 \theta \psi^{33} \\ A &= \frac{2\omega^2 I}{r}, \quad \xi = t - r. \end{aligned} \quad (16)$$

A direct calculation of the gravitational Poynting vector above leads to

$$\Theta_0^i = \frac{1}{16\pi} (A\omega)^2 \sin^2 \theta \cos^2 \theta \sin^2(\omega\xi) \left[\frac{x^i}{r} \right]. \quad (17)$$

The rate of change of energy averaged over one period of the oscillation is given by

$$\begin{aligned}\frac{dE}{dt} &= - \left\langle \oint_{\sigma} \Theta_0^i d\sigma_i \right\rangle \\ &= -\frac{1}{15} I^2 \omega^6\end{aligned}\tag{18}$$

consistent with the quadrupole radiation formula. An alternative derivation of the gravitational quadrupole radiation formula is provided below.

5 The quadrupole radiation formula

In this section we derive the quadrupole radiation formula using the gravitational Poynting vector (14). The solution of the linearized equations of general relativity (6) is given as:

$$\begin{aligned}\psi_{ab} &= -\frac{2\ddot{I}_{ab}(\xi)}{r}, \quad \psi_{a0} = -\frac{2\ddot{I}_{ab}n_b}{r}, \quad \psi_{00} = \frac{4m}{r} + \frac{2\ddot{I}_{ab}n_a n_b}{r} \\ n_a &= \frac{x^a}{r}, \quad \xi = t - r, \quad I_{ab}(\xi) = \int \rho x_a x_b d^3x\end{aligned}\tag{19}$$

where n_a is the radial unit vector and I_{ab} is the moment of inertia tensor of the system. In (19), $\dot{}$ denotes differentiation with respect to the retarded time ξ . The evaluation of the quadrupole radiation formula is commonly performed with the quadrupole moment tensor

$$Q_{ab} = \int \rho (3x_a x_b - \delta_{ab} r^2) d^3x\tag{20}$$

since it is symmetric and traceless. The moment of inertia tensor in (19) relates to the quadrupole moment tensor as

$$I_{ab} = \frac{1}{3} Q_{ab} + \frac{1}{3} \delta_{ab} \int \rho r^2 d^3x.\tag{21}$$

The second term in (21) doesn't contribute to radiation and can be dropped. Therefore, the energy flux can be obtained replacing $\psi_{\mu\nu}$ in (19) by

$$\Phi_{ab} = -\frac{2\ddot{Q}_{ab}}{3r}, \quad \Phi_{a0} = -\frac{2\ddot{Q}_{ab}n_b}{3r}, \quad \Phi_{00} = \frac{4m}{r} + \frac{2\ddot{Q}_{ab}n_a n_b}{3r}\tag{22}$$

The spatial derivatives of the fields Φ_{ab} involve only derivatives of Q_{ab} since the integration is carried out on a closed surface approaching infinity. On this surface the term proportional to $1/r^2$ is much smaller than that which is proportional to $1/r$. The components of the gravito-electromagnetic ‘tensor’ are given below:

$$\begin{aligned} \Pi_{\mu\nu} &= \Phi_{0\nu,\mu} - \Phi_{0\mu,\nu} \\ \Pi_{ad} &= \frac{2 \ddot{Q}_{db} n_a n_b}{3r} - \frac{2 \ddot{Q}_{ab} n_b n_d}{3r}, \quad \Pi_{0d} = \frac{2 \ddot{Q}_{ab} n_a n_b n_d}{3r} - \frac{2 \ddot{Q}_{da} n_a}{3r}. \end{aligned} \quad (23)$$

The gravitational radiation flux is then given as

$$16\pi\Theta_0^i = -\frac{4}{9r^2}n_i \left(\ddot{Q}_{db}\ddot{Q}_{kl} n_b n_k n_l n_d - \ddot{Q}_{db}\ddot{Q}_{dk} n_b n_k \right). \quad (24)$$

The calculation of the radiated power proceeds in the usual manner by averaging $n_b n_k n_l n_d$ and $n_b n_k$ over all spatial directions using [1]

$$\langle n_b n_k \rangle = \frac{1}{3}\delta_{bk}, \quad \langle n_b n_k n_l n_d \rangle = \frac{1}{15}(\delta_{bk}\delta_{ld} + \delta_{bl}\delta_{kd} + \delta_{bd}\delta_{lk}). \quad (25)$$

The spatial averaging of (24) yields

$$16\pi\Theta_0^i = \frac{4}{45r^2}n_i \ddot{Q}_{ab}\ddot{Q}_{ab}. \quad (26)$$

The power radiated by the system is given by the usual gravitational quadrupole radiation formula (in S.I. units)

$$\frac{dE}{dt} = -\oint_{\sigma} \Theta_0^i d\sigma_i = -\frac{G}{45c^5} \left(\ddot{Q}_{ab} \right)^2. \quad (27)$$

The symmetric energy-momentum (pseudo) tensor (14) allows us to define the angular momentum of the free gravitational field in the familiar scheme

$$M^{\mu\nu} = \int (x^\mu \Theta^{\nu\sigma} - x^\nu \Theta^{\mu\sigma}) dS_\sigma \quad (28)$$

where dS_σ is the element of a three dimensional space-like hypersurface.

6 Conclusion

It was shown that the field equations of general relativity in the Hilbert gauge can be derived from a Lagrangian density similar in structure to that of the electromagnetic field. This Lagrangian generates an energy-momentum (pseudo) tensor possessing the same properties as the electromagnetic energy-momentum tensor. The simple structure of the gravitational energy-momentum (pseudo) tensor constructed in this manner lends itself to an elegant derivation of Einstein's quadrupole radiation formula. The symmetric gravitational (pseudo) tensor allows the definition of angular momentum for the free gravitational field. It should be pointed out that the formalism presented in this paper is restricted to the Hilbert gauge. Although in principle all metrics can be transformed into a coordinate system satisfying the Hilbert gauge condition, future research should be aimed at constructing a gauge invariant gravito-electromagnetic formalism.

Acknowledgements

I would like to thank Professor W. Israel for his encouragement and Sean Bohun for his careful reading of the manuscript.

References

- [1] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press 1995)
- [2] J. Weber *General Relativity and Gravitational Waves* (Interscience Publishers, Inc., New York 1961)
- [3] F.I. Cooperstock and P.H. Lim, Phys. Rev. D 36 (1987) 330