Plebański–Demiański–like solutions in metric–affine gravity

arXiv:gr-qc/9803012v1 3 Mar 1998

Alberto García^{**}, Friedrich W. Hehl^{#†}, Claus Lämmerzahl^{⊲‡},

Alfredo Macías^{\$}, and José Socorro^{\$**}

* Departamento de Física,

CINVESTAV-IPN, Apartado Postal 14-740, C.P. 07000, México, D.F., Mexico

[#]Institute for Theoretical Physics,

University of Cologne, D–50923 Köln, Germany

[⊲] Fakultät für Physik, Universität Konstanz

Postfach 5560 M674, D–78434 Konstanz, Germany

[◊] Departamento de Física,

Universidad Autónoma Metropolitana-Iztapalapa,

Apartado Postal 55-534, C.P. 09340, México, D.F., Mexico.

^{\$} Instituto de Física de la Universidad de Guanajuato,

Apartado Postal E-143, C.P. 37150, León, Guanajuato, Mexico.

(August 23, 2021)

Abstract

We consider a (non–Riemannian) metric–affine gravity theory, in particular its nonmetricity–torsion sector "isomorphic" to the Einstein–Maxwell theory. We map certain Einstein–Maxwell electrovacuum solutions to it, namely the

^{*}E-mail: aagarcia@fis.cinvestav.mx

[†]E-mail: hehl@thp.uni-koeln.de

[‡]E-mail: claus@spock.physik.uni-konstanz.de

[§]E-mail: amac@xanum.uam.mx

^{**}E-mail: socorro@ifug4.ugto.mx

Plebański–Demiański class of Petrov type D metrics. *file plebdem3.tex, 1998-02-27*

PACS numbers: 04.50.+h; 04.20.Jb; 03.50.Kk

Typeset using $\text{REVT}_{E}X$

1. INTRODUCTION

Nowadays, there exists a revival of interest in metric–affine gravity (MAG) theories. It has been demonstrated that they contain the axi–dilatonic sector of *low energy string theory* [1] as special case. Moreover, the gravitational interactions involving the axion and dilaton may be derived from a geometrical action principle involving the curvature scalar with a non–Riemannian connection. In other words, the axi–dilatonic sector of the low energy string theory can be expressed in terms of a geometry with *torsion* and *nonmetricity* [2]. This formulation emphasizes the geometrical nature of the axion and dilaton fields and raises questions about the most appropriate geometry for the discussion of physical phenomena involving these fields.

Recently, it has been proposed that certain MAG models can be reduced to an *effective* Einstein–Proca system [3,4]. Indeed, we have in these kind of models, beside the orthonormal coframe of spacetime, effectively only one extra one–form (co-vector) field as additional degree of freedom.

Very important classes of Petrov type D solutions of the Einstein–Maxwell equations are the Plebański classes. The most general of them is the so–called Plebański–Demiański solution [5], which, as is well known, contains as special cases, among others, the Plebański– Carter, the Kerr–Newman, and the Kerr solutions [6]. In this paper we are going to map this complete space of general electrovacuum solutions to a metric–affine gravity model, which generalizes Einstein's general relativity. Thus, we are able to present solutions to this MAG model and to give to these solutions a physical interpretation.

One arrives at the metric-affine gauge theory of gravity if one gauges the affine group and additionally allows for a metric g [7]. The four-dimensional affine group A(4, R) is the semidirect product of the *translation* group R^4 and the *linear group* GL(4, R), that is, $GL(4, R) = R^4 \otimes GL(4, R)$. The spacetime of MAG encompasses two different post-Riemannian structures: the nonmetricity one-form $Q_{\alpha\beta} = Q_{i\alpha\beta} dx^i$ and the torsion two-form $T^{\alpha} = \frac{1}{2} T_{ij}^{\alpha} dx^i \wedge dx^j$. According to the Yang-Mills fashion, gauge Lagrangians of MAG are quadratic in curvature, torsion, and nonmetricity. One way to investigate the potentialities of such models is to look for *exact* solutions.

The search for exact solutions of MAG began with the work of Tresguerres [8,9], Tucker and Wang [10], Obukhov et al. [11], Vlashinsky et al. [12], and of Puntigam et al. [13]. Macías et al. [14], and Socorro et al. [15] mapped the Einstein–Maxwell sector of dilaton–gravity, emerging from low energy string theory, and found new soliton and multipole solutions of MAG. However, it is important to note that in order to incorporate the scalar dilaton field, one could, for instance, generalize the torsion kink of Baekler et al. [16], an exact solution with an external massless scalar field, or one could turn to the axi–dilatonic sector of MAG [1]. Moreover, solutions implying the existence of torsion shock waves have already been found by García et al. [17]. In this spirit, we are going to look for a wide class of solutions of the vacuum field equations of MAG. Note that such a solution with the additional electromagnetic field of a point charge has been presented in [13]. There it was confirmed that the electromagnetic field is not directly influenced by the post–Riemannian structures torsion and nonmetricity.

A general quadratic Lagrangian in MAG reads [4,7]:

$$V_{\text{MAG}} = \frac{1}{2\kappa} \left[-a_0 R^{\alpha\beta} \wedge \eta_{\alpha\beta} - 2\lambda_{\text{cosm}} \eta + T^{\alpha} \wedge * \left(\sum_{I=1}^{3} a_I^{(I)} T_{\alpha} \right) \right. \\ \left. + 2 \left(\sum_{I=2}^{4} c_I^{(I)} Q_{\alpha\beta} \right) \wedge \vartheta^{\alpha} \wedge * T^{\beta} + Q_{\alpha\beta} \wedge * \left(\sum_{I=1}^{4} b_I^{(I)} Q^{\alpha\beta} \right) \right] \\ \left. - \frac{1}{2} R^{\alpha\beta} \wedge * \left(\sum_{I=1}^{6} w_I^{(I)} W_{\alpha\beta} + \sum_{I=1}^{5} z_I^{(I)} Z_{\alpha\beta} \right) \right]$$
(1.1)

The signature of spacetime is (- + ++), the volume four-form $\eta := *1$, the two-form $\eta_{\alpha\beta} := *(\vartheta_{\alpha} \wedge \vartheta_{\beta})$, and the dimensionless coupling constants read

$$a_0, \ldots a_3, b_1, \ldots b_4, c_2, c_3, c_4, w_1, \ldots w_6, z_1, \ldots z_5.$$
 (1.2)

Moreover, κ is the gravitational and λ_{cosm} the cosmological constant. In suitable units, $\kappa = 1$, which will be assumed in future. In the curvature square term we introduced the antisymmetric part $W_{\alpha\beta} := R_{[\alpha\beta]}$ and the symmetric part $Z_{\alpha\beta} := R_{(\alpha\beta)}$ of the curvature two-form. In $Z_{\alpha\beta}$, we meet a purely post-Riemannian part. Weyl's segmental curvature ${}^{(4)}Z_{\alpha\beta} := R_{\gamma}{}^{\gamma} g_{\alpha\beta}/4 = g_{\alpha\beta} dQ$, with the Weyl covector $Q := Q_{\gamma}{}^{\gamma}/4$, has formally a similar structure as the electromagnetic field strength F = dA, but is physically quite different since it is related to Weyl rescalings.

For the torsion and nonmetricity field configurations, we concentrate on the simplest non-trivial case *with* shear. According to its irreducible decomposition [7], the nonmetricity contains two covector pieces, namely ${}^{(4)}Q_{\alpha\beta} = Q g_{\alpha\beta}$, the dilation piece, and

$$^{(3)}Q_{\alpha\beta} = \frac{4}{9} \left(\vartheta_{(\alpha} e_{\beta)} \rfloor \Lambda - \frac{1}{4} g_{\alpha\beta} \Lambda \right) , \quad \text{with} \quad \Lambda := \vartheta^{\alpha} e^{\beta} \rfloor \mathcal{Q}_{\alpha\beta}^{i} , \quad (1.3)$$

a proper shear piece. Accordingly, our ansatz for the nonmetricity reads

$$Q_{\alpha\beta} = {}^{(3)}Q_{\alpha\beta} + {}^{(4)}Q_{\alpha\beta} \,. \tag{1.4}$$

The torsion, in addition to its tensor piece, encompasses a covector and an axial covector piece. Let us choose only the covector piece as non-vanishing:

$$T^{\alpha} = {}^{(2)}T^{\alpha} = \frac{1}{3} \vartheta^{\alpha} \wedge T, \quad \text{with} \quad T := e_{\alpha} \rfloor T^{\alpha}.$$
(1.5)

Thus we are left with the three non-trivial one-forms Q, Λ , and T. We shall assume that this triplet of one-forms shares the spacetime symmetries, that is, its members are proportional to each other [11–14].

With propagating nonmetricity $Q_{\alpha\beta}$ two types of charge are expected to arise: One dilation charge related by the Noether procedure to the trace of the nonmetricity, the Weyl covector $Q = Q_i dx^i$. It represents the connection associated with gauging the scale transformations (instead of the U(1)-connection in the case of the Maxwell's field). Furthermore, nine shear charges are expected that are related to the remaining traceless piece $\mathcal{Q}_{\alpha\beta} := Q_{\alpha\beta} - Q g_{\alpha\beta}$ of the nonmetricity.

The Lagrangian (1.1) is very complicated, in particular on account of its curvature square pieces. Therefore we have to restrict its generality in order to stay within manageable limits. Our ansatz for the nonmetricity is expected to require a nonvanishing post–Riemannian term quadratic in the segmental curvature. Accordingly, in (1.1) we choose

$$w_1 = \ldots = w_6 = 0, \qquad z_1 = z_2 = z_3 = z_5 = 0,$$
 (1.6)

that is, only z_4 is allowed to survive.

The plan of the paper is as follows: In Sec. 2 a class of solutions, which is related to the Plebański–Demiański solution of the Einstein–Maxwell system, is presented, and in Sec. 3 we shall discuss the results and the further prospects of the theory.

2. PLEBAŃSKI-DEMIAŃSKI-LIKE SOLUTION IN MAG

We start from the coframe of Plebański and Demiański [5] which is specified in terms of the coordinates (τ, y, x, σ) :

$$\vartheta^{\hat{0}} = \frac{1}{H} \sqrt{\frac{Y}{\tilde{\Delta}}} \left(d\tau - x^2 d\sigma \right), \qquad (2.1)$$

$$\vartheta^{\hat{1}} = \frac{1}{H} \sqrt{\frac{\tilde{\Delta}}{Y}} \, dy \,, \tag{2.2}$$

$$\vartheta^{\hat{2}} = \frac{1}{H} \sqrt{\frac{\tilde{\Delta}}{X}} \, dx \,, \tag{2.3}$$

$$\vartheta^{\hat{3}} = \frac{1}{H} \sqrt{\frac{X}{\tilde{\Delta}}} \left(d\tau + y^2 d\sigma \right).$$
(2.4)

Here H = H(x, y), X = X(x), Y = Y(y), and $\tilde{\Delta} = \tilde{\Delta}(x, y)$ are unknown functions. The coframe is orthonormal,

$$g = o_{\alpha\beta} \,\vartheta^{\alpha} \otimes \vartheta^{b}$$
, with $o_{\alpha\beta} = \operatorname{diag}(-1, +1, +1, +1)$. (2.5)

Thus we find the following explicit expression for the metric:

$$g = \frac{1}{H^2} \left[-\frac{Y}{\widetilde{\Delta}} \left(d\tau - x^2 \, d\sigma \right)^2 + \frac{\widetilde{\Delta}}{Y} \, dy^2 + \frac{\widetilde{\Delta}}{X} \, dx^2 + \frac{X}{\widetilde{\Delta}} \left(d\tau + y^2 \, d\sigma \right)^2 \right] \,. \tag{2.6}$$

For the nonmetricity and torsion we assume that they are represented by a *triplet of one*forms, the Weyl covector Q, the covector Λ corresponding to the third irreducible nonmetricity piece, and the torsion trace T.

We substitute the local metric $o_{\alpha\beta}$, the coframe (2.1–2.4), the nonmetricity (1.4), and the torsion (1.5) into the two field equations following from the Lagrangian (1.1) with (1.6) by variation with respect to metric and connection. Then, provided the (rather weak) constraint

$$32a_0^2b_4 - 4a_0a_2b_4 + 64a_0b_3b_4 - 32a_2b_3b_4 + 48a_0b_4c_3 + 24b_4c_3^2 + 24b_3c_4^2 + 12a_0a_2b_3 + 48a_0b_3c_4 - 9a_0c_3^2 + 18a_0c_3c_4 + 3a_0c_4^2 + 6a_0^2a_2 + 24a_0^2c_4 = 0, \qquad (2.7)$$

on the coupling constants (1.2) is fulfilled, we find a general exact solution for the following expressions:

$$\frac{Q}{k_0} = \frac{\Lambda}{k_1} = \frac{T}{k_2} = \frac{H}{\sqrt{\tilde{\Delta}}} \left(\frac{N_{\rm e} y}{\sqrt{Y}} \,\vartheta^{\hat{0}} + \frac{N_{\rm g} x}{\sqrt{X}} \,\vartheta^{\hat{3}} \right) \,, \tag{2.8}$$

$$H(x,y) = 1 - \mu xy,$$

$$X(x) = (b - g^{2}) + 2nx - \epsilon x^{2} + 2m\mu x^{3} - \left(\frac{\lambda_{\text{cosm}}}{3a_{0}} + \mu^{2}(b + e^{2})\right) x^{4},$$

$$Y(y) = (b + e^{2}) - 2my + \epsilon y^{2} - 2n\mu y^{3} - \left(\frac{\lambda_{\text{cosm}}}{3a_{0}} + \mu^{2}(b - g^{2})\right) y^{4},$$

$$\tilde{\Delta}(x,y) = x^{2} + y^{2}.$$
(2.9)

Here $N_{\rm e}$ and $N_{\rm g}$ are the quasi–electric and quasi–magnetic nonmetricity–torsion charges of the source which fulfill

$$\frac{z_4 k_0^2}{2a_0} \left(N_{\rm e}^2 + N_{\rm g}^2 \right) = g^2 + e^2 \,. \tag{2.10}$$

The coefficients k_0, k_1, k_2 in (2.8) are determined by the dimensionless coupling constants (1.2) of the Lagrangian:

$$k_0 := \left(\frac{a_2}{2} - a_0\right) (8b_3 + a_0) - 3(c_3 + a_0)^2, \qquad (2.11)$$

$$k_1 := -9 \left[a_0 \left(\frac{a_2}{2} - a_0 \right) + (c_3 + a_0)(c_4 + a_0) \right], \qquad (2.12)$$

$$k_2 := \frac{3}{2} \left[3a_0(c_3 + a_0) + (8b_3 + a_0)(c_4 + a_0) \right] .$$
(2.13)

Then the constraint (2.7) can be put into the following more compact form

$$b_4 = \frac{a_0 k + 2c_4 k_2}{8k_0}$$
, with $k := 3k_0 - k_1 + 2k_2$, (2.14)

The constants μ , b, g, e, n, and m are free parameters. The parameter ϵ is related to the 2-dimensional spacelike xy-surface, it is $\epsilon = 1$ for spherical, $\epsilon = 0$ for flat, and $\epsilon = -1$ for hyperbolical geometry.

If we collect our results, then the nonmetricity and the torsion read as follows:

$$Q^{\alpha\beta} = \left[k_0 \, o^{\alpha\beta} + \frac{4}{9} \, k_1 \, \left(\vartheta^{(\alpha} e^{\beta)} \rfloor - \frac{1}{4} \, o^{\alpha\beta}\right)\right] \, \frac{H}{\sqrt{\tilde{\Delta}}} \left(\frac{N_{\rm e} \, y}{\sqrt{Y}} \, \vartheta^{\hat{0}} + \frac{N_{\rm g} \, x}{\sqrt{X}} \, \vartheta^{\hat{3}}\right) \,, \tag{2.15}$$

$$T^{\alpha} = \frac{k_2}{3} \,\vartheta^{\alpha} \wedge \,\frac{H}{\sqrt{\widetilde{\Delta}}} \left(\frac{N_{\rm e} \, y}{\sqrt{Y}} \,\vartheta^{\hat{0}} + \frac{N_{\rm g} \, x}{\sqrt{X}} \,\vartheta^{\hat{3}} \right) \,. \tag{2.16}$$

We recognize, see also (2.8), that the members Q, Λ, T of the triplet are proportional to each other. Therefore, we have in our model, besides the spacetime metric, effectively only one extra one-form as additional degree of freedom. This makes it clear why a mapping of our MAG model to the Einstein-Maxwell system and, accordingly, the use of the Plebański-Demiański ansatz is possible, i.e. both models have the same number of degrees of freedom. Indeed, using this ansatz of Plebański and Demiański for a stationary metric and a corresponding ansatz for nonmetricity and torsion, where we additionally assumed that only co-vector parts of these post-Riemannian structures are non-vanishing, we arrived at a general class of solutions for a MAG model.

The physical interpretation of the post-Riemannian parameters of the solution, as described above, is clear: The dilation ('Weyl') charges (related to ${}^{(4)}Q_{\alpha\beta}$) are described by $k_0N_{\rm e}$ and $k_0N_{\rm g}$, the shear charges (related to ${}^{(3)}Q_{\alpha\beta}$) by $k_1N_{\rm e}$ and $k_1N_{\rm g}$, and, eventually, the spin charges (related to ${}^{(2)}T^{\alpha}$) by $k_2N_{\rm e}$ and $k_2N_{\rm g}$, respectively.

The solution (2.1)-(2.5), (2.9), (2.15), and (2.16) found above, was checked with the help of the computer algebra system Reduce [18,19], using its Excalc package [20] for handling exterior differential forms, and by means of the Reduce–based GRG computer algebra system [21].

3. DISCUSSION

The physical motivation to go beyond classical Einstein gravity by means of MAG models is fairly clear and well founded, see the discussion in [22]. One may suspect that the spin–3 modes of the linear connection in the framework of MAG leads to acausalities. However, no detailed investigation has been done into this question so far. Also, in view of the problems of other theories, like supergravity and even string *field theory* [26] in this respect, it appears unfair to ask questions like that of the renormalizability of MAG.

Due to the fact that torsion couples to the *spin* of matter, a discussion of those experiments which may lead to restrictions on torsion also leads, due to (2.8), to restrictions on the two covector parts of the nonmetricity. Therefore, under our triplet ansatz — which certainly describes a highly idealized situation — it is not necessary to devise separate experiments testing the coupling of nonmetricity to the shear current of some matter model.

However, spin $\frac{1}{2}$ matter fields couple to the *axial* vector piece ${}^{(3)}T^{\alpha}$ of the torsion alone, (massless) gauge fields carry a helicity of 1 and do not couple to torsion at all, see e.g. [13]. For massive fields with spin $s = \frac{1}{2}, 1, \frac{3}{2}$, we can extract the following formula for the torsion T^{α} from the literature, see [23] and [24]:

$$T_{\text{as seen by spin } s>0}^{\alpha} = \left(1 - \frac{1}{2s}\right) T^{\alpha} + \frac{3}{2s} {}^{(3)}T^{\alpha} = \left(1 - \frac{1}{2s}\right) \left({}^{(1)}T^{\alpha} + {}^{(2)}T^{\alpha}\right) + \left(1 + \frac{1}{s}\right){}^{(3)}T^{\alpha} .$$
(3.1)

Thereby we recognize that for massive higher spin fields the trace part ${}^{(2)}T^{\alpha}$ of the torsion couples to the spin of these matter fields in the same way as the axial part ${}^{(3)}T^{\alpha}$, modulo numerical factors of the order of unity. Accordingly, we can assume that restrictions on axial torsion also restrict the trace part in a similar way. Analyzing known experiments, we find, with [25], $t_i \leq 1.5 \times 10^{-15} \,\mathrm{m}^{-1}$ and, consequently, $(k_2/k_0) \,Q_i \leq 1.5 \times 10^{-15} \,\mathrm{m}^{-1}$ and $(k_2/k_1) \Lambda_i \leq 1.5 \times 10^{-15} \,\mathrm{m}^{-1}$. Here $T = t_i \,dx^i$, $Q = Q_i \,dx^i$, and $\Lambda = \Lambda_i \,dx^i$.

On the other hand, we presented here a complete class of solutions of MAG. The physical interpretation of the parameters involved in (2.9) can be given as follows: μ is the acceleration parameter, b is related to the angular momentum of the solution, m is the mass and n the NUT parameter [6]. The quasi-electric and quasi-magnetic charges e and g, via (2.10), are related to the torsion and nonmetricity charges $N_{\rm e}$ and $N_{\rm g}$, respectively.

It is important to point out that the generalization of these results to the whole electrovacuum sector of MAG, i.e., including an electromagnetic field as source, is straightforward, and these results will be reported elsewhere.

We want to conclude with two remarks: First, one would like to know at which energy scale such a MAG framework can be regarded as an *effective* gravitational model. According to Ref. [7], the motivation for MAG came mainly from particle physics and the manifield description of an infinite tower of fermions. One may regard such a gauge theory of gravity with Weyl invariance as a small but decisive step towards quantum gravity. Circumstances under which spacetime might become non–Riemannian near Planck energies occur in string theory or in the inflationary model during the early epoch of our universe. The simplest such geometry is metric–affine geometry, in which nonmetricity appears as a field strength, side by side with torsion and curvature.

Secondly, on the one hand the axion-dilaton theory emerges at the low energy limit of string models. On the other hand such models represent one sector of the MAG models. Since these two models have one important sector in common, we should consider the MAG models, in a new perspective, as an effective low energy theory of quantum gravity.

4. ACKNOWLEDGMENTS

This research was supported by CONACyT, grants No. 3544–E9311, No. 3898P–E9608, and by the joint German–Mexican project Conacyt–DLR E130–2924 and DLR–Conacyt MXI 6 B0A 6A. Moreover, C.L. acknowledges support from DAAD and Conacyt, and J.S. from the ANUIES–DAAD agreement, Kennziffer A/98/04459.

REFERENCES

- [1] T. Dereli and R.W. Tucker, *Class. Quantum Grav.* **12** (1995) L31.
- [2] T. Dereli, M. Onder, and R.W. Tucker, *Class. Quantum Grav.* **12** (1995) L251.
- [3] T. Dereli, M. Onder, J. Schray, R.W. Tucker, and C. Wang, Class. Quantum Grav. 13 (1996) L103.
- [4] Yu.N. Obukhov, E.J. Vlachynsky, W. Esser, and F.W. Hehl, *Phys. Rev.* D56 (1997) 7769.
- [5] J. Plebański and M. Demiański, Ann. Phys. (N.Y.) 98 (1976) 98.
- [6] A. García and A. Macías: Black Holes as exact solution of the Einstein-Maxwell equations of Petrov type D. In "Black Holes: Theory and Observation". F.W. Hehl, C. Kiefer, and R. Metzler, eds. (Springer, Berlin, 1998) in print.
- [7] F.W. Hehl, J.D. McCrea, E.W. Mielke, and Y. Ne'eman, *Phys. Rep.* 258 (1995) 1.
- [8] R. Tresguerres, Z. Phys. C65 (1995) 347.
- [9] R. Tresguerres, *Phys. Lett.* A200 (1995) 405.
- [10] R.W. Tucker and C. Wang, *Class. Quantum Grav.* **12** (1995) 2587.
- [11] Yu.N. Obukhov, E.J. Vlachynsky, W. Esser, R. Tresguerres, and F.W. Hehl, *Phys. Lett.* A220 (1996) 1.
- [12] E.J. Vlachynsky, R. Tresguerres, Yu.N. Obukhov, and F.W. Hehl, *Class. Quantum Grav.* 13 (1996) 3253.
- [13] R.A. Puntigam, C. Lämmerzahl, F.W. Hehl, Class. Quantum Grav. 14 (1997) 1347.
- [14] A. Macías, E.W. Mielke and J. Socorro, Class. Quantum Grav. 15 (1998) 445.
- [15] J. Socorro, C. Lämmerzahl, A. Macías, and E.W. Mielke: "Multipole solutions in metric-affine gravity". Submitted for publication (1998).

- [16] P. Baekler, E.W. Mielke, R. Hecht, and F.W. Hehl, Nucl. Phys. B288 (1987) 800.
- [17] A. García, C. Lämmerzahl, A. Macías, E.W. Mielke, and J. Socorro: "Colliding waves in metric–affine gravity". *Phys. Rev.* D57 (1998) in print.
- [18] A.C. Hearn, *REDUCE User's Manual. Version 3.6.* Rand publication CP78 (Rev. 7/95)
 (RAND, Santa Monica, CA 90407-2138, USA, 1995).
- [19] D. Stauffer, F.W. Hehl, N. Ito, V. Winkelmann, and J.G. Zabolitzky: Computer Simulation and Computer Algebra – Lectures for Beginners. 3rd ed. (Springer, Berlin, 1993).
- [20] E. Schrüfer, F.W. Hehl, and J.D. McCrea, Gen. Relat. Grav. 19 (1987) 197.
- [21] V.V. Zhytnikov, GRG. Computer Algebra System for Differential Geometry, Gravity and Field Theory. Version 3.1 (Moscow, 1991) 108 pages.
- [22] Y. Ne'eman and F.W. Hehl, Class. Quantum Grav. 14 (1997) A251.
- [23] M. Seitz, Class. Quantum Gravity 3 (1986) 1265.
- [24] R. Spinosa, Class. Quantum Grav. 4 (1987) 473; Class. Quantum Grav. 4 (1987) 1799.
- [25] C. Lämmerzahl, Phys. Lett. A228 (1997) 223.
- [26] Y. Ne'eman: "Status of quantum gravity" (1997) in: Recent Developments in Gravitation and Mathematical Physics, Proceedings of the 2nd Mexican School on Gravitation and Mathematical Physics, Tlaxcala, Tlax. 1996. A. García et al., eds. (Science Network Publishing, Konstanz, Germany, 1997).