

# Gravitational light bending in Euclidean space

I E Bulyzhenkov

Institute of Spectroscopy RAS, Troitsk, M.o., 142092, Russia

Both the non-homogeneous slowness of electromagnetic waves in gravitational fields and the frequency red shift contribute to the gravitational light bending. This twofold contribution explains the measured deflection of light rays by the Sun under Euclidean geometry of space.

PACS numbers: 04.20.Cv

As known, the observed light deflection by the Sun is nearly twice what Newton's mechanics predicts for particles [1,2]. The hypothetical analogy between particles and light waves was employed by majority of authors for speculations about the curved three-space around sources of gravity.

However, there is no direct analogy in gravitation between the mechanical particle and the electromagnetic wave. The rest-mass particle keeps the measured energy  $m\sqrt{g_{oo}}/\sqrt{1-(dl/d\tau)^2}$  along its trajectory in a static gravitational field, for example, while the measured energy  $\hbar\omega = k_o/\sqrt{g_{oo}}$  of the electromagnetic wave changes along the light ray together with changes of the proper time rate  $d\tau = \sqrt{g_{oo}}dx^o$ . Moreover, the gravitational field works like a medium for electromagnetic waves and leads to their slowness,  $n = \sqrt{\varepsilon\bar{\mu}} = g_{oo}^{-1/2}$ , because  $\mathbf{D} = g_{oo}^{-1/2}\mathbf{E}$  and  $\mathbf{B} = g_{oo}^{-1/2}\mathbf{H}$  in covariant the Maxwell equations [3].

General equations for light rays in static gravitational fields ought to be derived from Fermat's principle [4],

$$\delta \int k_i dx^i = -\delta \int \hbar\omega n dl = -k_o \delta \int \frac{dl}{g_{oo}} = -k_o \delta \int \frac{\sqrt{du^2 + u^2 d\varphi^2}}{u^2(1-GMu)^2} = 0, \quad (1)$$

where  $n^2 k_o k^o = k_i k^i$ ,  $g_{oo} k^o = k_o = const$ ,  $k_i = -nk_o g_{oo}^{-1/2} dx^i/dl$ ,  $\sqrt{g_{oo}} = 1-GMu$ ,  $g_i = -g_{oi}/g_{oo} = 0$ ,  $dl = \sqrt{\delta_{ij} x^i x^j} = \sqrt{dr^2 + r^2 d\varphi^2}$  ( $r \equiv u^{-1}$ ,  $\varphi$ , and  $\vartheta = \pi/2$  are the spherical coordinates). It is worthy of note that both the non-homogeneous wave slowness,  $n = n(g_{oo}) \neq const$ , and the non-homogeneous frequency,  $\omega = \omega(g_{oo}) \neq const$  (red shift), are responsible in (1) for the twofold curvature of light rays in gravitational fields.

The variations of (1) with respect to  $u$  and  $\varphi$  leads to a couple of ray-equations,

$$(1-GMu)^4 \left[ \left( \frac{du}{d\varphi} \right)^2 + u^2 \right] = U_o^2 = const, \quad (2)$$

$$\frac{d^2 u}{d\varphi^2} + u = 2U_o^2 \frac{GM}{(1-GMu)^5}. \quad (3)$$

A family of solutions,  $u \equiv r^{-1} = r_o^{-1} \sin \varphi + 2GM r_o^{-2} (1 + \cos \varphi)$  and  $r_o^{-1} = U_o$ , for both these equations might be found in weak fields, if one ignores all terms nonlinear in  $GM r_o^{-1} \ll 1$ . A propagation of light from  $r(-\infty) = \infty, \varphi(-\infty) = \pi$  to  $r(+\infty) \rightarrow \infty, \varphi(+\infty) \rightarrow \varphi_\infty$  corresponds to the angular deflection  $\varphi_\infty = \arcsin[-2GM r_o^{-1} (1 + \cos \varphi_\infty)] \approx -4GM r_o^{-1}$  from the initial light direction. This result, derived under the flat three-interval  $dl$ , coincides with the measured deflection,  $-1,66'' \pm 0.18''$  [5], of light rays by the Sun ( $r_o = 6,96 \times 10^8 km$  and  $4GM r_o^{-1} = 1,75''$ ).

A four-interval equation for electromagnetic waves in non-stationary gravitational fields may be derived from the light wave-velocity,  $dl/d\tau = n^{-1}$ , in the non-dispersive medium or vacuum, *i.e.*

$$n^2(g_{oo}, g_i) \delta_{ij} dx^i dx^j = g_{oo} (dx^o - g_i dx^i)^2, \quad (4)$$

where the slowness  $n(g_{oo}, g_i)$  is determined by the contribution of gravity into the covariant Maxwell and wave equations. This non-stationary slowness defines the wave and ray equations,  $k_\mu k^\mu = (1 - n^2)(k_\mu V^\mu)^2$  and  $\delta \int k_\mu dx^\mu = 0$ , for light in arbitrary gravitational fields [4].

Note that Euclidean metrics,  $\gamma_{ij} \equiv g_{oi} g_{oj} g_{oo}^{-1} - g_{ij} = \delta_{ij}$ , is hold in (1) and (4) only for 3D space, rather than for 4D space-time. All components of the metric tensor  $g_{\mu\nu} \neq \text{diag} (+1, -1, -1, -1)$  are determined by a particular distribution of external (for an observer) matter. In other words, the three-space with universal Euclidean geometry may be common for all observers, while curved space-times are different pseudo-Riemannian manifolds for different observers.

Euclid's 3D geometry satisfies the observed conservation of a system three-momentum (and angular momentum) at all points of three-space. The homogeneous conservation law for a sum of the three-vectors is possible, in principle, for common three-spaces with constant curvatures (positive, negative, or zero). But only flat three-space, corresponded to all known experiments, may open a simple way for coupling gravity and electromagnetism [6].

- 
- [1] Einstein A 1911 *Über den Einfluss der Schwerkraft auf die Ausbreitung des Lichtes* *Annalen der Physik* **35** 898;
  - [2] Lenard P 1921 *Über die Ablenkung eines Lichtstrahls von seiner geradlinigen Bewegung durch die Attraktion eines Weltkörpers, an welchem er nahe vorbeigeht*, von J Soldner 1801 *Annalen der Physik* **65** 593
  - [3] Landau L D and Lifshitz E M 1971 *The classical theory of fields* (Oxford: Pergamon)
  - [4] Synge J L 1960 *Relativity: the General Theory* (Amsterdam: North-Holland Publishing Company)
  - [5] Will C M 1993 *Theory and experiment in gravitational physics* (Cambridge University Press)
  - [6] Bulzhenkov I E 1997, e-preprint, gr-qc/9708004.