## A Comment on the Paper, "Cosmic Background Radiation"

B.G. Sidharth

## G.P. Birla Observatory & Astronomical Research Centre B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500 063 (India)

In the paper [1] and its updated version [2] it was shown that a collection of Photons with nearly the same energy or frequency exhibit a condensation type of behaviour. To see what this means, our starting point is the formula for the average occupation number for photons of momentum  $\vec{k}$  for all polarizations [3]:

$$\langle n_{\vec{k}} \rangle = \frac{2}{e^{\beta \hbar \omega} - 1} \tag{1}$$

Let us specialize to a scenario in which all the photons have nearly the same energy so that we can write,

$$\langle n_{\vec{k}} \rangle = \langle n_{\vec{k'}} \rangle \delta(k - k'), \tag{2}$$

where  $\langle n'_k \rangle$  is given by (1), and  $k \equiv |\vec{k}|$ . The total number of photons N, in the volume V being considered, can be obtained in the usual way,

$$N = \frac{V[k]}{(2\pi)^3} \int_0^\infty dk 4\pi k^2 \langle n_k \rangle \tag{3}$$

where V is large. Inserting (2) in (3) we get,

$$N = \frac{2V}{(2\pi)^3} 4\pi k^{\prime 2} [\epsilon^{\Theta} - 1]^{-1} [k], \Theta \equiv \beta \hbar \omega, \qquad (4)$$

In the above,  $[k] \equiv [L^{-1}]$  is a dimensionality constant, introduced to compensate the loss of a factor k in the integral (3), owing to the  $\delta$ -function in (2): That is, a volume integral in  $\vec{k}$  space is reduced to a surface integral on the sphere  $[\vec{k}] = k'$ , due to our constraint that all photons have nearly the same energy.

We observe that,  $\Theta = \hbar \omega / KT \approx 1$ , since by (2), the photons have nearly the same energy  $\hbar \omega$ . We also introduce,

$$v = \frac{V}{N}, \lambda = \frac{2\pi c}{\omega} = \frac{2\pi}{k} \text{ and } z = \frac{\lambda^3}{v}$$
 (5)

 $\lambda$  being the wavelength of the radiation. We now have from (4), using (5),

$$(e-1) = \frac{vk^{\prime 2}}{\pi^2}[k] = \frac{8\pi}{k'z}[k]$$

Using (5) we get:

$$z = \frac{8\pi}{k'(e-1)} = \frac{4\pi}{(e-1)}[k] \tag{6}$$

From (6) we conclude that, when

$$\lambda = \frac{e-1}{4} = 0.4[L]$$
(7)

then,

$$z \approx 1 \tag{8}$$

or conversely. We are working in the cgs system, so that the units in (7) are cm. As pointed out, this could be further confirmed from other points of view.

We next observe that the author's 1997 cosmology used the ubiquitous zero point energy to deduce a model of an accelerating universe with a small cosmological constant (unlike in its previous 1960s version of Zeldovich and others) which lead to the famous cosmological constant problem [4].

At this time the ruling cosmological paradigm was exactly the opposite – a dark matter dominated decelerating universe [5, 6, 7]. It has also been known that the Zero Point Field leads to the Lamb shift [8, 9]. It is quite remarkable that radiations due to the Lamb shift have a frequency of around 1000 megacycles, which falls right within the microwave region to a few centimeters radiation. So there is a sea of microwave photons in the universe whose frequencies differ by small amounts. By the above argument these nearly mono energetic photons would condense to a value around 4 millimeters wavelength which is exactly the cosmic microwave background.

## References

- Sidharth, B.G. (2000). Chaos, Solitons & Fractals 11, 2000, pp.1471-1472.
- [2] Sidharth, B.G. and Valluri, S.R. (2015). Int. J. Th. Phys Vol.54(8), 2015, pp.2792-2797.
- [3] Huang, K. (1975). Statistical Mechanics (Wiley Eastern, New Delhi), pp.75ff.
- [4] Weinberg, S. (1979). *Phys. Rev. Lett.* 43, pp.1566.
- [5] Sidharth, B.G. (1999). Proc. of the Eighth Marcell Grossmann Meeting on General Relativity (1997) Piran, T. (ed.) (World Scientific, Singapore), pp.476–479.
- [6] Sidharth, B.G. (1998). Int. J. of Mod. Phys. A 13, (15), pp.2599ff.
- Sidharth, B.G. (1998). International Journal of Theoretical Physics Vol.37, No.4, pp.1307–1312.
- [8] Bjorken, J.D. and Drell, S.D. (1964). Relativistic Quantum Mechanics (Mc-Graw Hill, New York), pp.39.
- [9] Itzykson, C. and Zuber, J. (1980). Quantum-Field Theory (Mc-Graw Hill, New York), pp.139.