

**TODA CHAINS WITH TYPE  $A_m$  LIE ALGEBRA FOR  
MULTIDIMENSIONAL CLASSICAL COSMOLOGY WITH  
INTERSECTING  $p$ -BRANES**

V. R. GAVRILOV, V. N. MELNIKOV

*Centre for Gravitation and Fundamental Metrology VNIIMS,  
3-1 M. Ulyanovoy St., Moscow 117313, Russia  
E-mail: melnikov@rqs.phys.msu.su*

We consider a  $D$ -dimensional cosmological model describing an evolution of  $(n+1)$  Einstein factor spaces ( $n \geq 2$ ) in the theory with several dilatonic scalar fields and generalized electro-magnetic forms, admitting an interpretation in terms of intersecting  $p$ -branes. The equations of motion of the model are reduced to the Euler-Lagrange equations for the so called pseudo-Euclidean Toda-like system. We consider the case, when characteristic vectors of the model, related to  $p$ -branes configuration and their couplings to the dilatonic fields, may be interpreted as the root vectors of a Lie algebra of the type  $A_m$ . The model is reduced to the open Toda chain and integrated. The exact solution is presented in the Kasner-like form.

## 1 Introduction

Last years have witnessed a growth of interest to classical  $p$ -brane solutions of (bosonic sector of) supergravities in various dimensions.<sup>1-21</sup> This interest is inspired by a conjecture that  $D = 11$  supergravity is a low-energy effective field theory of eleven-dimensional fundamental  $M$ -theory, which (together with so called  $F$ -theory) is a candidate for unification of five known  $D = 10$  superstring theories.<sup>22-28</sup> Classical  $p$ -brane solutions may be considered as a method for investigation of interlinks between superstrings and  $M$ -theory.

In this paper we consider a generalized bosonic sector of early supergravity theories<sup>29</sup> in the form of multidimensional gravitational model with several dilatonic fields and Maxwell-like forms of various ranks. When  $D$ -dimensional space-time manifold is a product of several Einstein factor spaces, the most convenient to use  $\sigma$ -model approach (see, for instance,<sup>4,30,31</sup>) for constructing exact solutions with  $p$ -branes. It was shown<sup>18</sup> that for cosmological space-times the equations of motion to such  $\sigma$ -model are reduced to the Euler-Lagrange equations for a pseudo-Euclidean Toda-like Lagrange system. The methods for integrating of pseudo-Euclidean Toda-like systems were developed in the papers (see, for instance,<sup>32-34</sup> and references therein) devoted to the multidimensional cosmologies with multicomponent perfect fluid. Here we integrate the intersecting  $p$ -branes model reducible to Toda chain associated with Lie algebra of the type  $A_m$ .

## 2 The general model

Following the papers<sup>4,5,9-15,18-21</sup> we study a classical model in  $D$  dimensions described by the action

$$S = \int_{\mathcal{M}} d^D z \sqrt{|g|} \{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{1}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)_g^2 \}, \quad (1)$$

where  $g = g_{MN} dz^M \otimes dz^N$  is the metric on  $D$ -dimensional manifold  $\mathcal{M}$  ( $M, N = 1, \dots, D$ ),  $|g| = |\det(g_{MN})|$ . We denoted by  $\varphi = (\varphi^\alpha) \in \mathbb{R}^l$  a vector from dilatonic scalar fields,  $(h_{\alpha\beta})$  is a non-degenerate  $l \times l$  matrix ( $l \in \mathbb{N}$ ) and  $\lambda_a$  is a 1-form on  $\mathbb{R}^l$ :  $\lambda_a(\varphi) = \lambda_{a\alpha} \varphi^\alpha$ ,  $a \in \Delta$ ,  $\alpha = 1, \dots, l$ .

$$F^a = dA^a = \frac{1}{n_a!} F_{M_1 \dots M_{n_a}}^a dz^{M_1} \wedge \dots \wedge dz^{M_{n_a}} \quad (2)$$

is a  $n_a$ -form ( $n_a \geq 1$ ) on  $\mathcal{M}$ . In Eq. (1) we denoted

$$(F^a)_g^2 = F_{M_1 \dots M_{n_a}}^a F_{N_1 \dots N_{n_a}}^a g^{M_1 N_1} \dots g^{M_{n_a} N_{n_a}}, \quad (3)$$

$a \in \Delta$ , where  $\Delta$  is some finite set.

Equations of motion corresponding to (1) have the following form

$$R_{MN} - \frac{1}{2} g_{MN} R = T_{MN}, \quad (4)$$

$$\Delta[g] \varphi^\alpha - \sum_{a \in \Delta} \frac{\lambda_a^\alpha}{n_a!} e^{2\lambda_a(\varphi)} (F^a)_g^2 = 0, \quad (5)$$

$$\nabla_{M_1} [g] (e^{2\lambda_a(\varphi)} F^{a, M_1 \dots M_{n_a}}) = 0, \quad (6)$$

$a \in \Delta$ ;  $\alpha = 1, \dots, l$ . In Eq. (5)  $\lambda_a^\alpha = h^{\alpha\beta} \lambda_{a\beta}$ , where  $(h^{\alpha\beta})$  is a matrix inverse to  $(h_{\alpha\beta})$ . In (4) we denoted

$$T_{MN} = T_{MN}[\varphi, g] + \sum_{a \in \Delta} e^{2\lambda_a(\varphi)} T_{MN}[F^a, g], \quad (7)$$

where

$$T_{MN}[\varphi, g] = h_{\alpha\beta} \left( \partial_M \varphi^\alpha \partial_N \varphi^\beta - \frac{1}{2} g_{MN} \partial_P \varphi^\alpha \partial^P \varphi^\beta \right), \quad (8)$$

$$T_{MN}[F^a, g] = \frac{1}{n_a!} \left[ -\frac{1}{2} g_{MN} (F^a)_g^2 + n_a F_{M M_2 \dots M_{n_a}}^a F_N^{a, M_2 \dots M_{n_a}} \right]. \quad (9)$$

In Eqs. (5), (6)  $\Delta[g]$  and  $\nabla[g]$  are Laplace-Beltrami and covariant derivative operators corresponding to  $g$ .

We consider the manifold

$$\mathcal{M} = \mathbb{R} \times M_0 \times \dots \times M_n \quad (10)$$

with the metric

$$g = -e^{2\gamma(t)} dt \otimes dt + \sum_{i=0}^n e^{2x^i(t)} g^i, \quad (11)$$

where  $g^i = g_{m_i n_i}^i(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$  is a metric on  $M_i$  satisfying the equation

$$R_{m_i n_i} [g^i] = \xi_i g_{m_i n_i}^i, \quad (12)$$

$m_i, n_i = 1, \dots, d_i$ ;  $d_i = \dim M_i$ ,  $\xi_i = \text{const}$ ,  $i = 0, \dots, n$ ;  $n \in \mathbf{N}$ . So all  $(M_i, g^i)$  are Einstein spaces.

Each manifold  $M_i$  is supposed to be oriented and connected. Then the volume  $d_i$ -form

$$\tau_i = \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad (13)$$

and the signature

$$\text{sign det}(g_{m_i n_i}^i) = 1 \quad (14)$$

are correctly defined for all  $i = 0, \dots, n$ .

Let

$$\Omega_0 = \{\emptyset, \{0\}, \{1\}, \dots, \{n\}, \{0, 1\}, \dots, \{0, 1, \dots, n\}\} \quad (15)$$

be a set of all ordered subsets of  $I_0 \equiv \{0, \dots, n\}$ . For any  $I = \{i_1, \dots, i_k\} \in \Omega_0$ ,  $i_1 < \dots < i_k$ , we define a volume form

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k} \quad (16)$$

of rank

$$d(I) \equiv \sum_{i \in I} d_i = d_{i_1} + \dots + d_{i_k} \quad (17)$$

and a corresponding  $p$ -brane submanifold

$$M_I \equiv M_{i_1} \times \dots \times M_{i_k}, \quad (18)$$

where  $p = d(I) - 1$  ( $\dim M_I = d(I)$ ).

We adopt the following "composite electro-magnetic" Ansatz for fields of forms

$$F^a = \sum_{I \in \Omega_{a,e}} F^{(a,e,I)} + \sum_{J \in \Omega_{a,m}} F^{(a,m,J)}, \quad (19)$$

where

$$F^{(a,e,I)} = d\Phi^{(a,e,I)} \wedge \tau(I), \quad (20)$$

$$F^{(a,m,J)} = e^{-2\lambda_a(\varphi)} * \left( d\Phi^{(a,m,J)} \wedge \tau(J) \right), \quad (21)$$

$a \in \Delta$ ,  $I \in \Omega_{a,e}$ ,  $J \in \Omega_{a,m}$  and

$$\Omega_{a,e}, \Omega_{a,m} \subset \Omega_0. \quad (22)$$

(For empty  $\Omega_{a,v} = \emptyset$ ,  $v = e, m$ , we put  $\sum_{\emptyset} = 0$  in Eq. (19)). In Eq. (21)  $* = *[g]$  is the Hodge operator on  $(M, g)$ .

For the potentials in Eqs. (20), (21) we put

$$\Phi^s = \Phi^s(t), \quad (23)$$

$s \in S$ , where

$$S = S_e \sqcup S_m, \quad S_v \equiv \prod_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (24)$$

$v = e, m$ .

For dilatonic scalar fields we also put

$$\varphi^\alpha = \varphi^\alpha(t), \quad (25)$$

$\alpha = 1, \dots, l$ .

From Eqs. (20) and (21) we obtain the relations between dimensions of  $p$ -brane worldsheets and ranks of forms

$$d(I) = n_a - 1, \quad I \in \Omega_{a,e}, \quad (26)$$

$$d(J) = D - n_a - 1, \quad J \in \Omega_{a,m} \quad (27)$$

in electric and magnetic cases respectively.

We put the following restrictions on  $\Omega_{a,v}$ . Let

$$w_1 \equiv \{i \mid i \in \{1, \dots, n\}, d_i = 1\}. \quad (28)$$

The set  $w_1$  describes all 1-dimensional manifolds among  $M_i$  ( $i \geq 0$ ). We impose the following restrictions on the sets  $\Omega_{a,v}$  (22):

$$W_{ij}(\Omega_{a,v}) = \emptyset, \quad (29)$$

$a \in \Delta; v = e, m; i, j \in w_1, i < j$  and

$$W_j^{(1)}(\Omega_{a,m}, \Omega_{a,e}) = \emptyset, \quad (30)$$

$a \in \Delta; j \in w_1$ . Here

$$W_{ij}(\Omega_*) \equiv \{(I, J) | I, J \in \Omega_*, I = \{i\} \sqcup (I \cap J), J = \{j\} \sqcup (I \cap J)\}, \quad (31)$$

$i, j \in w_1, i \neq j, \Omega_* \subset \Omega_0$  and

$$W_j^{(1)}(\Omega_{a,m}, \Omega_{a,e}) \equiv \{(I, J) \in \Omega_{a,m} \times \Omega_{a,e} | \bar{I} = \{j\} \sqcup J\}, \quad (32)$$

$j \in w_1$ . In (32)

$$\bar{I} \equiv I_0 \setminus I \quad (33)$$

is a "dual" set. (The restrictions (29) and (30) are trivially satisfied when  $n_1 \leq 1$  and  $n_1 = 0$ , respectively, where  $n_1 = |w_1|$  is the number of 1-dimensional manifolds among  $M_i$ ).

It was shown<sup>18</sup> that after the following gauge fixing

$$\gamma = \gamma_0 \equiv \sum_{i=0}^n d_i x^i, \quad (34)$$

the Maxwell equations (6) for  $s \in S_e$  and Bianchi identities  $dF^s = 0$  for  $s \in S_m$  look as follows

$$\frac{d}{dt} \left( \exp[-2\gamma_0 + 2\chi_s \lambda_{a_s}(\varphi)] \dot{\Phi}^s \right) = 0, \quad (35)$$

where

$$\chi_s = +1, \quad v_s = e; \quad (36)$$

$$\chi_s = -1, \quad v_s = m. \quad (37)$$

Then

$$\dot{\Phi}^s = Q_s \exp[2\gamma_0 - 2\chi_s \lambda_{a_s}(\varphi)], \quad (38)$$

where  $Q_s$  are constants,  $s = (a_s, v_s, I_s) \in S$ . Let

$$Q_s \neq 0, \quad s \in S_*; \quad (39)$$

$$Q_s = 0, \quad s \in S \setminus S_*,$$

where  $S_* \subset S$  is a non-empty subset of  $S$ .

For fixed  $Q = (Q_s, s \in S_*)$  the equations of motion (4),(5) are equivalent (after the gauge fixing) to the Euler-Lagrange equations following from the Lagrangian

$$L_Q = \frac{1}{2} \bar{G}_{AB} \dot{x}^A \dot{x}^B - V_Q \quad (40)$$

under the zero-energy constraint

$$E_Q = \frac{1}{2} \bar{G}_{AB} \dot{x}^A \dot{x}^B + V_Q = 0. \quad (41)$$

We denoted

$$(x^A) = (x^i, \varphi^\alpha), \quad (42)$$

$$V_Q = -\frac{1}{2} \sum_{i=0}^n \xi^i d^i \exp[-2x^i + 2\gamma_0] + \frac{1}{2} \sum_{s \in S_*} Q_s^2 \exp[2\gamma_0 - 2\chi_s \lambda_{a_s}(\varphi)] \quad (43)$$

and

$$(\bar{G}_{AB}) = \begin{pmatrix} G_{ij} & 0 \\ 0 & h_{\alpha\beta} \end{pmatrix}, \quad (44)$$

$$G_{ij} = d_i \delta_{ij} - d_i d_j. \quad (45)$$

A system with Lagrangian of the type (40) is called the pseudo-Euclidean Toda-like system. <sup>34</sup> In the next section we consider a special case, when pseudo-Euclidean Toda-like system is reducible to the Toda lattice associated with a Lie algebra of the type  $A_m$ .

### 3 Integration of the models reducible to a classical open Toda chain

Now we consider the general model under the following restrictions

1. The factor space  $M_0$  has non-zero Ricci tensor and all remaining factor spaces  $M_1, \dots, M_n$  are Ricci flat, i.e.

$$\xi_0 \neq 0, \quad \xi_1 = \dots = \xi_n = 0.$$

2. All nonvanishing (with  $Q_s \neq 0$ )  $p$ -branes do not "live" in the factor space  $M_0$ , i.e.

$$0 \notin I_s, \quad \forall s \in S_*.$$

We introduce  $(n + l + 1)$ -dimensional real vector space  $\mathbf{R}^{n+l+1}$  with a canonical basis  $\{e_A\}$ ,  $A = 0, \dots, n + l$ , where  $e_0 = (1, 0, \dots, 0)$  etc. Hereafter we use the following vectors

1. The vector  $x(t)$  we need to calculate

$$x(t) = \sum_{A=0}^{n+l+1} x^A(t) e_A, \quad x^A(t) = (x^i(t), \varphi^\alpha(t)), \quad (46)$$

$i = 0, \dots, n$ ,  $\alpha = 1, \dots, l$ . We remind that  $\exp[x^i(t)]$  is the scale factor of the space  $M_i$ ,  $\varphi^\alpha(t)$  is the  $\alpha$ -th dilatonic scalar field.

2. The vector  $U_0$  induced by the curvature of the factor space  $M_0$

$$U_0 = \sum_{A=0}^{n+l+1} U_0^A e_A, \quad U_0^A = \left( -\frac{\delta_{i0}}{d_0}, 0, \dots, 0 \right), \quad (47)$$

$i = 0, \dots, n$ .

3. The vector  $U_s$  induced by the  $p$ -brane corresponding to  $s = (a_s, v_s, I_s) \in S_*$

$$U_s = \sum_{A=0}^{n+l+1} U_s^A e_A, \quad U_s^A = \left( \delta_{iI_s} - \frac{d(I_s)}{D-2}, -\chi_s \lambda_{a_s}^\alpha \right), \quad (48)$$

$i = 0, \dots, n$ ,  $\alpha = 1, \dots, l$ .

Let  $(\cdot, \cdot)$  be a symmetrical bilinear form defined in  $\mathbf{R}^{n+l+1}$  by the following manner

$$(e_A, e_B) = \bar{G}_{AB}. \quad (49)$$

The bilinear form  $(\cdot, \cdot)$  for the vectors  $U_0, U_s$  reads

$$(U_0, U_0) = \frac{1}{d_0} - 1, \quad (U_0, U_s) = 0, \quad \forall s \in S_*, \quad (50)$$

$$(U_s, U_{s'}) = d(I_s \cap I_{s'}) - \frac{d(I_s)d(I_{s'})}{D-2} + \chi_s \chi_{s'} \sum_{\alpha, \beta=1}^l \lambda_{a_s, \alpha} \lambda_{a_{s'}, \beta} h^{\alpha\beta}. \quad (51)$$

Using the notation  $(\cdot, \cdot)$  and the vectors (46)-(48), we may present the Lagrangian and the zero energy constraint for this special type of the model in the form

$$L_Q = \frac{1}{2}(\dot{x}, \dot{x}) - V_Q, \quad E_Q = \frac{1}{2}(\dot{x}, \dot{x}) + V_Q = 0, \quad (52)$$

where

$$V_Q = a^{(0)} e^{2(U_0, x)} + \sum_{s \in S_*} a^{(s)} e^{2(U_s, x)}. \quad (53)$$

We used the following constants in Eq. (53)

$$a^{(0)} = -\frac{1}{2} \xi_0 d_0, \quad a^{(s)} = \frac{1}{2} Q_s^2. \quad (54)$$

Let the vectors  $U_s \in \mathbf{R}^{n+l+1}$  satisfy the following conditions

$$(U_s, U_s) = U^2 > 0, \quad \forall s \in S_*, \quad (55)$$

$$(C_{ss'}) = \left( \frac{2(U_s, U_{s'})}{(U_{s'}, U_{s'})} \right) = \begin{pmatrix} 2 & -1 & 0 & \dots & 0 & 0 \\ -1 & 2 & -1 & \dots & 0 & 0 \\ 0 & -1 & 2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 2 & -1 \\ 0 & 0 & 0 & \dots & -1 & 2 \end{pmatrix}. \quad (56)$$

where  $(C_{ss'})$  is the Cartan matrix for the Lie algebra  $A_{|S_*|} = sl(|S_*|+1, C)$ . We mention that the number of the vectors  $U_0, U_s$  should not exceed the dimension of  $\mathbf{R}^{n+l+1}$ , i.e.

$$n + l - |S_*| \equiv n' \geq 0. \quad (57)$$

We choose in  $\mathbf{R}^{n+l+1}$  a basis  $\{f_0, f_s, f_p\}$ ,  $s \in S_*$ ,  $p = |S_*|+1, \dots, |S_*|+n'$ , with the properties

$$f_0 = \frac{U_0}{(U_0, U_0)}, \quad f_s = \frac{U_s}{(U_s, U_s)}, \quad s \in S_*, \quad (58)$$

$$(f_0, f_p) = (f_s, f_p) = 0, \quad (f_p, f_{p'}) = 0, \quad p \neq p', \quad (59)$$

$p, p' = |S_*|+1, \dots, |S_*|+n'$ . It should be noted that if  $n' = 0$ , then the vectors  $f_p$  do not appear.

Using the decomposition

$$x(t) = q^0(t) f_0 + \sum_{s \in S_*} q^s(t) f_s + \sum_{p=|S_*|+1}^{|S_*|+n'} q^p(t) f_p \quad (60)$$

with respect to this basis, we present the Lagrangian and the zero-energy constraint (52) in the form

$$L_Q = L_0 + L_{Q,T} + L_f, \quad E_Q = E_0 + E_{Q,T} + E_f = 0, \quad (61)$$



where

$$L_0 = \frac{(\dot{q}^0)^2}{2(U_0, U_0)} - V_0, \quad E_0 = \frac{(\dot{q}^0)^2}{2(U_0, U_0)} + V_0, \quad V_0 = a^{(0)} e^{2q^0}, \quad (62)$$

$$L_f = E_f = \frac{1}{2} \sum_{p=|S_*|+1}^{|S_*|+n'} (f_p, f_p) (\dot{q}^p)^2, \quad (63)$$

$$L_{Q,T} = \sum_{s,s' \in S_*} \frac{C_{ss'}}{4U^2} \dot{q}^s \dot{q}^{s'} - V_{Q,T}, \quad E_{Q,T} = \sum_{s,s' \in S_*} \frac{C_{ss'}}{4U^2} \dot{q}^s \dot{q}^{s'} + V_{Q,T}, \quad (64)$$

$$V_{Q,T} = \sum_{s' \in S_*} a^{(s)} \exp \left[ \sum_{s' \in S_*} C_{ss'} q^{s'} \right]. \quad (65)$$

Integration of equations of motion for  $q^0$  is straightforward and leads to the following result

$$e^{-q^0(t)} = F(t - t_0), \quad (66)$$

where  $t_0$  is an arbitrary constant and the function  $F_0$  has the form

$$F_0(t) = \sqrt{-2(U_0, U_0)a^{(0)}|t|}, \quad \text{if } a^{(0)} > 0, \quad E_0 = 0, \quad (67)$$

$$= \sqrt{|a^{(0)}/E_0|} \cosh[\sqrt{2(U_0, U_0)E_0}t], \quad \text{if } a^{(0)} < 0, \quad E_0 < 0, \quad (68)$$

$$= \sqrt{|a^{(0)}/E_0|} \sinh[\sqrt{2(U_0, U_0)E_0}|t|], \quad \text{if } a^{(0)} > 0, \quad E_0 < 0, \quad (69)$$

$$= \sqrt{|a^{(0)}/E_0|} \sin[\sqrt{-2(U_0, U_0)E_0}|t|], \quad \text{if } a^{(0)} > 0, \quad E_0 > 0. \quad (70)$$

For  $q^p$  we get

$$e^{-q^p(t)} = e^{A^p t + B^p}, \quad p = |S_*| + 1, \dots, |S_*| + n', \quad (71)$$

$$E_f = \frac{1}{2} \sum_{p=|S_*|+1}^{|S_*|+n'} (f_p, f_p) (A^p)^2, \quad (72)$$

where  $A^p$  and  $B^p$  are arbitrary constants.

Using the following translation

$$q^s \mapsto q^s - \ln C^s, \quad (73)$$

where the constants  $C^s$  satisfy

$$\sum_{s' \in S_*} C_{ss'} \ln C^{s'} = \ln[2U^2 a^{(s)}], \quad s \in S_*, \quad (74)$$

we get  $L_{Q,T}$  and  $E_{Q,T}$  in the form

$$L_{Q,T} = \frac{1}{4U^2}L_T, \quad E_{Q,T} = \frac{1}{4U^2}E_T, \quad (75)$$

where

$$L_T = \frac{1}{2} \sum_{s,s' \in S_*} C_{ss'} \dot{q}^s \dot{q}^{s'} - V_T, \quad E_T = \frac{1}{2} \sum_{s,s' \in S_*} C_{ss'} \dot{q}^s \dot{q}^{s'} + V_T, \quad (76)$$

$$V_T = \sum_{s' \in S_*} \exp \left[ \sum_{s' \in S_*} C_{ss'} q^{s'} \right]. \quad (77)$$

We notice that  $L_T$  represents the Lagrangian of a Toda chain associated with the Lie algebra  $A_{|S_*|} = sl(|S_*| + 1, C)$ , when the root vectors are put into the Chevalley basis and the coordinate describing the motion of the mass center is separated out. Using the result of Anderson<sup>35</sup> we present the solution of the equations of motion following from  $L_T$

$$\begin{aligned} e^{-q^s(t)} &\equiv F_s(t) \\ &= \sum_{r_1 < \dots < r_{m(s)}}^{|S_*|+1} v_{r_1} \cdots v_{r_{m(s)}} \Delta^2(r_1, \dots, r_{m(s)}) e^{(w_{r_1} + \dots + w_{r_{m(s)}})t}, \end{aligned} \quad (78)$$

where we denoted by  $m(s)$  the number of element  $s \in S_*$ . The  $m(s)$  is assigned to  $s$  in accordance with Eq. (56).  $\Delta^2(r_1, \dots, r_{m(s)})$  denotes the square of the Vandermonde determinant

$$\Delta^2(r_1, \dots, r_m) = \prod_{r_i < r_j} (w_{r_i} - w_{r_j})^2. \quad (79)$$

The constants  $v_r$  and  $w_r$  have to satisfy the relations

$$\prod_{r=1}^{|S_*|+1} v_r = \Delta^{-2}(1, \dots, |S_*| + 1), \quad \sum_{r=1}^{|S_*|+1} w_r = 0. \quad (80)$$

The energy of the Toda chain described by this solution is given by

$$E_T = \frac{1}{2} \sum_{r=1}^{|S_*|+1} w_r^2. \quad (81)$$

Taking into account the transformation (73), we get

$$e^{-q^s(t)} \equiv \tilde{F}_s(t) = \frac{F_s(t)}{C^s}, \quad s \in S_*, \quad (82)$$

$$E_{Q,T} = \frac{1}{4U^2} \sum_{r=1}^{|S_*|+1} w_r^2. \quad (83)$$

Finally, we obtain the following decomposition of the vector  $x(t)$

$$x(t) = \ln[F_0(t-t_0)] \frac{U_0}{1-1/d_0} - \sum_{s \in S_*} \ln[\tilde{F}_s(t)] \frac{U_s}{U^2} - \sum_{p=|S_*|+1}^{|S_*|+n'} [A^p t + B^p] f_p. \quad (84)$$

We introduce the following vectors in  $\mathbf{R}^{n+l+1}$

$$a \equiv \sum_{A=0}^{n+l} a^A e_A = \sum_{p=|S_*|+1}^{|S_*|+n'} A^p f_p, \quad b \equiv \sum_{A=0}^{n+l} b^A e_A = \sum_{p=|S_*|+1}^{|S_*|+n'} B^p f_p. \quad (85)$$

Their coordinates  $a^A$  and  $b^A$  with respect to the canonical basis satisfy the relations

$$(a, U_0) = \sum_{A,B=0}^{n+l} G_{AB} a^A U_0^B = 0, \quad (a, U_s) = \sum_{A,B=0}^{n+l} G_{AB} a^A U_s^B = 0, \quad (86)$$

$$(b, U_0) = \sum_{A,B=0}^{n+l} G_{AB} b^A U_0^B = 0, \quad (b, U_s) = \sum_{A,B=0}^{n+l} G_{AB} b^A U_s^B = 0, \quad (87)$$

$$(a, a) = \sum_{A,B=0}^{n+l} G_{AB} a^A a^B = 2E_f, \quad (88)$$

$s \in S_*$ . We note that, if  $n' = n + l - |S_*| = 0$ , we must put  $a = b = 0$ .

Using Eq. (84) with Eq. (85), the definitions (46)-(48), we obtain the scale factors  $\exp[x^i(t)]$  and the dilatonic fields  $\varphi^\alpha(t)$

$$e^{x^0(t)} = [F_0(t-t_0)]^{\frac{1}{d_0-1}} \prod_{s \in S_*} [\tilde{F}_s(t)]^{\frac{d(I_s)}{(D-2)U^2}} \exp[a^0 t + b^0], \quad (89)$$

$$e^{x^i(t)} = \prod_{s \in S_*} [\tilde{F}_s(t)]^{\frac{1}{U^2} \left( \frac{d(I_s)}{D-2} - \delta_{iI_s} \right)} \exp[a^i t + b^i], \quad i = 1, \dots, n, \quad (90)$$

$$\varphi^\alpha(t) = \prod_{s \in S_*} [\tilde{F}_s(t)]^{\frac{1}{U^2} \chi_s \lambda_{\alpha s}} \exp[a^{n+1+\alpha} t + b^{n+1+\alpha}], \quad \alpha = 1, \dots, l. \quad (91)$$

For the potentials  $\Phi^s$  we obtain

$$\dot{\Phi}^s(t) = Q_s \prod_{s' \in S_*} [\tilde{F}_{s'}(t)]^{-C_{ss'}}. \quad (92)$$

We remind that within this model  $t$  is the time in the harmonic gauge.

### Acknowledgments

The authors are grateful to M.Rainer for useful discussions.

### References

- [1] K.S. Stelle, "Lectures on Supergravity p-branes ", hep-th/9701088.
- [2] I.Ya. Aref'eva and O.A. Rytchkov, "Incidence Matrix Description of Intersecting p-brane Solutions", hep-th/9612236.
- [3] I.Ya. Aref'eva, M.G. Ivanov and O.A. Rytchkov, "Properties of Intersecting p-branes in Various Dimensions", hep-th/9702077.
- [4] V.D. Ivashchuk and V.N. Melnikov, "Intersecting p-brane Solutions in Multidimensional Gravity and M-theory", hep-th/9612089; *Grav. & Cosmol.* **2**, No 4, 297 (1996).
- [5] V.D. Ivashchuk and V.N. Melnikov, *Phys. Lett. B* **403**, 23 (1997).
- [6] H. Lü, S. Mukherji, C.N. Pope and K.-W. Xu, "Cosmological Solutions in String Theories", hep-th/9610107.
- [7] H. Lü, C.N. Pope, and K.W. Xu, "Liouville and Toda Solitons in M-theory", hep-th/9604058.
- [8] M. Cvetič and A.A. Tseytlin, *Nucl. Phys. B* **478**, 181 (1996).
- [9] V.D. Ivashchuk and V.N. Melnikov, "Sigma-model for the Generalized Composite p-branes", hep-th/9705036; *Class. Quantum Grav.* **14**, 3001 (1997).
- [10] V.D. Ivashchuk, V.N. Melnikov and M. Rainer, "Multidimensional  $\sigma$ -models with Composite Electric p-branes", gr-qc/9705005; *Grav. & Cosmol.* **4**, No 2, (1998).
- [11] K.A. Bronnikov, M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov, "Integrable Multidimensional Cosmology for Intersecting p-branes", *Grav. & Cosmol.* **3**, No 2, 105 (1997).
- [12] M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov, "Integrable Multidimensional Quantum Cosmology for Intersecting p-branes", gr-qc/9708031; *Grav. & Cosmol.* **3**, No 3, 240 (1997).
- [13] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnikov, "The Reissner-Nordström Problem for Intersecting Electric and Magnetic p-branes", gr-qc/9710054; *Grav. & Cosmol.* **3**, No 3, 200 (1997).

- [14] K.A.Bronnikov, U.Kasper and M.Rainer. "Intersecting electric and magnetic  $p$ -branes: spherically symmetric solutions", gr-qc/9708058.
- [15] K.A.Bronnikov, "Block-orthogonal Branes Systems", hep-th/9710207
- [16] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, "Non-extremal Intersecting  $p$ -branes in Various Dimensions", hep-th/9702079.
- [17] N. Ohta, "Intersection Rules for Non-extreme  $p$ -branes", hep-th/9702164.
- [18] V.D. Ivashchuk and V.N. Melnikov, "Multidimensional Classical and Quantum Cosmology with Intersecting  $p$ -branes", hep-th/9708157, *J. Math. Phys.* (1998).
- [19] V.D.Ivashchuk and V.N.Melnikov, "Madjumdar-Papaptrou Type Solutions in Sigma Model and Intersecting  $p$ -branes", hep-th/9802121.
- [20] V.D.Ivashchuk,S.-W.Kim and V.N.Melnikov, "Hyperbolic Kac-Moody Algebra from Intersecting  $p$ -branes", hep-th/9803006.
- [21] M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov, "Multidimensional Cosmology for Intersecting  $p$ -branes with Static Internal Spaces", gr-qc/98040. *Grav. & Cosmol.* **4**, No 2 (1998).
- [22] M.J. Duff, R.R. Khuri and J.X. Lu, *Phys. Rep.* **259**, 213 (1995).
- [23] N. Khvengia, Z. Khvengia, H. Lü and C.N. Pope, "Toward Field Theory of F-Theory", hep-th/9703012.
- [24] J.M. Schwarz, "Lectures on Superstring and M-theory Dualities", hep-th/9607201.
- [25] M.J. Duff, "M-theory (the Theory Formerly Known as Strings)", hep-th/9608117.
- [26] C. Hull and P. Townsend, *Nucl. Phys. B* **438**, 109 (1995);
- [27] P. Horava and E. Witten, *Nucl. Phys. B* **460**, 506 (1996).
- [28] C. Vafa, *Nucl. Phys. B* **469**, 403 (1996).
- [29] E. Cremmer, B. Julia and J. Scherk, *Phys. Lett. B* **76**, 409 (1978).
- [30] M. Rainer and A. Zhuk, *Phys. Rev. D* **54**, 6186 (1996).
- [31] M. Rainer," Multidimensional Scalar-Tensor-Theories and Minisuperspace Approach", Proc. Int. Workshop on Modern Modified Theories of Gravitation and Cosmology (Beer Sheva, June 29-30,1997), publ. in: *Hadronic Journal* **21**, 351 (1998).
- [32] V.D. Ivashchuk and V.N. Melnikov, *Int. J. Mod. Phys. D* **3**, No 4, 795 (1994).
- [33] U. Bleyer, V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, *Nucl. Phys. B* **429**, 177 (1994).
- [34] V.R. Gavrillov, V.D. Ivashchuk and V.N. Melnikov, *J. Math. Phys.* **36**, 5829 (1995).
- [35] Anderson A *J. Math. Phys.* **37**, 1349 (1996).