

Energy-momentum tensor of quasiparticles in the effective gravity in superfluids.

G.E. Volovik

Low Temperature Laboratory, Helsinki University of Technology
P.O.Box 2200, FIN-02015 HUT, Finland

and

L.D. Landau Institute for Theoretical Physics, Moscow

December 2, 2024

Abstract

The problem of the energy-momentum conservation for matter in the gravitational field is discussed on the example of the effective gravity, which arises in superfluids. The "gravitational" field experienced by the relativistic-like massless quasiparticles which form the "matter" (phonons in superfluid 4He and low-energy fermions in superfluid 3He-A), is induced by the flow of the superfluid "vacuum". It appears that the energy-momentum conservation law for quasiparticles, has the covariant form $T_{\nu;\mu}^{\mu} = 0$. "Pseudotensor" of the energy-momentum for the "gravitational field" (superfluid condensate) appears to depend on "matter". The inflationary stage of the expansion of the Universe can be modelled using the expanding Bose-condensate.

Introduction. The effective gravity of the Sakharov type [1] can occur in many condensed matter systems. This allows us to use these systems for modelling the specific properties of the general relativity [2, 3, 4, 5, 6, 7]. The important class of the effective gravity theories, which naturally arises in condensed matter, is that which is induced by the gapless (massless) quasiparticles. An example is the effective gravity in superfluid $^3\text{He-A}$ [7]. The energy spectrum of fermionic quasiparticles in this superfluid contains

the topologically stable point nodes (Fermi points) [8]. In the vicinity of the gap node the quasiparticle spectrum becomes fully relativistic

$$g^{\mu\nu}(p_\mu - eA_\mu)(p_\nu - eA_\nu) = 0 . \quad (1)$$

The topological stability of the gap nodes is crucial: the small deformations of the quantum vacuum do not destroy the Fermi points but deform their characteristics, the slopes $g^{\mu\nu}$ and shifts A_μ in Eq.(1), which play the part of the gravitational field and different types of the gauge fields correspondingly. The gravity and gauge fields are thus the natural dynamical collective modes of the fermionic quantum vacuum with Fermi points.

Using such an effective gravity we discuss the particular problems related to the energy-momentum tensor of matter in the gravitational field.

Basic equations. For the energy-momentum problem the quasiparticle energy spectrum only is important, so that the quantum statistics of quasiparticles is irrelevant. So, we can consider the Landau two-fluid model of superfluidity, in which one component of the fluid is formed by free quasiparticles (Bose or Fermi) moving on the background of the superfluid vacuum play the part of matter. The effective gravitational field acting on quasiparticles is produced by the motion of the another component of the liquid – the superfluid condensate, which is characterized by the density ρ and superfluid velocity $\mathbf{v}_{(s)}$. The latter is not necessarily curl-free. Let us recall some basic equations of two-fluid dynamics, using the simplest possible model with the following energy of the liquid [9]:

$$\mathcal{E} = \int d^3r \left(\frac{1}{2} \rho \mathbf{v}_{(s)}^2 + \epsilon(\rho) + \sum_{\mathbf{p}} E(\mathbf{p}, \mathbf{r}) f(\mathbf{p}, \mathbf{r}) \right) . \quad (2)$$

Here the first term is the kinetic energy of the superfluid condensate; the second term is its energy density as a function of ρ ; $f(\mathbf{p}, \mathbf{r})$ is the distribution function of quasiparticles; and the quasiparticle energy spectrum is:

$$E(\mathbf{p}, \mathbf{r}) = E_0(\mathbf{p}, \rho(\mathbf{r})) + \mathbf{p} \cdot \mathbf{v}_{(s)}(\mathbf{r}) . \quad (3)$$

where the second term is the Doppler shift in the moving condensate, and we assumed that the quasiparticle energy $E_0(\mathbf{p}, \mathbf{r})$ in the superfluid frame (in the frame where the superfluid vacuum is at rest) depends on the coordinate only through the density ρ .

The mass current density or the linear momentum density consists of the vacuum current and the momentum of quasiparticles

$$\mathbf{j} = \rho \mathbf{v}_{(s)} + \mathbf{P} , \quad \mathbf{P} = \sum_{\mathbf{p}} \mathbf{p} f(\mathbf{p}, \mathbf{r}) , \quad (4)$$

so that the continuity equation is

$$\dot{\rho} + \vec{\nabla}(\rho \mathbf{v}_{(s)} + \mathbf{P}) = 0 . \quad (5)$$

The superfluid velocity of the vacuum obeys the London equation:

$$\dot{\mathbf{v}}_{(s)} + \vec{\nabla} \frac{\delta \mathcal{E}}{\delta \rho} - \frac{\mathbf{j}}{\rho} \times (\vec{\nabla} \times \mathbf{v}_{(s)}) = 0 . \quad (6)$$

In superfluid ^4He the superfluid velocity is potential, so that vorticity $\vec{\nabla} \times \mathbf{v}_{(s)}$ is nonzero only in the presence of quantized vortices. In superfluid $^3\text{He-A}$ the vorticity can be continuous and the London equation in this form can be obtained if one neglects the axial anomaly and the dependence of the energy on the anisotropy vector (see Eqs.(6.6)-(6.7) in Ref. [8]). The last term in equation (6) states that in the absence of anomalies the vorticity moves with the center-of-mass velocity \mathbf{j}/ρ .

The distribution function f of the quasiparticles is determined by the kinetic equation:

$$\dot{f} - \frac{\partial E}{\partial \mathbf{r}} \cdot \frac{\partial f}{\partial \mathbf{p}} + \frac{\partial E}{\partial \mathbf{p}} \cdot \frac{\partial f}{\partial \mathbf{r}} = J_{coll} . \quad (7)$$

Momentum conservation. From above equations one obtains the time evolution of the momentum density for each of two subsystems: the superfluid background (vacuum) and quasiparticles (matter). The momentum evolution of the superfluid vacuum is

$$\partial_t(\rho \mathbf{v}_{(s)}) = -\nabla_i(j_i \mathbf{v}_{(s)}) - \rho \vec{\nabla} \left(\frac{\partial \epsilon}{\partial \rho} + \sum_{\mathbf{p}} f \frac{\partial E_0}{\partial \rho} \right) + P_i \vec{\nabla} v_{(s)i} . \quad (8)$$

Since the collision integral conserves the momentum of quasiparticles, the evolution of the momentum density of quasiparticles, as obtained from the kinetic equation, is:

$$\partial_t \mathbf{P} = -\nabla_i(v_{(s)i} \mathbf{P}) - \nabla_i \left(\sum_{\mathbf{p}} \mathbf{p} f \frac{\partial E_0}{\partial p_i} \right) - \sum_{\mathbf{p}} f \vec{\nabla} E_0 - P_i \vec{\nabla} v_{(s)i} . \quad (9)$$

Though the momentum of each subsystem is not conserved because of the interaction with the other subsystem, the total momentum is conserved:

$$\partial_t j_i = \partial_t (\rho v_{(s)i} + P_i) = -\nabla_i P_{ik} , \quad (10)$$

with the stress tensor

$$P_{ik} = j_i v_{(s)k} + v_{(s)i} P_k + \sum_{\mathbf{p}} p_k f \frac{\partial E_0}{\partial p_i} + \delta_{ik} G , \quad G = \rho \left(\frac{\partial \epsilon}{\partial \rho} + \sum_{\mathbf{p}} f \frac{\partial E_0}{\partial \rho} \right) - \epsilon . \quad (11)$$

"Relativistic" quasiparticles . Let us consider the "relativistic" quasiparticles – phonons in superfluid ^4He or fermionic quasiparticles in superfluid ^3He . Their energy spectrum is linear in the limit of low energy:

$$E(\mathbf{p}, \mathbf{r}) = cp + \mathbf{p} \cdot \mathbf{v}_{(s)} , \quad \text{or} \quad (E - \mathbf{p} \cdot \mathbf{v}_{(s)})^2 - c^2 p^2 = 0 . \quad (12)$$

This equation is correct for phonons in superfluid ^4He , with c being the speed of sound. In the $^3\text{He-A}$ the low energy spectrum is also linear but anisotropic (see [8]). The simplified equation (12) is valid in $^3\text{He-A}$ only for the homogeneous order parameter and is obtained after shifting of the momentum and rescaling. The function c in this case is some combination of the slopes of the energy spectrum.

Effective metric. Since the energy spectrum has the relativistic form, let us rewrite the time evolution of the quasiparticle momentum density in Eq.(9) in the covariant form, using the superfluid system as the effective background metric. The corresponding metric follows from the quasiparticle spectrum in Eq.(12), which can be written in a general Lorentzian form

$$g^{\mu\nu} p_\mu p_\nu = 0 . \quad (13)$$

Here $p_0 = -E$, and the metric elements are

$$g^{00} = 1 , \quad g^{0i} = v_s^i , \quad g^{ik} = -(c^2 \delta^{ik} - v_{(s)i}^i v_{(s)k}^k) . \quad (14)$$

The metric of the effective space in which quasiparticles propagate along the geodesics, i.e. $ds^2 = g_{\mu\nu} dx^\mu dx^\nu = 0$, is correspondingly

$$g_{00} = \left(1 - \frac{v_{(s)}^2}{c^2} \right) , \quad g_{0i} = \frac{v_{(s)i}}{c^2} , \quad g_{ik} = -\frac{1}{c^2} \delta_{ik} . \quad (15)$$

Except for the conformal factor, it coincides with acoustic metric discussed for sound propagation in normal [2, 3] and superfluid [10] liquids. In our case the quasiparticles are not necessarily the phonons.

The system of quasiparticles play the part of the matter in general relativity. That is why one can expect that the momentum "conservation law" for matter has the usual covariant form

$$T_{\nu;\mu}^{\mu} = 0 \ , \quad \text{or} \quad \frac{1}{\sqrt{-g}} \partial_{\mu} \left(T_{\nu}^{\mu} \sqrt{-g} \right) - \frac{1}{2} T^{\alpha\beta} \partial_{\nu} g_{\alpha\beta} = 0 \ , \quad (16)$$

where $\sqrt{-g} = c^{-3}$. The terms which are not the total derivatives represent the forces acting on the matter from the effective gravitational field. Let us check if this equation can be applied to our case, where the effective gravity is induced by the superflow.

Energy-momentum tensor.

Let us introduce the components of the energy-momentum tensor:

$$\begin{aligned} \sqrt{-g} T_i^0 &= -P_i \ , \quad \sqrt{-g} T_0^0 = \sum_{\mathbf{p}} f E \ , \quad \sqrt{-g} T_i^k = - \sum_{\mathbf{p}} p_i f \frac{\partial E}{\partial p_k} \ , \\ \sqrt{-g} T_0^i &= P^i c^2 + v_{(s)}^i \sum_{\mathbf{p}} f E_0 + v_{(s)}^k \sum_{\mathbf{p}} p_k f \frac{\partial E}{\partial p_i} \ . \end{aligned} \quad (17)$$

$$\begin{aligned} \sqrt{-g} T^{00} &= \sum_{\mathbf{p}} f E_0 \ , \quad \sqrt{-g} T^{0i} = c^2 P^i + v_{(s)}^i \sum_{\mathbf{p}} f E_0 \ , \\ \sqrt{-g} T^{ik} &= c^2 \sum_{\mathbf{p}} p_i f \frac{\partial E_0}{\partial p_k} + c^2 (v_{(s)}^k P^i + v_{(s)}^i P^k) + v_{(s)}^i v_{(s)}^k \sum_{\mathbf{p}} f E_0 \ . \end{aligned} \quad (18)$$

Using the above expression for the quasiparticle energy-momentum tensor, one can be rewrite the equation for the quasiparticle momentum density, Eq.(9), and the analogous equation for the evolution of quasiparticles energy density in the following form:

$$\partial_{\mu} \left(\sqrt{-g} T_{\nu}^{\mu} \right) - \mathbf{P} \cdot \partial_{\nu} \mathbf{v}_{(s)} - \frac{\partial_{\nu} c}{c} \sum_{\mathbf{p}} f E_0 = 0 \ . \quad (19)$$

The last two terms are the forces acting on the subsystem of quasiparticles from the superfluid background, if the latter is inhomogeneous. These forces

exactly reproduce the forces acting on the matter from the gravitational field in Eq.(16):

$$\frac{1}{2}\sqrt{-g}T^{\alpha\beta}\partial_\nu g_{\alpha\beta} = \mathbf{P} \cdot \partial_\nu \mathbf{v}_{(s)} + \frac{\partial_\nu c}{c} \sum_{\mathbf{p}} f E_0 . \quad (20)$$

Thus the energy-momentum tensor of quasiparticles satisfies the covariant equation (16). This result does not depend on the dynamic properties of the superfluid condensate (gravity field) and follows only from the "relativistic" spectrum of quasiparticles.

Thermal equilibrium. In thermal equilibrium the quasiparticle distribution function is $f = (\exp(E(\mathbf{p}, \mathbf{r}) - \mathbf{p} \cdot \mathbf{v}_{(n)})/T \pm 1)^{-1}$. It is determined by the temperature T and the normal component velocity $\mathbf{v}_{(n)}$, which reflect the conservation of correspondingly the energy and momentum by the collision term in kinetic equation. The Doppler shifted energy is $E(\mathbf{p}, \mathbf{r}) - \mathbf{p} \cdot \mathbf{v}_{(n)} = cp - \mathbf{p} \cdot \mathbf{w}$, where $\mathbf{w} = \mathbf{v}_{(n)} - \mathbf{v}_{(s)}$ is the counterflow velocity of the normal component with respect to the superfluid one.

The quasiparticle momentum density in thermal equilibrium is

$$\mathbf{P} = \sum_{\mathbf{p}} \mathbf{p} f(\mathbf{p}, \mathbf{r}) = \rho_n^{(0)} \frac{\mathbf{w}}{\left(1 - \frac{w^2}{c^2}\right)^3} , \quad (21)$$

where $\rho_n^{(0)}$ is normal fluid density in the limit of zero counterflow, $\mathbf{w} = 0$. For the spin-1/2 fermions

$$\rho_n^{(0)} = \frac{7}{90} \pi^2 \frac{T^4}{c^5} = \frac{7}{90} \pi^2 \frac{T^4}{c^2} \sqrt{-g} . \quad (22)$$

The quasiparticle energy density is

$$\sum_{\mathbf{p}} E_0 f = \frac{\varepsilon^{(q)} + P^{(q)}}{1 - \frac{w^2}{c^2}} - P^{(q)} = \frac{3}{4} \rho_n^{(0)} c^2 \frac{1 + \frac{1}{3} \frac{w^2}{c^2}}{\left(1 - \frac{w^2}{c^2}\right)^3} , \quad (23)$$

where $P^{(q)}$ is the quasiparticle contribution to pressure

$$P^{(q)} = \frac{1}{4} \frac{\rho_n^{(0)} c^2}{\left(1 - \frac{w^2}{c^2}\right)^2} , \quad \frac{\partial P^{(q)}}{\partial \mathbf{v}_n} = \frac{\partial P^{(q)}}{\partial \mathbf{w}} = \mathbf{P} . \quad (24)$$

The quasiparticle contribution to the stress tensor is

$$\sum_{\mathbf{p}} p_k f \frac{\partial E_0}{\partial p_i} = (\varepsilon^{(q)} + P^{(q)}) \frac{w_i w_k}{c^2 - w^2} + P^{(q)} \delta_k^i . \quad (25)$$

where $\varepsilon^{(q)} = 3P^{(q)} = \rho^{(q)}c^2$ is the energy density of "matter" and $\rho^{(q)}$ is its "mass".

The energy-momentum tensor for quasiparticles has the following general form

$$T^{\mu\nu} = (\varepsilon + P)u^\mu u^\nu - Pg^{\mu\nu} . \quad (26)$$

Here u^α and $u_\alpha = g_{\alpha\beta}u^\beta$ are the 4-velocities of the "matter" which satisfy equation $u_\alpha u^\alpha = 1$:

$$u^0 = \frac{1}{\sqrt{1 - \frac{w^2}{c^2}}} , \quad u^i = \frac{v_{(n)}^i}{\sqrt{1 - \frac{w^2}{c^2}}} , \quad u_i = -\frac{w_i}{c^2 \sqrt{1 - \frac{w^2}{c^2}}} , \quad u_0 = \frac{1 + \frac{\mathbf{w} \cdot \mathbf{v}_{(s)}}{c^2}}{\sqrt{1 - \frac{w^2}{c^2}}} , \quad (27)$$

and ε is the "covariant" energy density of quasiparticles: $\varepsilon = \varepsilon^{(q)}/\sqrt{-g}$. For the 1/2-spin fermionic quasiparticles one has

$$\varepsilon = \frac{\varepsilon^{(q)}}{\sqrt{-g}} = \frac{7}{120}\pi^2 \frac{T^4}{\left(1 - \frac{w^2}{c^2}\right)^2} , \quad P = \frac{\varepsilon}{3} . \quad (28)$$

This corresponds to the effective local temperature

$$T_{\text{eff}} = \frac{T}{\sqrt{\tilde{g}_{00}}} , \quad \tilde{g}_{00} = g_{00}(\mathbf{v}_{(n)} = 0) = 1 - \frac{w^2}{c^2} , \quad (29)$$

where \tilde{g}_{00} is the effective metric in the frame comoving with the normal component (the local frame, where $\mathbf{v}_{(n)} = 0$ and $\mathbf{w} = -\mathbf{v}_{(s)}$).

The components of the energy-momentum tensor are

$$\begin{aligned} T_i^0 &= -(\varepsilon + P)\frac{w_i}{c^2 - w^2} , \quad T_i^k = -(\varepsilon + P)\frac{v_{(n)}^k w_i}{c^2 - w^2} - P\delta_i^k , \\ T_0^i &= (\varepsilon + P)v_{(n)}^i \frac{c^2 + \mathbf{w} \cdot \mathbf{v}_{(s)}}{c^2 - w^2} , \quad T^{00} = \frac{(\varepsilon + P)}{1 - w^2/c^2} - P , \quad T_0^0 = T^{00} + \frac{\mathbf{P} \cdot \mathbf{v}_{(s)}}{\sqrt{-g}} . \end{aligned} \quad (30)$$

The energy-momentum pseudotensor.

The equation for the energy-momentum of quasiparticles is covariant but this equation does not represent any conservation law. On the contrary, the total energy and momentum of the system, quasiparticles + superfluid condensate (gravity field), are conserved, but this conservation laws cannot be expressed in the covariant form. This is another problem of the general

relativity, which probably also can be attacked using the condensed matter analogy. In the condensed matter the energy-momentum tensor of the system exists on the microscopic level, i.e. it can be expressed using the microscopic variables. But in many cases, for example in ferromagnets [11], it cannot be expressed in terms of the local effective variables, such as magnetization. The effective gravity field as the local low-energy collective mode can be just an another example.

Let us consider the analog of the pseudotensor of the energy-momentum of the gravitational field, which is in our case the part of the total energy momentum tensor coming from the superfluid motion of the condensate. The conservation law for the total energy-momentum can be written as

$$\partial_\mu (\sqrt{-g}T_\nu^\mu + \sqrt{-g}t_\nu^\mu) = 0 , \quad (31)$$

where t_ν^μ is the pseudotensor of energy-momentum. From the basic equations for superfluids it follows that

$$\begin{aligned} \sqrt{-g}t_i^0 &= -\rho v_{(s)i} , \\ \sqrt{-g}t_i^k &= -j^k v_{(s)i} - \delta_i^k \left[\rho \left(\frac{\partial \epsilon}{\partial \rho} + \sum_{\mathbf{p}} f \frac{\partial E_0}{\partial \rho} \right) - \epsilon \right] , \\ \sqrt{-g}t_0^0 &= \frac{1}{2} \rho \mathbf{v}_{(s)}^2 + \epsilon(\rho) , \\ \sqrt{-g}t_0^i &= j^i \left(\frac{\partial \epsilon}{\partial \rho} + \sum_{\mathbf{p}} f \frac{\partial E_0}{\partial \rho} + \frac{\mathbf{v}_{(s)}^2}{2} \right) . \end{aligned} \quad (32)$$

It appears, that the pseudotensor of the "gravitational field" depends on matter.

Let us introduce the Lagrangian

$$L = \frac{1}{2} \rho \mathbf{v}_{(s)}^2 + \epsilon(\rho) + \sum_{\mathbf{p}} E f + \rho \partial_t \Phi . \quad (33)$$

where Φ is the phase of the condensate related to the superfluid velocity:

$$\vec{\nabla} \Phi = \frac{m}{\hbar} \mathbf{v}_{(s)} . \quad (34)$$

Here m is the mass of the initial boson in the superfluid Bose condensate, and further we put $\hbar/m = 1$. Then the above pseudotensor of the "gravitational

field" can be written in the compact form:

$$\sqrt{-g}t_\nu^\mu = L\delta_\nu^\mu - \frac{\partial L}{\partial \nabla_\mu \Phi} \nabla_\nu \Phi - \delta_\nu^\mu \sum_{\mathbf{p}} E f . \quad (35)$$

Hubble expansion.

The expanding solution of the motion equations is

$$\mathbf{v}_{(s)}(\mathbf{r}, t) = \mathbf{v}_{(n)}(\mathbf{r}, t) = \tilde{H}(t)\mathbf{r} , \quad \rho = \rho(t) . \quad (36)$$

The Hubble parameter and its connection to the density of the liquid are found from Eqs.(5,6):

$$\tilde{H}(t) = -\frac{\partial_t \rho}{3\rho} , \quad \partial_t \tilde{H} + \tilde{H}^2 = 0 . \quad (37)$$

which give

$$\tilde{H}(t) = (t - t_0)^{-1} , \quad \rho(t) = \text{Const } (t - t_0)^{-3} \quad (38)$$

Note that the Hubble parameter in the radiation-dominated Universe is $H(t) = (1/2)(t - t_0)^{-1} = (1/2)\tilde{H}(t)$.

Since the quasiparticle momentum $\mathbf{P} = 0$ (because $\mathbf{v}_{(s)} = \mathbf{v}_{(n)}$), the Eq.(9) is automatically satisfied, while the energy conservation law for quasiparticles gives the following equation for the quasiparticles energy density $\varepsilon^{(q)} = 3P^{(q)} (= \rho^{(q)} c^2)$:

$$\partial_t \varepsilon^{(q)} + \left(4\tilde{H} - \frac{\partial_t c}{c} \right) \varepsilon^{(q)} = 0 . \quad (39)$$

Since $\varepsilon^{(q)} = \varepsilon/c^3 \propto T^4/c^3$, one obtains equation for T :

$$\partial_t \left(\frac{T}{c} \right) + \tilde{H} \left(\frac{T}{c} \right) = 0 . \quad (40)$$

which gives $T(t) \propto c(t)/(t - t_0)$.

The time dependence of the speed of light is determined by its dependence on the density ρ : $c(t) = c(\rho(t))$. The cosmological scenario of the expansion of the radiation-dominated Universe, with $T(t) \propto (t - t_0)^{-1/2}$, is obtained if $c \propto (t - t_0)^{1/2}$. If the speed c comes from the compressibility of the superfluid vacuum, then such behavior of c can result from the following

equation of states for the superfluid condensate: $\epsilon(\rho) \sim \rho^{2/3}$ which gives $c(\rho) \sim \rho^{-1/6} \propto (t - t_0)^{1/2}$. Note that in the cases of the Bose condensation of the almost ideal Bose gas and of the superfluidity of the almost ideal Fermi gas, the equation of states are correspondingly $\epsilon(\rho) \sim \rho^2$ and $\epsilon(\rho) \sim \rho^{5/3}$ with $c(\rho) \sim \rho^{1/2} \propto (t - t_0)^{-3/2}$ and $c(\rho) \sim \rho^{1/3} \propto (t - t_0)^{-1}$.

If one introduces the comoving coordinate frame: $\mathbf{r} = \tilde{\mathbf{r}}\tilde{a}(t)$ with $\partial_t \tilde{a} = \tilde{a}\tilde{H}$, one obtains the following effective metric in this frame

$$ds^2 = dt^2 - a^2 d\tilde{\mathbf{r}}^2, \quad a = \frac{\tilde{a}}{c}, \quad \frac{\partial_t \tilde{a}}{\tilde{a}} = \tilde{H}. \quad (41)$$

Since $\tilde{a} \propto t - t_0$, the behavior corresponding to the radiation-dominated Universe occurs if $c \propto (t - t_0)^{1/2}$. Then $a = \tilde{a}/c \propto (t - t_0)^{1/2}$ and $\partial_t a/a = H = (1/2)\tilde{H}$. Such case occurs if $\epsilon(\rho) \sim \rho^{2/3}$.

Another special case is when $c \propto t - t_0$. In this case $a = \text{Const}$, and $T = \text{Const}$. This means that the effective space-time for quasiparticles is Minkowski flat space-time, inspite of the nontrivial dynamics of the background superfluid condensate. Such case occurs if $\epsilon(\rho) \sim \rho^{1/3}$ which gives $c(\rho) \sim \rho^{-1/3} \propto t - t_0$.

In the case of the superfluidity of the almost ideal Bose and Fermi gases, the integral $\int dt a^{-1}(t)$ diverges at the origin. Thus both cases correspond to the power-law inflation.

The next step is to derive the effective action for the "gravity field". The effective gravity theory is obtained by integration over the fermionic degrees of freedom in the presence of the background field [1]. The latter field becomes dynamical and represents the low-frequency collective modes of the system of interacting particles – ^3He or ^4He atoms. In the case discussed here the result is well known: the hydrodynamic equations for the background superfluid vacuum in Eqs.(5,6), which do not depend on the details of the microscopic interactions between the atoms, but which are far from being covariant and relativistic. To have the relativistic equations for the effective gravity field one must consider such hypothetical condensed mater systems where the main contribution to the path integral comes from the "relativistic" quasiparticles. If in addition these fermions are "massless", there is an extra symmetry: with respect to the multiplication of the metric by an arbitrary function $g_{\mu\nu} \rightarrow a^2 g_{\mu\nu}$. Such conformal symmetry of quasiparticles should lead to the conformal invariant and covariant action for the superfluid condensate, which contains the Weyl tensor (see e.g. [12]).

I am grateful to Alexei Morozov and Alexei Starobinsky for discussions after which the error in the previous version of the manuscript became evident.

References

- [1] A.D. Sakharov, Sov. Phys. Doklady **12**, 1040 (1968).
- [2] W.G. Unruh, Phys. Rev. Lett. **46**, 1351 (1981).
- [3] M. Visser, Class. Quant. Grav. **15**, 1767 (1998).
- [4] M.O. Katanaev and I.V. Volovich, "Scattering on Dislocations and Cosmic Strings in the Geometric Theory of Defects", gr-qc/9801081
- [5] R. Bausch and R. Schmitz, Phys. Rev. Lett., **80**, 2257 (1998).
- [6] T. Jacobson and G.E. Volovik, Phys. Rev. **D 58**, 064021(1-7) (1998).
- [7] G.E. Volovik, "Induced Gravity in Superfluid ^3He ", cond-mat/9806010.
- [8] G. E. Volovik, "*Exotic properties of superfluid ^3He* ", World Scientific, Singapore - New Jersey - London - Hong Kong, 1992.
- [9] I.M. Khalatnikov, "An Introduction to the Theory of Superfluidity", (Benjamin, New York), 1965.
- [10] K.N.Ilinski and A.S.Stepanenko, "From Bose condensation to quantum gravity and back", cond-mat/9803233.
- [11] G.E. Volovik, J. Phys. C **20**, L83 (1987).
- [12] N.D. Birell and P.C.W. Davies, Quantum Fields in Curved Space (Cambridge University Press, Cambridge, 1982).