Domain walls and torsion potentials.

L.C. Garcia de Andrade¹

Abstract

A Lagrangian for flat domain walls in spaces with Cartan torsion and electromagnetic fields is proposed.The Lagrangian is very similar to a recently proposed Lagrangian for domain walls in a Chern-Simons electrodynamics in 2+1 dimensions.We show that in the first approximation of the torsion scalar potential the field equations are reduced to a Klein-Gordon type field equation for the torsion potential and the electromagnetic wave equation.A planar symmetric solution representing a parallel plates electric capacitor interacting with the electric field is given.The photon mass is proportional to the torsion potential and in the time dependent case the angular momentum is computed and is shown to be connected with torsion in analogy with the spin-torsion relation which appears in Einstein-Cartan gravity.When the curvature Ricci scalar is introduced we are able to show that the torsion potential can be associated with Higgs massive vectorial bosons.

¹Departamento de Física Teorica - UERJ. Rua São Fco. Xavier 524, Rio de Janeiro, RJ Maracanã, CEP:20550-003, Brasil. E-Mail.: GARCIA@SYMBCOMP.UERJ.BR

Recently vortices and domain walls in a Chern-Simons (CS) theory with magnetic moment interactions were investigated by Antillon, Escalona and Torres $[1]$. In their paper they work in $2+1$ dimensional spacetime. Here following recently investigation on non-Riemannian domain walls in space-times with torsion $[2, 3, 4]$ $[2, 3, 4]$ we worked out avery similar Lagrangian like the one of reference one with the exception that dual field in the (CS) Lagrangian is replaced by torsion and we work in the full four-dimensional spacetime.The equations obtained for the torsion potential and the electromagnetic field are reduced to non-linear Klein-Gordon field equation and to the Proca field equation in the case of the electromagnetic vector potetencial.The controversy concerning the interaction of the electromagnetic fields and torsion [[5](#page-4-0), [6,](#page-4-0) [7\]](#page-5-0) here reachs a slightly different approach since the interaction between photons and torsion appears intermediated by a spin-0 massive boson given by the torsion potential ,besides in the first approximation of the torsion potential the Proca equation is reduced to the Maxwell equation and the photon is massless and the theory is gauge invariant in first approximation.As an application we solve the field equations in the first order in the torsion potential for the case of the parallel plates Capacitor.This simple example is analogous to the case of the Casimir effect [\[8](#page-5-0)] with torsion to be investigated in a future research.Of course throughout the paper the metric is considered to be flat.Let us assume that our Lagrangian is given by

$$
L = -\frac{1}{4}F^2 + \frac{1}{2}|D_{\mu}\phi|^2 - V(\phi)
$$
 (1)

where the bars denotes the modulus since we considering that the torsion potential ϕ is real but the covariant derivative defined by

$$
D_{\mu} = \partial_{\mu} - if A_{\mu} - ig S_{\mu} \tag{2}
$$

is complex.Here $F^2 = F_{\mu\nu}F^{\mu\nu}$ is the electromagnetic invariant and $F_{\mu\nu} =$ $\partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ is the electromagnetic field tensor. The torsion potential generates the torsion vector through the relation $S_{\mu} = \partial_{\mu} \phi$. Substitution of the definition (2) into the Lagrangian (1) reduces the previous Lagrangian to

$$
L = \frac{1}{2}(\partial_{\mu}\phi)^{2} - \frac{1}{4}F^{2} - \frac{1}{2}(f^{2}A^{2} + g^{2}S^{2})\phi^{2} - V(\phi)
$$
 (3)

or

$$
L = \frac{1}{2}(1+f^2)(\partial\phi)^2 + \frac{1}{2}g^2A^2\phi^2 - \frac{1}{4}F^2 - V(\phi)
$$
 (4)

Variation of this Lagrangian with respect to torsion potential and the electromagnetic potential vector A_μ yields the following non-linear field equations

$$
\nabla^2 A^{\mu} + \frac{1}{2} g^2 \phi^2 A^{\mu} = 0
$$
\n(5)

and

$$
\partial^2 \phi - \frac{2f^2 \phi}{(1+f^2 \phi^2)} (\partial \phi)^2 = \frac{g^2 A^2 \phi}{1+f^2 \phi^2} \tag{6}
$$

where we have supressed the indices and have already considered the domain wall potential $V(\phi)$ as zero since we will not need it in our next application.Notice that the equation (5) yields a mass for the photon like $m_{\gamma} = g\phi$. Thus the photon mass can be expressed in terms of the torsion potential.This situation is similar to the photon mass dependence on torsion developed some years ago by Sivaram and myself[[9\]](#page-5-0).One should notice that equation([4\)](#page-1-0) is the Proca equation for the massive photon while the second is a non-linear field equation which reduces to the Klein-Gordon type field equation in the case that the terms quadratic in the torsion potential are dropped out.In this case also we note that the Proca equation is reduced to the massless wave equation which in the time independent case reads

$$
\nabla^2 A^\mu = 0\tag{7}
$$

As an example let us solve these equations inthe static case of a parallel plate capacitor interacting with the scalar torsion field. Since the capacitor electrostatic potential is given by $\phi^{el} = qd$ where d is the separation between the plates and q is the electric charge.Here we have taken the Parallel plate orthogonal to the z-coordinate axis,the Klein-Gordon-Cartan equation is

$$
\phi'' - g^2 d^2 q^2 \phi = 0 \tag{8}
$$

where the double lines denote second derivative in z . To solve this Klein-Gordon field equation for torsion potential,let us use the following ansatz

$$
\phi = e^{-\alpha z} \tag{9}
$$

Since we assume that the torsion field vanishes at infinity. Substitution of (9) into (8) yields

$$
\alpha = gqd \tag{10}
$$

which yields the following solution for the torsion potential $\phi = e^{-gqdz}$ which displays the coupling between the electric charge and the torsion coupling constant.In the time independent case we have an expression for the electromagnetic vector potential in terms of the scalar field as

$$
A^{0} = \frac{(\nabla . S)^{\frac{1}{2}} (1 + f^{2} \phi^{2})^{\frac{1}{2}}}{g\phi}
$$
 (11)

From this expression it is very easy to compute the electrostatic field as

$$
E = -\nabla A^0 = -(\nabla . S)^{\frac{1}{2}} \frac{(1 - \frac{1}{2}\phi)}{g\phi^2}
$$
(12)

where to simplify matters we have considered that the divergence of the torsion is constant.In the static case the energy can be computed from the Lagrangian above as

$$
T_{\mu\nu} = (1 + f^2)\partial_{\mu}\phi \partial_{\nu}\phi - \eta_{\mu\nu}L \tag{13}
$$

The energy is

$$
\epsilon = \frac{1}{4}(E^2 - B^2) - \frac{1}{2}g^2 A^2 \phi^2 \tag{14}
$$

In the time dependent case a relation between the angular momentum of the scalar field and torsion is obtained by using the definition of the angular momentum

$$
J_{ij} = \int (T_{0i}x_j - T_{0j}x_i)d^4x \tag{15}
$$

By making use of equation (13) and the definition of the torsion as the gradient of the scalar field yields

$$
J_{ij} = (1 + f^2) \int S_0 (S_i x_j - S_j x_i) d^4 x \tag{16}
$$

One must notice that this expression is similar to the relation between spin and torsion which appears in Einstein-Cartan gravity and to the 2+1 Kerr solution discovered by Jackiw [\[11](#page-5-0)]. It is interesting to note that when the curvature scalar is introduced into the Lagrangian one is able to show that the torsion potential is in fact a massive Higgs field.This can be show by considering the following lagrangian

$$
L = R(S) - m^2 \phi^2 - V(\phi)
$$
 (17)

where in considering just the flat metric terms and remiding that the Ricci scalar is given by $R(S) = \partial S + S^2$ where S represents the torsion vector, the Lagrangian [\(13\)](#page-3-0) reduces to

$$
L = \partial^2 \phi - (\partial \phi)^2 - m^2 \phi^2 - V(\phi)
$$
 (18)

The Euler-Lagrange equation for this Lagrangian yields

$$
\partial_{\mu}\partial^{\mu}\phi + m^2\phi = -\frac{\partial V}{\partial \phi} \tag{19}
$$

Which is the equation for a massive Higgs field given by the torsion potential.Recently in a similar approach Kleinert [\[10\]](#page-5-0) discussed the spontaneous symmetry breaking for the electroweak bosons induced by the torsion potential,nevertheless in his case the Higgs potential did not coincide with the torsion potential.A more detailed investigation of our model as well a investigation of the Casimir effect in spacetimes with torsion may appear elsewhere.

Acknowledgements

I am very much indebt to Professor A.Wang,I.D. Soares and H.Kleinert for helpful discussions on the subject of this paper. Thanks are also due to Universidade do Estado do Rio de Janeiro (UERJ) for financial Support.

References

- [1] A.Antillon,J.Escalona and M.Torres,Phys.Rev.D55,10,(1997),6327.
- [2] L.C.Garcia de Andrade,J.Math.Phys.32,(1998)372.
- [3] L.C.Garcia de Andrade,Gen.Rel.and Grav.30,11,(1998)1629.
- [4] L.C.Garcia de Andrade,Mod.Phys.Lett.A27,(1997),2005.
- [5] A.Unzicker,Teleparallel Space-time with Defects yields Geometrization of Electrodynamics with quantized charges,Los Alamos Electronic archives[,gr-qc/9612061.](http://arxiv.org/abs/gr-qc/9612061)
- [6] L.C.Garcia de Andrade,Gen.Rel.and Grav,(1990),8,882.
- [7] R.A.Puntigam,C.Lammerzhal and F.W.Hehl,Class.and Quantum Grav.14,5(1997),1347.
- [8] Mostepanenko,The Casimir Effect.
- [9] L.C.Garcia de Andrade and C.Sivaram,Spontaneous Symmetry Breaking in massive electrodynamics induced by torsion,dft-UERJ internal reports(1992).
- [10] H.Kleinert,Spontaneous Generation of Torsion Coupling of Electroweak Massive Gauge Bosons,(1998),Institut fur theoretische Physik,Freie University Berlin.
- [11] R.Jackiw in SILARG VII-Relativity and Gravitation:Classical and Quantum,(1990),Simposio Latino Americano de Relatividade Geral e Gravitacion,World Scientific.