

 P.Z. Aslanyan^{1,2}, V.N. Emelyanenko¹, G.G.Rikhvitzkaya¹

(1) Joint Institute for Nuclear Research
(2) Yerevan State University
† E-mail: petros@spp.jinr.ru

Abstract

In the metastable state sector the Λp , $\Lambda \pi \pi$ and $\Lambda p \pi$ resonances found in $n+{}^{12}C$ collisions at 7 GeV/c. Preliminary results show that the predicted peaks has been confirmed in $p^{12}C$ collision at 10 GeV/c. In the stable state sector candidates for S=-2 light and heavy dibaryons observed in $p+{}^{12}C$ collision at 10 GeV/c.

1 Introduction

Quark models with confinement predict the existence of exotic objects as, for instance, glueballs, hybrid states of quarks and gluons as well as multiquark hadrons [1]- [6]. It is important to stress that the existence of multiquarks is a necessary consequence of modern ideas of non-perturbative QCD vacuum and is a matter of principle to comprehend the characteristics of matter both at microscopic and cosmological levels. New particles or states of matter containing 2-or more strange quarks have inspired a lot of experiments at BNL(AGS), CERN, FNAL, SEBAF, KEK and JINR.

Stable and metastable strange dibaryons were searched a long time ago at LHE JINR, too. A thorough analysis of the merits and demerits of the available detection techniques and data analysis procedures has convinced us that the propane bubble chamber technique is most adequate to the physics problem at the present reconnaissance stage. A reliable identification of dibaryons needs a multivertex kinematic analysis which is in turn is feasible only using 4π -detectors and high precision measurements of the sought objects.

The investigations of multiquark states have been carried out in two directions.

1.1

Multiquark resonance formation via the compression mechanism reduces to the phase transition of normal density super strange hadronic matter revealing itself as multiquark resonances [1-3].

The problem of experimental examination of multibaryon strange resonances was started at LHE from 1962 up to now. The effective mass spectra of 17 strange multiquark systems were studied for neutron exposure n ${}^{12}C \rightarrow \Lambda X$ at average momentum $\langle p_n \rangle = 7.0 \text{ GeV/c}$, and our group succeeded in finding resonance-like peaks [7-10](Table 1) only in five of them $\Lambda p, \Lambda p\pi, \Lambda\Lambda, \Lambda\Lambda p, \Lambda\pi^+\pi^+$. Most significant evidence is included in the Review of Particle Properties.

Up to now, the same group is going on to collect statistics(more 3 time) for strange V^0 particles on the photographs of the JINR 2m propane bubble chamber exposed to a 10 GeV/c proton beam because it is the necessary to improve a statistical significance of the identified resonance peaks and to search for new strange multiquark resonance states.

below.						
Resonanse	М	Γ	Signifi-	σ	Bag model	predictions
system	MeV/c^2	MeV/c^2	cance	$(\mu b/^{12}C)$	М	J^p
			N_{sd}		(MeV/c^2)	
Λp	2095.0 ± 2.0	$7.0{\pm}2.0$	$5.70{\pm}1.20$	$55.0{\pm}16.0$	2110	1-
Λp	2181.0 ± 2.0	$3.2{\pm}0.5$	$4.36{\pm}1.21$	$60.0{\pm}15.0$	2169	1^{+}
Λp	2223.6 ± 1.8	$22.0{\pm}1.9$	$6.24{\pm}1.23$	$40.0{\pm}12.0$	2230	0^{+}
Λp	2263.0 ± 3.0	15.6 ± 2.3	8.55 ± 1.35	85.3 ± 20.0	2241	2^{+}
Λp	2356.0 ± 4.0	$98.6 {\pm} 2.5$	13.81 ± 1.39	$65.0{\pm}17.0$	2253	1^{+}
Λp	2129.2 ± 0.3	$0.7 {\pm} 0.16$	$11.37 {\pm} 1.37$	$90.0{\pm}20.0$	2128	1^{+}
$\Lambda p\pi^{\pm}$	2495.2 ± 8.7	204.5 ± 5.6	$12.86{\pm}1.68$	70.5 ± 15.0	2500	$0^{-}, 1^{-}, 2^{-}$
ΛΛ	2365.3 ± 9.6	47.2 ± 15.1	$4.2{\pm}1.40$	24.2 ± 7.0	2365	
$\Lambda\Lambda p$	≈ 3568.3	<60		16.1 ± 5.2	3570	$5/2^{+}$
$\Lambda \pi^+ \pi^+$	1704.9 ± 0.9	$18.0 {\pm} 0.5$	5.3 ± 1.6	$19.0 {\pm} 0.6$	1710	$1/2^{-}$
$\Lambda \pi^+ \pi^+$	2071.6 ± 4.0	172.9 ± 12.4	10.3 ± 1.5	$88.0{\pm}27.0$	2120	$1/2^{-}$
$\Lambda \pi^+ \pi^+$	2604.8 ± 4.8	85.9 ± 21.5	5.2 ± 1.4	$31.9 {\pm} 9.0$	2615	$3/2^{-}$

Table 1: The effective mass spectra in collisions of 4.0 GeV/c π^- and neutrons of a 7.0 GeV/c average momentum with ${}^{12}C$ nuclei, have led to the discovery of the peaks presented below.

Fig.1(a,b,c) show the preliminary experimental effective-mass spectrum of the $\Lambda \pi$, Λp and $\Lambda p\pi$ produced by pC interaction.

1.2

A simple consideration of symmetry and the properties of color-magnetic interactions argues in flavor of increasing binding in three flavor matter-systems containing u,d and s quarks. These objects known an H particles(uuddss). This was first proposed by Jaffe in 1977 using the MIT bag model [3-6]. The possibility that H dibaryon matter may exist in the core of a neutron star was also pointed out. The estimated binding energy of the H particle is model dependent and ranges from positive (unbound) to negative strong bound states(-650MeV). The heavy isotriplet stable dibaryon H (I=1, J=0+, Y=0, B=2, S=-2) of a 2370 MeV/c^2 mass is predicted by the soliton Skirme-like model [1,3].

The search for weak decay channels for stable S=-2,3 dibaryon states is being continued to date. A few events, detected on the photographs of the propane bubble chamber exposed to a 10 GeV/c proton beam, were interpreted as H dibaryons [11-16]. There are two groups of events interpreted as S=-2 stable dibaryons(Table 2):

a) The first group is formed of three neutral, S=-2 stable dibaryons, the masses of which are below $\Lambda\Lambda$ threshold;

b) The second group is formed of neutral and positively charged S=-2 heavy stable dibaryons (Figs.2,3). The masses of all the three dibaryons coincide within the errors are over the $\Lambda\Lambda$, ΞN , $\Lambda\Sigma$ threshold.

Ν	Channel of decay	Mass H	C.L.	References
		(MeV/c^2)	of fit	
		Dibaryon	%	
1	$H^0 \to \Sigma^- p$	2172 ± 15	99	Z.Phys.C 39, 151(1988).
2	$H^0 \to \Sigma^- p, \Sigma^- \to n\pi^-$	2146 ± 1	30	JINR RC,
	$H_1^0 \to H^0(2146)\gamma$	2203 ± 6	51	N $1(69)$ -95-61,1995.
3	$H^0 \rightarrow \Sigma^- p, \Sigma^- \rightarrow n\pi^-$	2218 ± 12	69	Phys.Lett B235(1990),208.
4	$H^0 \to \Sigma^- p, \Sigma^- \to n\pi^-$	2385 ± 31	34	Phys.Lett B316(1993),593.
5	$H^+ \to p \pi^0 \Lambda^0, \Lambda^0 \to p \pi^-$	2376 ± 10	87	Phys.Lett B316(1993),593
6	$H^+ \to p \pi^0 \Lambda^0, \Lambda^0 \to p \pi^-$	2580 ± 108	86	Nucl.Phys.75B(1999),63.
	$H^+n \to \Sigma^+\Lambda^0 \ n, \Lambda^0 \to p\pi^-$	2410 ± 90	6	
7	$H^+ \to p\gamma \Lambda^0, \Lambda^0 \to p\pi^-$	2448 ± 47	73	JINR Com.(2002)
	$H^+ \to p \pi^0 \Lambda^0, \Lambda^0 \to p \pi^-$	2488 ± 48	72	E1-2001-265

Table 2: Mass and weak decay channels for the registration of dibaryons.

References

- [1] Ts. Sakai et al., Osaka U., nucl-th/9912063, Dec. 1999.
- [2] P.J.G. Mulders, Aerts A.Th.M., de J.J. Swart, Phys.Rev., 1980, D.21, p.2553.
- [3] J. Schaffner-Bielich et al., Phys. Rev. C, v.55, n. 6, p. 3038-3046, 1997.
- [4] R.L. Jaffe, Phys. Rev. Lett. 38(1977)195.
- [5] T.Goldman et al., Phys.Rev.Lett., 1987, 59, p.627.
- [6] A. Faessler, U. Straub, Annalen der Physik, 1990, 47, p.439.
- [7] B.A. Shahbazian et al., Nucl. Physics, A374(1982), p. 73c-93.c.
- [8] B.A. Shahbazian ,JINR Commun., E1-82-446,1982, International Conference on Hypernuclear and Kaon Physics, Heidelberg, 1982.
- [9] B.A. Shakhbazian et al., (Dubna, JINR). JINR-D1-81-107, 1981.
- [10] B.A. Shahbazian et al., JINR Rapid Communications, No. 2[48]-91.
- [11] B.A. Shahbazian et al., JINR Rapid Communications, No. 1[69]-95.
- [12] B.A. Shahbazian et al., Phys. Lett. B316(1993)593.
- [13] P.Zh.Aslanian et al., JINR Rapid Communications, N 1(87)-98,1998.
- [14] P.Zh.Aslanyan et al., Nucl. Phys.B(Proc.Suppl.) 75B (1999)63-65.
- [15] P.Z. Aslanian et al., ISBN 981-02-4733-8, 2001, Int. Conference, Bologna'2000, 29 May - 3 June, Italy.
- [16] P.Zh.Aslanian et al., JINR Communications, E1-2001-265,2002.

