

Measurement of branching fractions and charge asymmetries in B^+ decays to $\eta\pi^+$, ηK^+ , $\eta\rho^+$ and $\eta'\pi^+$, and search for B^0 decays to ηK^0 and $\eta\omega$

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges-Pous,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ A. W. Borgland,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groyzman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. Cuhadar-Donszelmann,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ D. Thiessen,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² V. N. Ivanchenko,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ Sh. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ A. Lu,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzki,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ S. Yang,¹⁹ S. Jayatilleke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ I. M. Derrington,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² J. L. Harton,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² B. Spaan,²³ D. Altenburg,²⁴ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ E. Feltresi,²⁴ A. Hauke,²⁴ H. M. Lacker,²⁴ E. Maly,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ A. Sarti,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² M. J. Charles,³³ G. J. Grenier,³³ U. Mallik,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ T. C. Petersen,³⁵ M. Pierini,³⁵ S. Plaszczynski,³⁵ S. Rodier,³⁵ P. Roudeau,³⁵ M. H. Schune,³⁵ A. Stocchi,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ K. A. George,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹ R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ M. A. Winter,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ M. T. Naisbit,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³ V. B. Koptchev,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³

S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,⁵⁰ * G. De Nardo,⁵⁰ F. Fabozzi,⁵⁰ * C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malcès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ G. Simi,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,^{61,62} A. D’Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ F. F. Wilson,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Graziani,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn’ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ A. K. Mohapatra,⁶⁷ D. R. Muller,⁶⁷ C. P. O’Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ J. Strube,^{54,67} D. Su,⁶⁷ M. K. Sullivan,⁶⁷ J. M. Thompson,⁶⁷ J. Va’vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ H. Kim,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ M. Bomben,⁷⁴ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,⁷⁴ † L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ F. Martinez-Vidal,⁷⁵ R. S. Panvini,⁷⁶ † Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ B. Mellado,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ J. Wu,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ M. G. Greene,⁸⁰ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

- ¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA
¹⁵University of California at Riverside, Riverside, California 92521, USA
¹⁶University of California at San Diego, La Jolla, California 92093, USA
¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA
¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA
¹⁹California Institute of Technology, Pasadena, California 91125, USA
²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
²¹University of Colorado, Boulder, Colorado 80309, USA
²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Imperial College London, London, SW7 2AZ, United Kingdom
³³University of Iowa, Iowa City, Iowa 52242, USA
³⁴Iowa State University, Ames, Iowa 50011-3160, USA
³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁸Queen Mary, University of London, E1 4NS, United Kingdom
³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
⁴²University of Maryland, College Park, Maryland 20742, USA
⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁵McGill University, Montréal, Quebec, Canada H3A 2T8
⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁷University of Mississippi, University, Mississippi 38677, USA
⁴⁸Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7
⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵³Ohio State University, Columbus, Ohio 43210, USA
⁵⁴University of Oregon, Eugene, Oregon 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁸Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
⁶¹Princeton University, Princeton, New Jersey 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁸Stanford University, Stanford, California 94305-4060, USA
⁶⁹State University of New York, Albany, New York 12222, USA
⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
⁷¹University of Texas at Austin, Austin, Texas 78712, USA
⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁵IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6

⁷⁸*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*

⁷⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*

⁸⁰*Yale University, New Haven, Connecticut 06511, USA*

(Dated: November 1, 2018)

We present measurements of branching fractions and charge asymmetries for six B -meson decay modes with an η or η' meson in the final state. The data sample corresponds to 232 million $B\bar{B}$ pairs collected with the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- B Factory at SLAC. We measure the branching fractions (in units of 10^{-6}): $\mathcal{B}(B^+ \rightarrow \eta\pi^+) = 5.1 \pm 0.6 \pm 0.3$, $\mathcal{B}(B^+ \rightarrow \eta K^+) = 3.3 \pm 0.6 \pm 0.3$, $\mathcal{B}(B^0 \rightarrow \eta K^0) = 1.5 \pm 0.7 \pm 0.1$ (< 2.5 at 90% C.L.), $\mathcal{B}(B^+ \rightarrow \eta\rho^+) = 8.4 \pm 1.9 \pm 1.1$, $\mathcal{B}(B^0 \rightarrow \eta\omega) = 1.0 \pm 0.5 \pm 0.2$ (< 1.9 at 90% C.L.), and $\mathcal{B}(B^+ \rightarrow \eta'\pi^+) = 4.0 \pm 0.8 \pm 0.4$, where the first uncertainty is statistical and second systematic. We also determine the charge asymmetries for the charged modes: $\mathcal{A}_{ch}(B^+ \rightarrow \eta\pi^+) = -0.13 \pm 0.12 \pm 0.01$, $\mathcal{A}_{ch}(B^+ \rightarrow \eta K^+) = -0.20 \pm 0.15 \pm 0.01$, $\mathcal{A}_{ch}(B^+ \rightarrow \eta\rho^+) = 0.02 \pm 0.18 \pm 0.02$, and $\mathcal{A}_{ch}(B^+ \rightarrow \eta'\pi^+) = 0.14 \pm 0.16 \pm 0.01$.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

Charmless B decays are becoming increasingly useful to test the accuracy of theoretical predictions, for example based on QCD factorization [1, 2] or flavor SU(3) symmetry [3, 4]. In this Letter we present measurements of branching fractions and, when applicable, charge asymmetries for six charmless B decays: $B^+ \rightarrow \eta\pi^+$, $B^+ \rightarrow \eta K^+$, $B^0 \rightarrow \eta K^0$, $B^0 \rightarrow \eta\omega$, $B^+ \rightarrow \eta\rho^+$ and $B^+ \rightarrow \eta'\pi^+$, of which the last four were not observed before [5, 6, 7, 8]. Some of these decays may proceed through CKM-suppressed $b \rightarrow u$ and loop (“penguin”) $b \rightarrow s$ transitions with amplitudes of comparable size. Interference between these amplitudes can lead to direct CP violation measurable in charge asymmetries \mathcal{A}_{ch} [2]. The measured branching fractions and charge asymmetries may also be sensitive to the effect of non-Standard-model heavy particles entering the loop [9].

Charmless B decays with kaons are usually expected to be dominated by $b \rightarrow s$ loop amplitudes, while $b \rightarrow u$ tree amplitudes are typically larger for the decays with pions and ρ mesons. However, the $B \rightarrow \eta K$ decays are especially interesting since they are suppressed relative to the abundant $B \rightarrow \eta' K$ decays due to destructive interference between two penguin amplitudes [10]. The CKM-suppressed $b \rightarrow u$ tree amplitudes may interfere significantly with $b \rightarrow s$ penguin amplitudes of similar sizes, possibly leading to large direct CP violation in $B^+ \rightarrow \eta\rho^+$, $B^+ \rightarrow \eta\pi^+$ and $B^+ \rightarrow \eta'\pi^+$ [11]; numerical estimates are available in a few cases [2, 3, 12]. We search for such direct CP violation by measuring the charge asymmetry $\mathcal{A}_{ch} \equiv (\Gamma^- - \Gamma^+)/(\Gamma^- + \Gamma^+)$ in the rates $\Gamma^\pm = \Gamma(B^\pm \rightarrow f^\pm)$ for each charged final state f^\pm .

Finally, phenomenological fits to the branching fractions and charge asymmetries of charmless B decays can be used to understand the relative importance of tree and penguin contributions and may provide sensitivity to the CKM angle γ [3, 4].

The results presented here are obtained from extended unbinned maximum likelihood (ML) fits to data collected with the *BABAR* detector [13] at the PEP-II asymmetric e^+e^- collider [14] located at the Stanford Linear Accelerator Center.

The analysis uses an integrated luminosity of 211 fb^{-1} , corresponding to 232 million $B\bar{B}$ pairs, recorded at the $\Upsilon(4S)$ resonance (center-of-mass energy $\sqrt{s} = 10.58 \text{ GeV}$).

Charged particles are detected and their momenta measured by a combination of a vertex tracker consisting of five layers of double-sided silicon microstrip detectors and a 40-layer central drift chamber, both operating in the 1.5-T magnetic field of a superconducting solenoid. We identify photons and electrons using a CsI(Tl) electromagnetic calorimeter (EMC). Charged particle identification (PID) is provided by an internally reflecting ring imaging Cherenkov detector (DIRC) covering the central region of the detector, the average energy loss (dE/dx) in the tracking devices and by the EMC. A K/π separation better than two standard deviations (σ) is achieved for all momenta.

We select η , η' , ω , K_S^0 and π^0 candidates through the decays $\eta \rightarrow \gamma\gamma$ ($\eta_{\gamma\gamma}$), $\eta \rightarrow \pi^+\pi^-\pi^0$ ($\eta_{3\pi}$), $\eta' \rightarrow \eta\gamma\pi^+\pi^-$ ($\eta'_{\eta\pi\pi}$), $\eta' \rightarrow \rho^0\gamma$ ($\eta'_{\rho\gamma}$), $\omega \rightarrow \pi^+\pi^-\pi^0$, $K_S^0 \rightarrow \pi^+\pi^-$ and $\pi^0 \rightarrow \gamma\gamma$. We impose the following requirements on the invariant mass in MeV of the particle candidates' final states: $490 < m_{\gamma\gamma} < 600$ for $\eta_{\gamma\gamma}$, $520 < m_{\pi\pi\pi} < 570$ for $\eta_{3\pi}$, $910 < (m_{\eta\pi\pi}, m_{\rho\gamma}) < 1000$ for η' , $735 < m_{\pi\pi\pi} < 825$ for ω , $510 < m_{\pi\pi} < 1070$ for ρ^0 , $470 < m_{\pi\pi} < 1070$ for ρ^+ , $486 < m_{\pi\pi} < 510$ for K_S^0 and $120 < m_{\gamma\gamma} < 150$ for π^0 . These cuts are loose for the invariant mass variables used in the ML fit, and tight for those that are not. For K_S^0 candidates we require at least 3σ three-dimensional separation between the decay vertex and the e^+e^- collision point. For the vector resonances ω and ρ^+ we also use the helicity-frame decay angle θ_H . The helicity frame is defined as the vector-meson rest frame with polar axis along the direction of the boost from the B rest frame. For ω , θ_H is the polar angle of the normal to the decay plane, and for ρ it is the polar angle of the charged daughter momentum. We define $\mathcal{H} \equiv \cos\theta_H$ and require $-0.75 < \mathcal{H} < 0.95$ for ρ^+ .

All tracks from resonance candidates are required to have PID consistent with pions. For the B^+ decays to

$\eta\pi^+$, ηK^+ and $\eta'\pi^+$, the primary charged track must have an associated DIRC Cherenkov angle within 3.5σ of the expected value for either a π or K hypothesis. The discrimination between primary π and K is performed in the ML fits.

A B -meson candidate is characterized kinematically by the energy-substituted mass $m_{\text{ES}} = (\frac{1}{4}s - \mathbf{p}_B^2)^{\frac{1}{2}}$ and energy difference $\Delta E = E_B - \frac{1}{2}\sqrt{s}$, where (E_B, \mathbf{p}_B) is the B -meson 4-momentum vector, and all values are expressed in the $\Upsilon(4S)$ frame. Signal events peak at zero for ΔE , and at the B nominal mass for m_{ES} . The resolution on ΔE (m_{ES}) is about 30 MeV (3.0 MeV). We require $|\Delta E| \leq 0.2$ GeV and $5.25 \leq m_{\text{ES}} \leq 5.29$ GeV.

Backgrounds arise primarily from random combinations in continuum $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) events. To reject these events we make use of the angle θ_{T} between the thrust axis of the B candidate in the $\Upsilon(4S)$ frame and that of the rest of the charged tracks and neutral clusters in the event. The distribution of $|\cos\theta_{\text{T}}|$ is sharply peaked near 1 for combinations drawn from jet-like $q\bar{q}$ pairs, and nearly uniform for the almost isotropic B -meson decays; we require $|\cos\theta_{\text{T}}| < 0.9$ (< 0.65 for $\eta'_{\rho\gamma}\pi^+$). Further discrimination from continuum in the ML fit is obtained from a Fisher discriminant \mathcal{F} that is described in detail elsewhere [6].

Where necessary, we use additional event selection criteria to reduce $B\bar{B}$ backgrounds from several charmless final states. Specifically, we require that photons have energies in ranges uncharacteristic of these backgrounds and, in $\eta_{\gamma\gamma}\omega$ and $\eta_{\gamma\gamma}K_s^0$, we eliminate $\eta_{\gamma\gamma}$ candidates that share a photon with any π^0 candidate having momentum between 1.9 and 3.1 GeV in the $\Upsilon(4S)$ frame.

Multiple candidates are found in less than 30% of the events, in which case we choose the candidate with the smallest value of a χ^2 constructed from the deviations of the daughter resonance masses from their nominal values.

We use Monte Carlo (MC) simulation [15] for an initial estimate of the residual $B\bar{B}$ background and to identify the few (mostly charmless) decays that may survive the candidate selection and have characteristics similar to the signal. We find these contributions to be negligible for several of our modes. Where they are not negligible, namely for $\eta_{\gamma\gamma}\pi^+$, $\eta_{\gamma\gamma}K^+$, $\eta\rho^+$ and $\eta'_{\rho\gamma}\pi^+$, we include a component in the ML fit to account for them.

We obtain yields and \mathcal{A}_{ch} for each decay chain from a ML fit with the following input observables: ΔE , m_{ES} , \mathcal{F} , and m_{res} (the mass of the η , η' , ρ^+ , or ω candidate). For ω and ρ^+ decays we also use \mathcal{H} and, for charged modes with a primary charged track, the PID variables S_{π} and S_K , defined as the number of standard deviations between the measured DIRC Cherenkov angle and that expected for pions and kaons, respectively.

For each event i , hypothesis j (signal, continuum background, $B\bar{B}$ background), and flavor k (primary π^+ or

K^+), we define the probability density function (PDF)

$$\mathcal{P}_{jk}^i = \mathcal{P}_j(m_{\text{ES}}^i)\mathcal{P}_j(\Delta E_k^i [S_k^i])\mathcal{P}_j(\mathcal{F}^i)\mathcal{P}_j(m_{\text{res}}^i [H^i]) \quad (1)$$

The bracketed variables S and \mathcal{H} pertain to modes with a primary charged track or vector resonance daughters, respectively. Known correlations between ΔE_k and S_k , and between m_{res} and \mathcal{H} , are included in the PDF. The likelihood function is

$$\mathcal{L} = \exp\left(-\sum_{j,k} Y_{jk}\right) \prod_i^N \left[\sum_{j,k} Y_{jk} \mathcal{P}_{jk}^i \right], \quad (2)$$

where Y_{jk} is the yield of events of hypothesis j and flavor k , to be found by maximizing \mathcal{L} . N is the number of events in the sample. Free parameters of the fit are the signal and background yields, $q\bar{q}$ background PDF parameters (see below), and for charged modes the signal and $q\bar{q}$ background charge asymmetries.

For the signal and $B\bar{B}$ background components we determine the PDF parameters from simulation. For background from continuum (and non-peaking combinations from B decays) we obtain the PDF from $(m_{\text{ES}}, \Delta E)$ sideband data for each decay chain, before applying the fit to data in the signal region; we refine this PDF by letting as many of its parameters as feasible free to vary in the final fit. We parameterize each of the functions $\mathcal{P}_{\text{sig}}(m_{\text{ES}})$, $\mathcal{P}_{\text{sig}}(\Delta E_k)$, $\mathcal{P}_j(\mathcal{F})$, $\mathcal{P}(S_k)$ and the peaking components of $\mathcal{P}_j(m_{\text{res}})$ with either a Gaussian, the sum of two Gaussians or an asymmetric Gaussian function as required to describe the distribution. Slowly varying distributions (mass, energy or helicity-angle for combinatorial background) are represented by one or a combination of linear, quadratic and phase-space motivated functions [6]. The peaking and combinatorial components of the ω and ρ^+ mass spectra each have their own \mathcal{H} shapes. Control samples with similar topologies as our signal modes (e.g. $B \rightarrow D(K\pi\pi)\pi$) are used to verify or adjust the simulated resolutions evaluated from MC.

Before applying the fitting procedure to the data we subject it to several tests. In particular we evaluate possible biases in the yields from our neglect of small residual correlations among discriminating variables in the PDFs. This is achieved by fitting ensembles of simulated $q\bar{q}$ experiments drawn from the PDF into which we have embedded the expected number of signal and $B\bar{B}$ background events, randomly extracted from the fully simulated MC samples. The measured biases are listed in Table I.

The branching fraction for each decay is obtained from the measured yield, corrected for the fit bias and for the selection efficiency, and the number of $B\bar{B}$ pairs. We assume equal decay rates of the $\Upsilon(4S)$ to B^+B^- and $B^0\bar{B}^0$. In Table I we show for each decay mode the measured branching fraction together with the event yield and efficiency, and \mathcal{A}_{ch} when applicable. The purity is the ratio

TABLE I: Fitted signal yield Y_S in events (ev.), estimated purity P , measured bias (see text), detection efficiency ϵ , daughter branching fraction product ($\prod \mathcal{B}_i$), significance \mathcal{S} (with systematic uncertainties included), measured branching fraction \mathcal{B} , and signal charge asymmetry \mathcal{A}_{ch} for each mode. The quantities in parentheses are 90% C.L. upper limits.

| Mode | Y_S (ev.) | P (%) | Bias (ev.) | ϵ (%) | $\prod \mathcal{B}_i$ (%) | \mathcal{S} (σ) | \mathcal{B} (10^{-6}) | \mathcal{A}_{ch} |
|--------------------------------|-------------------|---------|------------|----------------|---------------------------|----------------------------|-----------------------------------------------------------------|---------------------------------------------|
| $\eta_{\gamma\gamma}\pi^+$ | 153_{-22}^{+24} | 30 | +7 | 33 | 39 | 7.9 | $4.8_{-0.7}^{+0.8}$ | -0.04 ± 0.14 |
| $\eta_{3\pi}\pi^+$ | 76_{-15}^{+16} | 32 | +6 | 24 | 23 | 5.6 | $5.6_{-1.2}^{+1.3}$ | -0.32 ± 0.20 |
| $\eta\pi^+$ | | | | | | 9.7 | $5.1 \pm 0.6 \pm 0.3$ | $-0.13 \pm 0.12 \pm 0.01$ |
| $\eta_{\gamma\gamma}K^+$ | 116_{-19}^{+21} | 29 | +8 | 32 | 39 | 6.1 | 3.6 ± 0.7 | -0.19 ± 0.16 |
| $\eta_{3\pi}K^+$ | 37_{-12}^{+13} | 24 | +5 | 23 | 23 | 2.8 | $2.6_{-1.0}^{+1.1}$ | -0.22 ± 0.33 |
| ηK^+ | | | | | | 6.7 | $3.3 \pm 0.6 \pm 0.3$ | $-0.20 \pm 0.15 \pm 0.01$ |
| $\eta_{\gamma\gamma}K^0$ | 17_{-7}^{+9} | 27 | +3 | 28 | 14 | 2.3 | $1.6_{-0.9}^{+1.0}$ | |
| $\eta_{3\pi}K^0$ | 5_{-3}^{+5} | 28 | +1 | 21 | 8 | 1.4 | $1.1_{-0.9}^{+1.3}$ | |
| ηK^0 | | | | | | 2.6 | $1.5 \pm 0.7 \pm 0.1$ (< 2.5) | |
| $\eta_{\gamma\gamma}\omega$ | 13_{-6}^{+7} | 32 | +1 | 14 | 35 | 2.5 | $1.1_{-0.5}^{+0.6}$ | |
| $\eta_{3\pi}\omega$ | 2_{-5}^{+7} | 6 | -1 | 11 | 20 | 0.6 | $0.6_{-1.0}^{+1.3}$ | |
| $\eta\omega$ | | | | | | 2.5 | $1.0 \pm 0.5 \pm 0.2$ (< 1.9) | |
| $\eta_{\gamma\gamma}\rho^+$ | 126_{-32}^{+34} | 12 | +18 | 16 | 39 | 3.7 | $7.3_{-2.2}^{+2.4}$ | 0.10 ± 0.23 |
| $\eta_{3\pi}\rho^+$ | 65_{-20}^{+22} | 15 | +3 | 11 | 23 | 3.4 | $10.6_{-3.5}^{+3.7}$ | -0.14 ± 0.31 |
| $\eta\rho^+$ | | | | | | 4.7 | $8.4 \pm 1.9 \pm 1.1$ | $0.02 \pm 0.18 \pm 0.02$ |
| $\eta'_{\eta\pi\pi}\pi^+$ | 69_{-12}^{+13} | 42 | +9 | 27 | 18 | 5.6 | $5.5_{-1.1}^{+1.2}$ | 0.09 ± 0.18 |
| $\eta'_{\rho\gamma}\pi^+$ | 30_{-15}^{+16} | 13 | +9 | 17 | 30 | 1.4 | $1.8_{-1.2}^{+1.3}$ | 0.58 ± 0.44 |
| $\eta'\pi^+$ | | | | | | 5.4 | $4.0 \pm 0.8 \pm 0.4$ | $0.14 \pm 0.16 \pm 0.01$ |

of the signal yield (Y_S) to the effective background plus signal ($Y_B^{\text{eff}} + Y_S$), which we estimate as the square of the uncertainty in the signal yield ($Y_B^{\text{eff}} + Y_S \equiv \sigma_{Y_S}^2$).

The statistical uncertainties in the signal yield and \mathcal{A}_{ch} are taken as the change in the central value when the quantity $-2 \ln \mathcal{L}$ increases by one unit from its minimum value. The significance is taken as the square root of the difference between the value of $-2 \ln \mathcal{L}$ (with systematic uncertainties included) for zero signal and the value at its minimum.

For each mode the measurements for separate daughter decays are combined by adding the values of $-2 \ln \mathcal{L}$ as functions of branching fraction, taking proper account of the correlated and uncorrelated systematic uncertainties described below [6]. For $\eta\omega$ and ηK^0 we quote 90% confidence level (C.L.) upper limits, taken to be the branching fraction below which lies 90% of the total of the likelihood integral in the positive branching fraction region.

In Fig. 1 we show projections onto m_{ES} and ΔE of subsamples enriched with a mode-dependent threshold requirement on the signal likelihood (computed without of the variable plotted) that optimizes the sensitivity.

Most of the uncertainties arising from lack of knowledge of the PDFs have been included in the statistical uncertainty since most background parameters are free in the fit. For the signal the uncertainties in PDF parameters are estimated from the consistency of fits to MC and data in control modes with similar final states. Varying the signal PDF parameters within these errors, we estimate the mode-dependent uncertainties due to the signal PDFs to be 1–8 events. We verify the validity of




FIG. 1: The B candidate m_{ES} (left) and ΔE (right) projections obtained with a cut on the signal likelihood (see text) for $B^+ \rightarrow \eta\rho^+$ (a, b), $B^+ \rightarrow \eta'\pi^+$ (c, d), and combined $B^+ \rightarrow \eta\pi^+$ and $B^+ \rightarrow \eta K^+$ (e, f). Points with uncertainties represent the data, solid curves the full fit functions, dashed curves the background functions and the dotted curves the background plus signal ηK^+ functions.

the fit procedure and PDF shapes by demonstrating that the likelihood for each fit is consistent with the distribution found in simulation.

The uncertainty in the fit bias correction is taken to be half of the correction itself. Similarly we estimate the uncertainty from modeling the $B\bar{B}$ backgrounds by taking half of the difference between the signal yield fitted with and without the $B\bar{B}$ background component.

Uncertainties in our knowledge of the reconstruction efficiency, found from auxiliary studies on inclusive control samples [6], include 0.6% per primary track, 0.8% per track from a resonance, 1.5% per photon, and 2.1% for a K_s^0 . Our estimate of the systematic uncertainty in the number of $B\bar{B}$ pairs is 1.1%. Published data [16] provide the uncertainties in the B -daughter product branching fractions (1–3%). The uncertainties in the efficiency of the event selection are 1% (4% in $B^+ \rightarrow \eta'_{\rho^+} \pi^+$) for the requirement on $\cos\theta_T$ and 1% for PID. Using several large inclusive kaon and B -decay samples, we find a systematic uncertainty for \mathcal{A}_{ch} of 1.1%, due mainly to the dependence of reconstruction efficiency on the charge, for the high momentum pion from $B^+ \rightarrow \eta\pi^+$, ηK^+ and $\eta'\pi^+$. The corresponding number for the softer charged pion from the ρ^+ in $B^+ \rightarrow \eta\rho^+$ is 2%.

In this Letter, we have presented improved measurements of branching fractions for six charmless B -meson decays. All branching fractions are in agreement with theoretical predictions. The previously unobserved $B^+ \rightarrow \eta\rho^+$ and $B^+ \rightarrow \eta'\pi^+$ decay modes are seen with significance 4.7σ and 5.4σ , respectively. For the charged modes, we also determine the charge asymmetries. These are found to be consistent with zero within their uncertainties.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support BABAR. We wish to acknowledge support from the University of Colorado Undergraduate Research Opportunities Program. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A. P. Sloan Foundation, Research Corporation, and

Alexander von Humboldt Foundation.

* Also with Università della Basilicata, Potenza, Italy
 † Deceased

- [1] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003) and references therein.
- [2] M.-Z. Yang and Y.-D. Yang, Nucl. Phys. B **609**, 469 (2001); M. Beneke and M. Neubert, Nucl. Phys. B **651**, 225 (2003);
- [3] H. K. Fu *et al.*, Phys. Rev. D **69**, 074002 (2004).
- [4] C.-W. Chiang, M. Gronau, and J. L. Rosner, Phys. Rev. D **68**, 074012 (2003); C.-W. Chiang *et al.*, Phys. Rev. D **69**, 034001 (2004); C.-W. Chiang *et al.*, Phys. Rev. D **70**, 034020 (2004).
- [5] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **92**, 061801 (2004).
- [6] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. D **70**, 032006 (2004).
- [7] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **93**, 181806 (2004).
- [8] Belle Collaboration, P. Chang *et al.*, hep-ex/0412043.
- [9] A. Datta and D. London, Phys. Lett. B **595**, 453 (2004); M. Ciuchini *et al.*, Phys. Rev. D **67**, 075016 (2003).
- [10] H. J. Lipkin, Phys. Lett. B **254**, 247 (1991).
- [11] M. Bander, D. Silverman, and A. Soni, Phys. Rev. Lett. **43**, 242 (1979); S. Barshay, D. Rein, and L.M. Sehgal, Phys. Lett. B **259**, 475 (1991); A.S. Dighe, M. Gronau, and J.L. Rosner, Phys. Rev. Lett. **79**, 4333 (1997).
- [12] G. Kramer, W.F. Palmer, and H. Simma, Nucl. Phys. B **428**, 77 (1994); A. Ali, G. Kramer, and C.-D. Lü, Phys. Rev. D **59**, 014005 (1999).
- [13] BABAR Collaboration, B. Aubert *et al.*, Nucl. Instr. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [14] PEP-II Conceptual Design Report, SLAC-R-418 (1993).
- [15] The BABAR detector Monte Carlo simulation is based on GEANT4: S. Agostinelli *et al.*, Nucl. Instr. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [16] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).