


EXCLUSIVE $\rho\rho$ PRODUCTION IN TAGGED $\gamma\gamma$ INTERACTIONS AT LEP

I. VOROBIEV ^a (on behalf of the L3 Collaboration)
Carnegie Mellon University, Pittsburgh, PA 15213



Exclusive $\rho\rho$ production in two-photon collisions between a quasi-real, γ , and a virtual, γ^* , photons is studied at LEP at centre-of-mass energies $89 \text{ GeV} \leq \sqrt{s} \leq 209 \text{ GeV}$ with a total integrated luminosity of 854.7 pb^{-1} . The cross sections of the $\gamma\gamma^* \rightarrow \rho\rho$ processes are determined as a function of the photon virtuality, Q^2 , and the two-photon centre-of-mass energy, $W_{\gamma\gamma}$, in the kinematic region: $0.2 \text{ GeV}^2 \leq Q^2 \leq 30 \text{ GeV}^2$ and $1.1 \text{ GeV} \leq W_{\gamma\gamma} \leq 3 \text{ GeV}$.

1 Introduction

Recently a QCD model was proposed ¹ for calculating the cross section of exclusive meson pair production in gamma-gamma interactions with one highly virtual photon having $Q^2 \gg W_{\gamma\gamma}$. In this model, the exclusive process is factorisable into a perturbative, calculable, short distance scattering $\gamma\gamma^* \rightarrow q\bar{q}$ or $\gamma\gamma^* \rightarrow gg$ and non-perturbative matrix elements, which are called generalized distribution amplitudes, describing the transition of the two partons into hadron pairs.

The L3 Collaboration performed a series of measurements of neutral and charged ρ -meson pair production in processes

$$e^+e^- \rightarrow e^\pm e^\mp_{tag} \gamma\gamma^* \rightarrow e^\pm e^\mp_{tag} \rho\rho,$$

where one electron/positron is tagged either by Luminosity Monitor (LUMI) or Very Small Angle Tagger (VSAT) ($\rho^0\rho^0$ with LUMI - Ref.², $\rho^+\rho^-$ with LUMI - Ref.³, $\rho^0\rho^0$ with VSAT - Ref.⁴, $\rho^+\rho^-$ with VSAT - Ref.⁵). LUMI covers Q^2 range $1.2 - 8.5 \text{ GeV}^2$ at LEPI, $8.8 - 30 \text{ GeV}^2$ at LEPII, and VSAT (available only at LEPII) - $0.2 - 0.85 \text{ GeV}^2$.

^a *Present address:* CERN, CH-1211 Geneva 23, Switzerland; e-mail: Igor.Vorobiev@cern.ch




Figure 1: Mass distributions for the selected $\pi^+\pi^-\pi^0\pi^0$ events with VSAT tag: a) the four-pion system mass, $W_{\gamma\gamma}$; b) the $\pi^\pm\pi^0$ combinations (four entries per event); c) correlation between the $\pi^-\pi^0$ and $\pi^+\pi^0$ pairs (two entries per event) and d) correlation between the $\pi^+\pi^-$ and $\pi^0\pi^0$ pairs.

2 Measurements

Measurements with LUMI were done at LEPI (\sqrt{s} around 91 GeV) with integrated luminosity $L = 148.7pb^{-1}$ and at LEPII ($161 \text{ GeV} \leq \sqrt{s} \leq 209 \text{ GeV}$) with $L = 706.0pb^{-1}$, measurements with VSAT - at LEPII ($183 \text{ GeV} \leq \sqrt{s} \leq 209 \text{ GeV}$) with $L = 684.8pb^{-1}$. Underlying selected reactions are

$$e^+e^- \rightarrow e^\pm e^\mp_{tag} \pi^+\pi^-\pi^+\pi^- \quad (1)$$

$$e^+e^- \rightarrow e^\pm e^\mp_{tag} \pi^+\pi^-\pi^0\pi^0 \quad (2)$$

934 events of reaction (1) were selected in LUMI sample and 1938 - in VSAT one. For reaction (2) - 343 and 414 events respectively. Example of distributions for selected events (for reaction (2) with VSAT) is shown in Figure 1. Mass of charged pion pairs (Figure 1b) shows a strong signal of ρ^\pm production, and two-dimensional distribution of such masses (Figure 1c) presents concentration of events in region of $\rho^+\rho^-$ signal. No structure is observed in the correlation plot of $\pi^+\pi^-$ and $\pi^0\pi^0$ mass combinations (Figure 1d). For further analysis and cross section determination events in the region $1.1 \text{ GeV} \leq W_{\gamma\gamma} \leq 3 \text{ GeV}$ were left (Figure 1a).

In order to determine the differential $\rho\rho$ production rates, a maximum likelihood fit was performed in intervals of Q^2 and $W_{\gamma\gamma}$ using a box method. Data were fit to a sum of non-interfering




Figure 2: The $\rho\rho$ production cross section as a function of Q^2 : a) differential cross section of the process $e^+e^- \rightarrow e^+e^-\rho\rho$ and b) cross section of the process $\gamma\gamma^* \rightarrow \rho\rho$. c)d)e) Cross section of the process $\gamma\gamma^* \rightarrow \rho\rho$ as a function of $W_{\gamma\gamma}$ in three Q^2 intervals.

contributions from the processes, generated by Monte Carlo in simple model of isotropic production and phase space decay. For data sample with LUMI the processes used in fit were: for the reaction (1): $\gamma\gamma^* \rightarrow \rho^0\rho^0$, $\gamma\gamma^* \rightarrow \rho^0\pi^+\pi^-$, $\gamma\gamma^* \rightarrow \pi^+\pi^-\pi^+\pi^-$ (non resonant); for the reaction (2): $\gamma\gamma^* \rightarrow \rho^+\rho^-$, $\gamma\gamma^* \rightarrow \rho^\pm\pi^0\pi^0$, $\gamma\gamma^* \rightarrow \pi^+\pi^-\pi^0\pi^0$ (non resonant). For data sample with VSAT it was found that a fit is improved if to include additional channels - for the reaction (1): $\gamma\gamma^* \rightarrow f_2(1270)\rho^0$, $\gamma\gamma^* \rightarrow f_2(1270)\pi^+\pi^-$ and for the reaction (2): $\gamma\gamma^* \rightarrow a^\pm_2(1320)\pi^0$ (we do not determine the cross sections of these processes). The inputs to the fit were the six possible two-pion mass combinations in an event. Fit provides a good description of all mass and angular distributions. From numbers of $\rho\rho$ events, obtained by fit in intervals of Q^2 and $W_{\gamma\gamma}$ and corrected by efficiencies and background, cross sections of $\rho\rho$ pair production were calculated.

3 Results

The cross section dependence on Q^2 and $W_{\gamma\gamma}$ for both $\rho^0\rho^0$ and $\rho^+\rho^-$ is shown on Figure 2. Q^2 dependence of differential cross section can be well described by the approximate formula

$d\sigma_{ee}/dQ^2 \sim 1/Q^n(Q^2 + \langle W_{\gamma\gamma} \rangle^2)^2$, following from QCD-based calculations⁶, where n is expected to be equal 2. In the range $Q^2 > 1.2 \text{ GeV}^2$ fit gives values $n = 2.4 \pm 0.3$ for $\rho^0\rho^0$ and $n = 2.5 \pm 0.4$ for $\rho^+\rho^-$, which are compatible with the expected value 2. $\rho^0\rho^0$ production at $Q^2 > 1.2 \text{ GeV}^2$ was analyzed in paper⁷. Q^2 dependence of the differential cross section is perfectly reproduced by formulas with three phenomenological parameters, one of which, $C_1 = 1.20 \pm 0.23 \text{ GeV}^2$, gives normalization of $\rho\rho$ generalized distribution amplitudes.

Ratio of cross sections $\sigma_{e^+e^- \rightarrow e^+e^-\rho^+\rho^-} / \sigma_{e^+e^- \rightarrow e^+e^-\rho^0\rho^0}$ (in the kinematic region $1.2 \text{ GeV}^2 \leq Q^2 \leq 8.5 \text{ GeV}^2$, $1.1 \text{ GeV} \leq W_{\gamma\gamma} \leq 2.1 \text{ GeV}$) is 1.8 ± 0.5 , which is compatible with the factor 2, expected for an isospin $I = 0$ state. Contrary to this measurement in high Q^2 region, in previous measurements done for $Q^2 \sim 0$ it was found that $\rho^0\rho^0$ cross section is few times higher than $\rho^+\rho^-$ one. And really, when we go to smaller Q^2 , the relative magnitude of $\rho^+\rho^-$ and $\rho^0\rho^0$ production changes in the vicinity of $Q^2 \approx 1 \text{ GeV}^2$, suggesting different ρ -pair production mechanisms at low and high Q^2 . In the whole Q^2 range fit to approximate formula gives $n = 2.3 \pm 0.15$ for $\rho^+\rho^-$ and $n = 2.9 \pm 0.14$ for $\rho^0\rho^0$ (fit is shown in Figure 2a by lines).

The measured cross section of the process $\gamma\gamma^* \rightarrow \rho\rho$ as a function of Q^2 is shown in Figure 2b. It is well described by GVDM model both for $\rho^+\rho^-$ and $\rho^0\rho^0$ at $Q^2 > 1 \text{ GeV}^2$, and for $\rho^0\rho^0$ - in the whole Q^2 range. The fit for $\rho^0\rho^0$ in the whole Q^2 range finds cross section $13.6 \pm 0.7 \text{ nb}$ at $Q^2 = 0$. The change of relative magnitude of $\rho^+\rho^-$ and $\rho^0\rho^0$ processes is clearly visible, when comparing low- and high- Q^2 regions. A ρ -pole description (is not shown on the picture) is excluded for both $\rho^0\rho^0$ and $\rho^+\rho^-$ data.

The cross sections of the processes $\gamma\gamma^* \rightarrow \rho\rho$ are plotted in Figure 2c-e as a function of $W_{\gamma\gamma}$. For $W_{\gamma\gamma} \leq 2.1 \text{ GeV}$ on Figure 2c there is a clear enhancement of $\rho^0\rho^0$ production relative to $\rho^+\rho^-$. Ratio $\sigma_{e^+e^- \rightarrow e^+e^-\rho^+\rho^-} / \sigma_{e^+e^- \rightarrow e^+e^-\rho^0\rho^0} = 0.6 \pm 0.1$ in this region.

4 Conclusions

This is the first measurement of tagged exclusive $\rho^0\rho^0$ production and of $\rho^+\rho^-$ production at high Q^2 . The measurements allow to follow the evolution of cross sections over a Q^2 -region of two orders of magnitude. A QCD-based form is found to provide a good parametrization in the entire Q^2 interval, where the differential cross sections show a monotonic decrease of more than four orders of magnitude (for the $W_{\gamma\gamma}$ range $1.1 \text{ GeV} \leq W_{\gamma\gamma} \leq 3 \text{ GeV}$). The Q^2 dependence of the process $\gamma\gamma^* \rightarrow \rho^0\rho^0$ is well reproduced by parametrization based on GVDM model over the entire Q^2 -region. But $\rho^+\rho^-$ data cannot be satisfactory described by such a parametrization in the whole Q^2 -region, only at $Q^2 > 1 \text{ GeV}^2$. A ρ -pole description is excluded for both $\rho^0\rho^0$ and $\rho^+\rho^-$ data. The relative magnitude of $\rho^+\rho^-$ and $\rho^0\rho^0$ production changes in the vicinity of $Q^2 \approx 1 \text{ GeV}^2$, suggesting different ρ -pair production mechanisms at low and high Q^2 . These data lay a new experimental grounds for obtaining information about QCD going from non-perturbative to the perturbative regime.

References

1. M. Diehl *et al*, *Phys. Rev. Lett.* **81**, 1782 (1998); N. Kivel, L. Mankiewicz and M.V. Polyakov, *Phys. Lett. B* **467**, 263 (1999); A. Freund, *Phys. Rev. D* **61**, 074010 (2000).
2. L3 Coll., P. Achard *et al.*, *Phys. Lett. B* **568**, 11 (2003).
3. L3 Coll., P. Achard *et al.*, *Phys. Lett. B* **597**, 26 (2004).
4. L3 Coll., P. Achard *et al.*, *Phys. Lett. B* **604**, 48 (2004).
5. L3 Coll., P. Achard *et al.*, Preprint CERN-PH-EP/2005-012, March 18, 2005.
6. M. Diehl, T. Gousset and B. Pire, *Phys. Rev. D* **62**, 073014 (2000).
7. I.V. Anikin, B. Pire and O.V. Teryaev, *Phys. Rev. D* **69**, 014018 (2004).