

Evidence for Parton k_T Effects in High- p_T Particle Production

L. Apanasevich,⁴ J. Bacigalupi,¹ W. Baker,³ M. Begel,⁹ S. Blusk,⁸ C. Bromberg,⁴
 P. Chang,⁵ B. Choudhary,² W. H. Chung,⁸ L. de Barbaro,⁹ W. DeSoi,⁹ W. Długosz,⁵
 J. Dunlea,⁹ E. Engels, Jr.,⁸ G. Fanourakis,⁹ T. Ferbel,⁹ J. Ftacnik,⁹ D. Garelick,⁵
 G. Ginther,⁹ M. Glaubman,⁵ P. Gutierrez,⁶ K. Hartman,⁷ J. Huston,⁴ C. Johnstone,³
 V. Kapoor,² J. Kuehler,⁶ C. Lirakis,⁵ F. Lobkowicz,⁹ P. Lukens,³ S. Mani,¹ J. Mansour,⁹
 A. Maul,⁴ R. Miller,⁴ B. Y. Oh,⁷ G. Osborne,⁹ D. Pellett,¹ E. Prebys,⁹ R. Roser,⁹
 P. Shepard,⁸ R. Shivpuri,² D. Skow,³ P. Slattery,⁹ L. Sorrell,⁴ D. Striley,⁵ W. Toothacker,⁷
 N. Varelas,⁹ D. Weerasundara,⁸ J. J. Whitmore,⁷ T. Yasuda,⁵ C. Yosef,⁴ M. Zieliński,⁹
 V. Zutshi²

(Fermilab E706 Collaboration)

¹*University of California-Davis, Davis, California 95616*

²*University of Delhi, Delhi, India 110007*

³*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

⁴*Michigan State University, East Lansing, Michigan 48824*

⁵*Northeastern University, Boston, Massachusetts 02115*

⁶*University of Oklahoma, Norman, Oklahoma 73019*

⁷*Pennsylvania State University, University Park, Pennsylvania 16802*

⁸*University of Pittsburgh, Pittsburgh, Pennsylvania 15260*

⁹*University of Rochester, Rochester, New York 14627*

(October 1, 2018)

Abstract

Inclusive π^0 and direct-photon cross sections in the kinematic range $3.5 < p_T < 12$ GeV/ c with central rapidities (y_{cm}) are presented for 530 and 800 GeV/ c proton beams and a 515 GeV/ c π^- beam incident on Be targets. Current Next-to-Leading-Order perturbative QCD calculations fail to adequately describe the data for conventional choices of scales. Kinematic distributions from these hard scattering events provide evidence that the interacting partons carry significant initial-state parton transverse momentum (k_T). Incorporating these k_T effects phenomenologically greatly improves the agreement between calculations and the measured cross sections.

PACS numbers: 12.38.Qk, 13.85.Qk, 13.85.Ni, 14.20.Dh

Typeset using REVTeX

In recent years, perturbative QCD (PQCD) has been tested in a wide variety of processes involving strong interactions at short distances, and increasing attention is now being directed towards areas that may be sensitive to shortcomings in the current theoretical description [1]. The high statistics samples of hard scattering data accumulated by Fermilab fixed-target experiment E706 provide an opportunity to probe such issues. This paper presents comparisons of PQCD calculations to our data on the production of direct photons and π^0 's with large transverse momenta (p_T). Direct-photon data have long been expected to provide an accurate determination of the distributions of gluons in hadrons, especially at large longitudinal momentum fraction (x), where information has proven difficult to obtain from other measurements. Inclusive meson production at large p_T probes a different mix of hard scattering processes and provides insights into parton fragmentation. For conventional choices of parameters, our data are not described satisfactorily by Next-to-Leading-Order (NLO) PQCD calculations [2]. Resolving the observed discrepancies is important for improving the understanding of both parton distribution functions (PDF) and parton fragmentation functions (FF).

Several interesting aspects of QCD contributions beyond Leading-Order (LO) can be investigated experimentally through studies of processes sensitive to transverse motion of the partons prior to the hard scatter. This k_T is presumably due to effects of hadron size (primordial k_T) as well as initial-state gluon radiation. Measurements of Drell-Yan pair production [3] and direct di-photon production [4] have demonstrated the presence of substantial effective k_T , (significantly larger than can be attributed to primordial k_T), and have revealed a significant \sqrt{s} dependence of $\langle k_T \rangle$. A resummation of soft gluon emissions has recently been used to reproduce the size of the effect observed in the WA70 direct di-photon data [5]. Other data also suggest $\langle k_T \rangle$ values larger than those expected from NLO PQCD calculations. Recent comparisons of p_T spectra from charm-particle hadroproduction to NLO PQCD results provide evidence that supplemental k_T may be required to properly describe the data [6]. Likewise, it has been suggested that the observed pattern of discrepancies between data from various direct-photon experiments and results from NLO PQCD calculations could be related to k_T effects [7].

E706 is designed to measure the production of direct photons, neutral mesons, and associated particles at high p_T . The apparatus features a large lead and liquid argon electromagnetic calorimeter and a charged particle spectrometer [8]. The experiment accumulated ≈ 10 events/pb of π^- beam data at 515 GeV/ c , ≈ 9 events/pb of proton beam data at 530 GeV/ c , and ≈ 11 events/pb of proton beam data at 800 GeV/ c on Be, Cu, and H targets (primarily Be). A variety of event selection triggers sensitive to high- p_T electromagnetic showers were employed (using different prescale factors) to accumulate data over a broad range of p_T .

The steep p_T dependences of neutral meson and direct-photon production make the measured cross sections very sensitive to uncertainties in the energy calibration. Therefore, achieving a precise and accurate calibration of the response of the electromagnetic calorimeter was essential to the success of E706. As a result of detailed studies of the data acquired, the uncertainty in the calibration of the energy response of the calorimeter was reduced to less than 0.5% [9].

The single-photon sample is composed of those photons not identified as elements of reconstructed two photon decays of π^0 or η meson candidates. The direct-photon signal

is extracted from the single-photon sample via statistical subtraction of the background contributions. These backgrounds are primarily due to photons from unreconstructed decays of neutral mesons. Failure to correctly identify a photon as originating from a π^0 or η decay occurs when the other photon from that decay converts in the target region, escapes the fiducial volume of the calorimeter, or is otherwise not reconstructed. Sources of direct-photon background have been modeled using the HERWIG event generator [10] and a detailed GEANT simulation [11] of the spectrometer response. These Monte Carlo generated events have been weighted to accurately represent our measured neutral meson production spectra.

The p_T dependences of inclusive π^0 and direct-photon cross sections are shown in Figs. 1, 2, and 3. The results of NLO PQCD calculations (using BKK FF for the π^0 [12]) are compared with the data [13]. For simplicity, all QCD scales (renormalization, factorization, and, where appropriate, fragmentation) have been set equal. The broken curves in Fig. 1 represent the results of NLO PQCD calculations using conventional choices of scales (and GRV PDF [14]). The calculations are quite sensitive to the scales (an indication of the importance of still higher order contributions), but even for rather small scales, the NLO calculations fail to describe our data. Using other recent PDF [15,16] in the calculations also does not adequately account for the discrepancy between the NLO PQCD results and our data (broken curves in Fig. 2) [17]. Differences between LO [18] and NLO calculations are likewise not large when compared to the difference observed between either and the measured cross sections (Fig. 3). However, PQCD at NLO may not adequately account for soft gluon radiation that imparts an effective transverse momentum to the incident partons.

Kinematic distributions for high-mass pairs of particles directly probe the transverse momentum of incident partons in hard scattering events. The p_T distribution of π^0 pairs produced in 515 GeV/c π^- -nucleon collisions is shown in the inset of Fig. 4. The results of LO PQCD calculations in which the incident partons have Gaussian transverse momentum distributions with $\langle k_T \rangle = 1.3$ GeV/c (dashed curve) and $\langle k_T \rangle = 1.7$ GeV/c (solid curve) are shown in Fig. 4 [18]; the dotted curve does not include any incident parton k_T . For these high-mass π^0 pairs, Fig. 4 also shows the distribution of the angle between the two π^0 's in the transverse plane ($\Delta\phi$), and provides a comparison to LO PQCD, both with and without supplemental k_T . Fragmentation itself generates significant pair- p_T and contributes to the width of the $\Delta\phi$ distributions, but supplemental $\langle k_T \rangle > 1$ GeV/c provides a much better description of the data.

Another kinematic variable that is sensitive to k_T is the out-of-plane momentum, p_{out} (the component of the momentum of one high- p_T particle, perpendicular to the plane defined by the incident beam direction and the direction of the other high- p_T particle). The p_{out} distributions for $\gamma\pi^0$ pairs produced in proton-nucleon interactions are shown in the upper part of Fig. 5, compared to LO results without k_T , and with $\langle k_T \rangle$ values chosen to bracket the data. The p_{out} distributions for π^0 pairs are also shown in Fig. 5 (lower plots). These distributions (pair- p_T , $\Delta\phi$, and p_{out}) show clear evidence for the presence of significant k_T (> 1 GeV/c) in the hard scattering interactions. The corresponding distributions for our other data samples also support this conclusion [19].

Our preliminary analyses of the kinematic distributions of pairs of direct photons, as well as studies of the distribution of the fractional momentum carried by individual charged particles in jets recoiling against isolated photons, also show evidence of substantial k_T , as do our comparisons of the measured high- p_T charged- D cross section to NLO PQCD results [8].

All these results suggest a supplemental $\langle k_T \rangle$ of order 1 GeV/ c .

Since the inclusive spectra fall rapidly with increasing p_T , the introduction of k_T -smearing has a significant effect on predicted cross sections. To approximate the effect of supplemental k_T smearing on the inclusive NLO PQCD calculations for direct-photon (and π^0) production, we calculated k_T factors (as functions of p_T) for different values of $\langle k_T \rangle$, by computing ratios of results from LO PQCD calculations [18] for different $\langle k_T \rangle$ values compared to results without k_T [20]. These same k_T factors were then applied to the results of NLO PQCD calculations. As indicated by the solid curves in Figs. 1, 2, and 3, reasonable representations of both the direct-photon and π^0 results are obtained using $\langle k_T \rangle$ values > 1 GeV/ c [21]. The kinematic distributions exhibit a pattern consistent with increasing $\langle k_T \rangle$ as \sqrt{s} increases, a trend reflected in the choices of $\langle k_T \rangle$ factors employed in the theory curves (solid curves) shown in the inclusive cross section plots.

As an illustration of the sensitivity of our data to the gluon distribution, Fig. 6 compares our direct-photon cross sections to NLO PQCD calculations using CTEQ4M and CTEQ4HJ PDF [16]. Once soft-gluon effects are satisfactorily taken into account, either approximately as in this paper or in a more theoretically rigorous manner, our data can be used to help discriminate between PDF that otherwise provide acceptable descriptions of the data sets used in Ref. [16].

In conclusion, we have measured the inclusive production of high- p_T neutral mesons and direct photons by 530 GeV/ c and 800 GeV/ c proton and 515 GeV/ c π^- beams. Current NLO PQCD calculations (which exhibit substantial dependences on QCD scales) fail to account for the measured cross sections using conventional choices of scales. Significant k_T effects (> 1 GeV/ c) have been observed in the kinematic distributions of high-mass pairs of π^0 's, as well as high-mass $\gamma\pi^0$ pairs. A simple implementation of supplemental parton k_T in PQCD calculations, using k_T values consistent with observations, provides a reasonable description of the inclusive cross sections. Our high statistics direct-photon data samples are directly sensitive to the gluon distribution at large x values. An improved theoretical understanding of soft-gluon effects in inclusive direct-photon production will facilitate the global determination of the gluon distribution function.

We are pleased to acknowledge the many contributions of the Fermilab staff. This research has been supported in part by the U. S. Department of Energy, the National Science Foundation, and the Universities Grants Commission of India.

REFERENCES

- [1] P. Burrows *et al.*, hep-ex/9612003; R. Brock, in *ICHEP '96 - Proceedings of the 28th International Conference on High Energy Physics*, edited by Z. Ajduk and A. K. Wróblewski, (World Scientific, Singapore, 1997), Vol. I, pp. 69-88; G. Sterman, *ibid.* Vol. I, pp. 103-120.
- [2] P. Aurenche *et al.*, Phys. Lett. **140B**, 87 (1984); E. L. Berger and J. Qiu, Phys. Rev. D **44**, 2002 (1991); F. Aversa *et al.*, Nucl. Phys. **B327**, 105 (1989).
- [3] B. Cox and P. K. Malhotra, Phys. Rev. D **29**, 63 (1984); G. Moreno *et al.*, Phys. Rev. D **43**, 2815 (1991).
- [4] WA70 Collaboration, E. Bonvin *et al.*, Phys. Lett. B **236**, 523 (1990); CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **70**, 2232 (1993).
- [5] P. Chiappetta, R. Fergani, and J.Ph. Guillet, Phys. Lett. B **348**, 646 (1995).
- [6] S. Frixione *et al.*, Nucl. Phys. **B431**, 453 (1994).
- [7] J. Huston *et al.*, Phys. Rev. D **51**, 6139 (1995).
- [8] L. Apanasevich *et al.*, Phys. Rev. D **56**, 1391 (1997).
- [9] L. Apanasevich *et al.*, FERMILAB-Pub-97/142-E (1997).
- [10] G. Marchesini *et al.*, Comput. Phys. Commun. **67**, 465 (1992).
- [11] R. Brun *et al.*, GEANT3 User's Guide CERN DD/EE/84-1, (1987).
- [12] J. Binnewies, B. A. Kniehl, and G. Kramer, Phys. Rev. D **52**, 4947 (1995).
- [13] Each theory calculation presented in this paper has been adjusted to account for nuclear effects. Our conclusions are fully supported by our hydrogen target data samples. For the inclusive high- p_T π^0 and direct-photon calculations, theory results were adjusted by $A^{\alpha-1}$, assuming $\alpha = 1.08$ and $\alpha = 1.04$, respectively.
- [14] M. Glück, E. Reya, and A. Vogt, Z. Phys. C **53**, 127 (1992); **53**, 651 (1992).
- [15] A. D. Martin, W. J. Stirling, and R. G. Roberts, Phys. Lett. B **354**, 155 (1995); M. Glück, E. Reya, and A. Vogt, Z. Phys. C **67**, 433 (1995).
- [16] H. L. Lai *et al.*, Phys. Rev. D **55**, 1280 (1997).
- [17] The variation in these results does not reflect the full uncertainty in the knowledge of the gluon distribution at large x values since little data have been available in this domain, and there is substantial overlap in the data used and the techniques employed by different groups in evaluating the various PDF.
- [18] J. F. Owens, Rev. Mod. Phys. **59**, 465 (1987).
- [19] M. Begel, Ph.D. thesis, University of Rochester (in preparation).
- [20] A possible variation of $\langle k_T \rangle$ with p_T , expected in PQCD, has been neglected in this analysis.
- [21] The $\langle k_T \rangle$ values that provide a corresponding level of agreement between LO direct-photon calculations (not shown) and our inclusive cross sections are somewhat larger ($\approx 10\%$) than the corresponding values for the NLO comparisons.




FIG. 3. The π^0 and direct-photon cross sections as functions of p_T for 800 GeV/c proton-nucleon interactions compared to LO and NLO PQCD results. The solid curves show NLO results adjusted for supplemental $\langle k_T \rangle$. (Note that the units for the π^0 and γ results differ by a factor of 1000.)




FIG. 4. The $\Delta\phi$ distribution for high-mass π^0 pairs produced in 515 GeV/ c π^- -nucleon interactions compared to curves showing LO PQCD results using various $\langle k_T \rangle$ values. The inset shows the pair- p_T distribution for such pairs and the corresponding results of LO PQCD calculations.

p Be at 530 GeV/c

p Be at 800 GeV/c




FIG. 5. The out-of-plane momentum distributions for high-mass pairs produced in proton-nucleon interactions at 530 and 800 GeV/c compared to results of LO PQCD calculations (using CTEQ4L PDF) for several $\langle k_T \rangle$ values.




FIG. 6. Direct-photon inclusive cross sections as functions of p_T for 530 and 800 GeV/ c proton-nucleon interactions compared to results of NLO PQCD calculations using CTEQ4HJ (dot-dashed curve) and CTEQ4M (solid curve) PDF. Factors for supplemental $\langle k_T \rangle$ are included. (Note that the units for the 530 and 800 GeV/ c results differ by a factor of 100.)