

$I = 2$ Pion Scattering Length with the Wilson Fermion

S. Aoki,¹ M. Fukugita,² S. Hashimoto,³ K-I. Ishikawa,^{1,4} N. Ishizuka,^{1,4} Y. Iwasaki,⁴
 K. Kanaya,^{1,4} T. Kaneko,³ Y. Kuramashi,³ M. Okawa,⁵ T. Onogi,⁶ S. Tominaga,⁴
 N. Tsutsui,³ A. Ukawa,^{1,4} N. Yamada,³ T. Yoshié,^{1,4}

(JLQCD Collaboration)

¹ *Institute of Physics, University of Tsukuba , Tsukuba, Ibaraki 305-8571, Japan*

² *Institute for Cosmic Ray Research, University of Tokyo , Kashiwa, Chiba 277-8582, Japan*

³ *High Energy Accelerator Research Organization (KEK) , Tsukuba, Ibaraki 305-0801, Japan*

⁴ *Center for Computational Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan*

⁵ *Department of Physics, Hiroshima University, Higashi-Hiroshima , Hiroshima 739-8526, Japan*

⁶ *Yukawa Institute for Theoretical Physics, Kyoto University , Kyoto 606-8502, Japan*

(February 8, 2020)

Abstract

The calculation of the $I = 2$ pion scattering length in quenched lattice QCD is revisited. The calculation is carried out with the Wilson fermion action employing Lüscher's finite size scaling method at $\beta = 5.9, 6.1$, and 6.3 corresponding to the range of lattice spacing $a \simeq 0.12 - 0.07$ fm. We obtain in the continuum limit $a_0 m_\pi = -0.0410(69)$, which is consistent with the prediction of chiral perturbation theory $a_0 m_\pi = -0.0444(10)$.

PACS number(s): 12.38.Gc, 11.15.Ha

Typeset using REVTeX

Lattice calculations of S -wave scattering lengths of the two-pion system are an important step to understand dynamical effects of strong interactions. There are already a number of calculations for the $I = 2$ process with either Staggered [1,2] or Wilson fermion action [1,3]. While these calculations gave results that are in gross agreement with the prediction of chiral perturbation theory (CHPT) [4], they were made on coarse and small lattices. More importantly, the continuum extrapolation was not made. Aiming to improve on these points, we carried out a calculation of the $I = 2$ S -wave scattering length in quenched lattice QCD. A preliminary result was reported in Ref. [5], in which some disagreement with the CHPT prediction was mentioned. In the mean time Liu *et.al.* carried out a similar calculation with the improved gauge and the Wilson fermion actions on anisotropic lattices [6].

We employ the standard plaquette action for gluons and the Wilson action for quarks, and explore the parameter range $m_\pi/m_\rho \sim 0.55 - 0.80$ for the chiral extrapolation and $a \sim 0.12 - 0.07$ fm for the continuum extrapolation. This is compared with the parameters of Liu *et.al.* which range $m_\pi/m_\rho \sim 0.7 - 0.9$ and $a_s \sim 0.4 - 0.2$ fm. Our calculations are made for parameters significantly closer to the chiral limit. In this short report we give the final result of our analysis.

The numbers of configurations (lattice sizes) are 187 ($16^3 \times 64$), 120 ($24^3 \times 64$), and 100 ($32^3 \times 80$) for $\beta = 5.9, 6.1$, and 6.3 , respectively. Quark propagators are solved with the Dirichlet boundary condition in the time direction and the periodic boundary condition in the space directions. The pion mass covers the range of $450 - 900$ MeV. The lattice constant is estimated from the ρ meson mass, which was obtained in our previous study [8], to be $a^{-1} = 1.64(2), 2.29(4), 3.02(5)$ (GeV) at $\beta = 5.9, 6.1$, and 6.3 . Our calculations were carried out on the Fujitsu VPP500/80 supercomputer at KEK.

The energy eigenvalue of a two-pion system in a finite periodic box L^3 is shifted by the finite-size effect. Lüscher presented a relation between the energy shift ΔE and the S -wave scattering length a_0 , given by [7]

$$-\Delta E \cdot \frac{m_\pi L^2}{4\pi^2} = T + C_1 \cdot T^2 + C_2 \cdot T^3 + O(T^4) , \quad (1)$$

where $T = a_0/(\pi L)$. The constants are $C_1 = -8.9136$ and $C_2 = 62.9205$ computed from geometry of the lattice. Since T has a small value, typically $\sim -10^{-2}$ in our simulation, we can safely neglect the higher order terms $O(T^4)$.

The energy shift ΔE can be obtained from the ratio

$$R(t) = G(t)/D(t) , \quad (2)$$

where

$$G(t) = \langle \pi^+(t) \pi^+(t) W^-(t_1) W^-(t_2) \rangle$$

$$D(t) = \langle \pi^+(t) W^-(t_1) \rangle \langle \pi^+(t) W^-(t_2) \rangle . \quad (3)$$

In order to enhance signals against the noise we use wall sources for π^- , which are denoted by W^- in (3), by fixing gauge configurations to the Coulomb gauge. The two wall sources are placed at different time slices t_1 and t_2 to avoid contaminations from Fierz-rearranged terms in the two-pion state which would occur for $t_1 = t_2$. We set $t_2 = t_1 + 1$ and $t_1 = 8, 10, 13$ for $\beta = 5.9, 6.1, 6.3$.

An example of $R(t)$ is plotted in Fig. 1 for $\beta = 6.3$ and $\kappa = 0.1513$ corresponding to $m_\pi = 433(4)\text{MeV}$. We see a clear, almost linear fall-off as a function of t till $t = 80$ even for a small energy $\Delta E \approx 20 \text{ MeV}$, showing that our wall sources work well for the two-pion state.

The energy shift ΔE is obtained from the linear term in the expansion of $R(t)$:

$$R(t) = Z \cdot \left(1 - \Delta E \cdot \tau + \mathcal{O}(\tau^2) \right) , \quad (4)$$

where $\tau = t - t_2$. The quadratic and higher order terms have no simple relations to ΔE due to effects from intermediate off-shell two-pion states [2] and quenching effects [9]. We first attempt to fit the data with the form

$$(\mathbf{Sqr}) \quad Z \cdot \left(1 - \Delta E \cdot \tau + E' \cdot \tau^2 \right) . \quad (5)$$

We find that this fit (**Sqr**) is quite ill-determined, since the two terms correlate so strongly, resulting in unacceptably large errors in ΔE and E' . We then attempt to fit with

$$\begin{aligned} (\mathbf{Exp}) \quad & Z \cdot \exp(-\Delta E \cdot \tau) , \\ (\mathbf{Lin}) \quad & Z \cdot \left(1 - \Delta E \cdot \tau \right) . \end{aligned} \quad (6)$$

These fitting forms give well-determined ΔE , while it may be contaminated by contributions from the second order term. We also include a fit of the form

$$(\mathbf{Old}) \quad Z - \Delta E \cdot \tau \quad (7)$$

into our attempts for completeness, since this was used in our preliminary report [5]. Note, however, that this form is theoretically correct only when Z is close to unity. The results for ΔE (and E' in case (**Sqr**)) are given in Table I for $\beta = 5.9$, Table II for $\beta = 6.1$, and Table III for $\beta = 6.3$. We take the same fitting range for the four fits, $t = 21 - 42$ for $\beta = 5.9$, $t = 25 - 50$ for $\beta = 6.1$, and $t = 27 - 62$ for $\beta = 6.3$. The value of χ^2 for each fitting is always small, and does not discriminate among fits. We do not consider case (**Sqr**) further because of very large errors, although the resulting central values for the energy shift are consistent with those from (**Exp**) and (**Lin**). The problem we must consider is whether we can remove contaminations of the second order term for ΔE from (**Exp**) and (**Lin**).

Figures 2, 3, and 4 show a_0/m_π as a function of the pion mass obtained at each β , with their numerical values tabulated in Table I, II, and III. We observe a large difference between **(Exp)** and **(Lin)**, indicating that contributions from the $O(\tau^2)$ term are indeed non-negligible and largely affect the determination of ΔE . The common in all figures of a_0/m_π versus m_π is that the data show a behavior linear in m_π^2 . We then fit

$$a_0/m_\pi = A + B \cdot m_\pi^2 \quad (8)$$

to extract the value A in the chiral limit. From the view point of CHPT we may in principle have a term $m_\pi^2 \log(m_\pi^2/\Lambda^2)$ added to (8). If we include this term with a free coefficient into the fit, however, the coefficients correlate too strongly that the fit is invalidated, producing a large error also for A . It is difficult to distinguish m_π^2 and $m_\pi^2 \log(m_\pi^2)$ within the range of m_π^2 that concerns us and the limited statistics. Since we do not see any significant curvature in the figure of a_0/m_π versus m_π , we simply drop this logarithmic term which itself vanishes at the chiral limit. We also note that for the Wilson fermion action the term $\propto 1/m_\pi^2$ may also exist, arising from explicit breaking of chiral symmetry, and also from quenching effects [9]. We do not see a $1/m_\pi^2$ effect, as our simulation is perhaps well away from $m_\pi^2 = 0$ and such a term is already damped into noise for the range of our simulation. Hence we do not include this term into our fit. In order to detect these two additional terms a simulation is needed close to the chiral limit with much higher statistics.

In Fig. 5 we present a_0/m_π in the chiral limit as a function of the lattice spacing, together with continuum extrapolations. This figure demonstrates a sizable scaling violation, but exhibits a very clean linear dependence as a function of a . It is interesting to observe that the difference between **(Exp)** and **(Lin)**, which are quite sizable on finite lattices, vanishes approaching the continuum limit. This shows that the second order term $O(\tau^2)$ included in (4) becomes irrelevant as $\Delta E \tau$ becomes sufficiently small; one may use any formula correct to the first order in τ to extract the ΔE . On the other hand, the extrapolation with **(Old)** gives a value somewhat different from the other two in the continuum limit, indicating that the departure of Z from unity could be non-negligible (although at $1.2 - 1.5\sigma$).

As our final value for the scattering length in the continuum limit at physical pion mass we take the result from **(Exp)**, which agrees with that from **(Lin)** but has a larger statistical error:

$$a_0/m_\pi = -2.09(35) \text{ 1/GeV}^2, \quad a_0 m_\pi = -0.0410(69), \quad (9)$$

where a rather large error arises from the continuum extrapolation. This result is compared with the CHPT prediction :

$$a_0/m_\pi = -2.265(51) \text{ (1/GeV}^2), \quad a_0 m_\pi = -0.0444(10) \quad (10)$$

The scattering length we derived at the continuum limit agrees well with the prediction of CHPT. The difference seen in the fitting formula of **(Old)** and **(Lin)** accounts for the 1.5σ difference of the lattice result from the CHPT prediction mentioned in our preliminary report, which is based on an incorrect extrapolation formula **(Old)**.

We remark that our results also agree with those of Liu *et.al.* [6], which use a long chiral extrapolation:

$$a_0/m_\pi = -1.75(38) \text{ (1/GeV}^2\text{)} , \quad a_0 m_\pi = -0.0342(75) \quad \text{for Scheme I} \quad (11)$$

$$a_0/m_\pi = -2.34(46) \text{ (1/GeV}^2\text{)} , \quad a_0 m_\pi = -0.0459(91) \quad \text{for Scheme II} \quad (12)$$

where two values (Scheme I and II) refer to their two different treatments for the finite volume corrections.

In this article we have reported a calculation of the scattering length for the $I = 2$ S -wave two pion system. We have shown that the result in the continuum limit is virtually independent of the choice of fitting functions used to extract ΔE from the ratio $R(t)$, and that it is consistent with the prediction of CHPT within our 15% statistical error.

REFERENCES

- [1] Y. Kuramashi, M. Fukugita, H. Mino, M. Okawa, and A. Ukawa, Phys. Rev. Lett. **71** (1993) 2387 ; M. Fukugita, Y. Kuramashi, M. Okawa, H. Mino, and A. Ukawa, Phys. Rev. **D52** (1995) 3003.
- [2] S.R. Sharpe, R. Gupta, and G.W. Kilcup, Nucl. Phys. **B383** (1992) 309.
- [3] R. Gupta, A. Patel, S.R. Sharpe, Phys. Rev. **D48** (1993) 388.
- [4] J. Gasser and H. Leutwyler, Phys. Lett. **B125** (1983) 325 ; J. Bijnens, G. Colangelo, G. Ecker, J. Gasser, and M.E. Sainio, Phys. Lett. **B374** (1996) 210; Nucl. Phys. **B508** (1997) 263. G. Colangelo, J. Gasser, and H. Leutwyler Nucl. Phys. **B603** (2001) 125.
- [5] JLQCD Collaboration, A. Aoki *et.al*, Nucl. Phys. (Proc. Suppl.) **83** (2000) 241.
- [6] C. Liu, J. Zhang, Y. Chen, and J.P. Ma, Nucl. Phys. **B624** (2002) 360.
- [7] M. Lüscher, Commun. Math. Phys. **105** (1986) 135; Commun. Math. Phys. **104** (1986) 177; *Selected topics in lattice theory*, Lectures given at Les Houches (1988); Nucl. Phys. **B354** (1991) 531.
- [8] JLQCD Collaboration (S. Aoki *et.al.*), Nucl. Phys. **B** (Proc. Suppl.) **53** (1997) 355.
- [9] C. Bernard and M. Golterman, Phys. Rev. **D53** (1996) 476.

FIGURES




FIG. 1. The ratio $R(t) = G(t)/D(t)$ at $\beta = 6.3$ and $\kappa = 0.1513$ corresponding to $m_\pi = 433(4)\text{MeV}$. The wall sources are located at $t = 13$ and 14 .




FIG. 2. The mass dependence of $a_0/m_\pi(1/\text{GeV}^2)$ at $\beta = 5.9$.




FIG. 3. The mass dependence of $a_0/m_\pi(1/\text{GeV}^2)$ at $\beta = 6.1$.




FIG. 4. The mass dependence of $a_0/m_\pi(1/\text{GeV}^2)$ at $\beta = 6.3$.




FIG. 5. $a_0/m_\pi(1/\text{GeV}^2)$ at the chiral limit at each lattice spacing. The CHPT prediction ($a_0/m_\pi = 2.265(51)$ GeV^2) is also plotted.

TABLES

$\beta = 5.9$				
κ m_π^2 (GeV ²)	Fit	ΔE ($\times 10^{-3}$)	E' ($\times 10^{-5}$)	a_0/m_π (1/GeV ²)
0.1585	Old	12.4(21)	—	−0.84(12)
0.2529(56)	Exp	20.9(40)	—	−1.29(20)
	Lin	14.5(19)	—	−0.96(11)
	Sqr	23.1(74)	29(21)	−1.40(35)
0.1580	Old	12.5(15)	—	−0.822(84)
0.3468(49)	Exp	19.9(27)	—	−1.20(13)
	Lin	14.0(13)	—	−0.905(72)
	Sqr	19.0(57)	14(15)	−1.16(27)
0.1575	Old	12.1(12)	—	−0.786(65)
0.4396(48)	Exp	18.5(21)	—	−1.108(98)
	Lin	13.3(11)	—	−0.849(56)
	Sqr	16.3(50)	8(13)	−1.00(24)
0.1570	Old	11.5(10)	—	−0.743(55)
0.5337(49)	Exp	17.0(17)	—	−1.017(79)
	Lin	12.48(92)	—	−0.794(47)
	Sqr	14.4(45)	5(12)	−0.89(22)
0.1565	Old	10.86(91)	—	−0.698(48)
0.6297(50)	Exp	15.6(15)	—	−0.931(67)
	Lin	11.69(82)	—	−0.741(42)
	Sqr	13.0(41)	3(10)	−0.81(20)
0.1560	Old	10.19(82)	—	−0.654(43)
0.7279(51)	Exp	14.2(13)	—	−0.855(59)
	Lin	10.92(75)	—	−0.692(38)
	Sqr	11.9(37)	2.6(95)	−0.74(19)

TABLE I. The results at $\beta = 5.9$. Four lines for each m_π are results with the fitting functions **Old**, **Exp**, **Lin**, and **Sqr**, which are defined in (5)–(7).

$\beta = 6.1$				
κ m_π^2 (GeV ²)	Fit	ΔE ($\times 10^{-3}$)	E' ($\times 10^{-5}$)	a_0/m_π (1/GeV ²)
0.15430	Old	8.45(98)	—	−1.13(11)
0.1925(42)	Exp	13.0(17)	—	−1.62(17)
	Lin	9.82(95)	—	−1.28(10)
	Sqr	14.2(37)	9.9(84)	−1.73(36)
0.15415	Old	8.17(79)	—	−1.080(89)
0.2329(42)	Exp	12.2(13)	—	−1.51(13)
	Lin	9.38(76)	—	−1.214(82)
	Sqr	13.0(33)	8.2(74)	−1.59(32)
0.15400	Old	7.88(71)	—	−1.035(79)
0.2732(42)	Exp	11.6(11)	—	−1.42(11)
	Lin	8.97(68)	—	−1.154(72)
	Sqr	12.2(30)	7.2(68)	−1.48(29)
0.15370	Old	7.38(62)	—	−0.960(68)
0.3539(44)	Exp	10.43(93)	—	−1.274(89)
	Lin	8.23(59)	—	−1.056(62)
	Sqr	11.0(26)	6.2(60)	−1.33(25)
0.15340	Old	6.96(56)	—	−0.987(60)
0.4355(46)	Exp	9.56(80)	—	−1.164(76)
	Lin	7.73(53)	—	−0.980(55)
	Sqr	10.2(24)	5.5(55)	−1.22(22)
0.15300	Old	6.48(49)	—	−0.831(53)
0.5465(49)	Exp	8.65(68)	—	−1.050(65)
	Lin	7.12(47)	—	−0.898(48)
	Sqr	9.3(21)	4.7(49)	−1.11(20)
0.15250	Old	5.96(43)	—	−0.760(45)
0.6897(52)	Exp	7.73(58)	—	−0.938(56)
	Lin	6.48(42)	—	−0.814(42)
	Sqr	8.2(19)	3.9(43)	−0.99(18)
0.15200	Old	5.48(40)	—	−0.697(41)
0.8385(55)	Exp	6.95(52)	—	−0.845(50)
	Lin	5.92(38)	—	−0.743(39)
	Sqr	7.4(17)	3.1(39)	−0.89(16)

TABLE II. The results at $\beta = 6.1$. Four lines for each m_π are results with the fitting functions **Old**, **Exp**, **Lin**, and **Sqr**, which are defined in (5)–(7).

$\beta = 6.3$				
κ m_π^2 (GeV ²)	Fit	ΔE ($\times 10^{-3}$)	E' ($\times 10^{-5}$)	a_0/m_π (1/GeV ²)
0.15130	Old	5.97(60)	—	−1.21(11)
0.1876(36)	Exp	8.19(89)	—	−1.58(14)
	Lin	6.71(60)	—	−1.34(10)
	Sqr	7.9(18)	2.4(36)	−1.54(29)
0.15115	Old	5.79(48)	—	−1.160(83)
0.2399(36)	Exp	7.78(71)	—	−1.48(11)
	Lin	6.43(49)	—	−1.267(81)
	Sqr	7.7(14)	2.6(28)	−1.48(22)
0.15100	Old	5.63(42)	—	−1.115(70)
0.2924(36)	Exp	7.42(60)	—	−1.400(93)
	Lin	6.19(42)	—	−1.206(69)
	Sqr	7.3(13)	2.3(24)	−1.39(19)
0.15075	Old	5.33(36)	—	−1.042(59)
0.3815(38)	Exp	6.87(51)	—	−1.282(76)
	Lin	5.80(36)	—	−1.118(58)
	Sqr	6.5(11)	1.5(21)	−1.23(16)
0.15050	Old	5.01(33)	—	−0.973(54)
0.4728(40)	Exp	6.34(45)	—	−1.177(67)
	Lin	5.42(33)	—	−1.038(52)
	Sqr	5.81(99)	0.8(19)	−1.10(15)
0.15000	Old	4.36(30)	—	−0.842(48)
0.6634(45)	Exp	5.37(39)	—	−0.996(58)
	Lin	4.70(30)	—	−0.894(46)
	Sqr	4.72(89)	0.0(17)	−0.90(14)

TABLE III. The results at $\beta = 6.3$. Four lines for each m_π are results with the fitting functions **Old**, **Exp**, **Lin**, and **Sqr**, which are defined in (5)–(7).

β	a (1/GeV)	Old	Exp	Lin	Sqr
5.9	0.493(7)	-0.96(10)	-1.51(16)	-1.093(90)	-1.58(36)
6.1	0.378(6)	-1.185(59)	-1.653(80)	-1.335(55)	-1.78(22)
6.3	0.302(5)	-1.335(76)	-1.745(99)	-1.466(74)	-1.77(21)
	$a \rightarrow 0$	-1.92(25)	-2.09(35)	-2.07(24)	-2.04(78)

TABLE IV. The values of $a_0/m_\pi(1/\text{GeV}^2)$ in the chiral limit for the each fitting function of $R(t)$ at each β and those in the continuum limit obtained by liner extrapolation in the lattice spacing. The fitting functions of $R(t)$ are defined in (5)–(7).