Screening and confinement in $U(1)^{N-1}$ Abelian effective theories

Tsuneo Suzuki¹ and M. N. Chernodub^{1,2}

¹Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan ²Institute of Theoretical and Experimental Physics, ITEP,

B. Cheremushkinskaja 25, Moscow, 117259, Russia

(Dated: November 4, 2018)

We discuss the effect of dynamical charged particles in $U(1)^{N-1}$ Abelian effective theories of QCD. Screening and confinement at zero temperature are explained qualitatively. The effect of dynamically charged particle is expressed in terms of an effective action of integer electric currents. When a screening is expected, the static potential shows a flattening in the long-range region and a linear behavior in the intermediate region. We show why the screening is better observed in the Polyakov-loop correlators than in the Wilson loops. The breaking of the adjoint string is explained without the Z(N) picture.

PACS numbers: 12.38.Aw,11.15.Ha,14.80.Hv

Understanding of non-perturbative aspects of strongly coupled field theories – such as QCD – is always a challenging task. From the study of confinement in QCD, it has been known that duality and topology are two important keywords. Namely 't Hooft idea [1] of monopole condensation after Abelian projection to $U(1)^{N-1}$ of SU(N)QCD has been shown to be successful in the infrared region. Abelian and monopole components seem to be dominant in the long-range region of quenched QCD [2]. An infrared effective monopole action has been derived in the continuum limit after a block-spin transformation of monopole currents [3, 4]. It is a quantum perfect action described by monopole currents. The monopole action shows that monopole entropy dominates over the energy. That is, monopole condensation occurs in the confinement phase. Using a quantum perfect operator [5], we can evaluate the string tension analytically and prove the continuum rotational invariance of the static potential [6].

Although the 't Hooft scenario seems correct in many respects, there remains an unsolved serious problem even in guenched QCD and also in full QCD. It is the screening-confinement problem extensively discussed in recent publications [7]. How can screening be explained in this scheme? Consider a static pair of charge 2 particles in quenched QCD. Since off-diagonal gluons can screen the charge 2 particle, the static potential is expected to show flattening in the long-range region, i.e., screening. Actually, an adjoint Wilson loop and a charge 2 Wilson loop show such behaviors in the strong coupling region. This problem was discussed extensively in Ref.[8]. The theory in terms of Abelian link fields or Abelian monopole currents alone becomes highly non-local if we integrate out off-diagonal gluon fields after an Abelian projection. Namely we have to keep all charge 2 Abelian Wilson loops in the effective action written by Abelian link fields to reproduce the screening of charge 2. Needless to say, such an Abelian effective action is useless. The same problem is more serious in the real full QCD,

since a fundamental charge also is screened. The authors in Ref.[8] suggest that the relevant quantity in the confinement mechanism is not the Abelian monopoles but the Z(N) center-vortices which can explain the screening problem [9].

The aim of this note is to show how the screening and confinement problem is solved qualitatively in the framework of $U(1)^{N-1}$ Abelian dynamics. We find the effect of dynamical charged particles can be described in terms of integer electric currents on the lattice.

Let us first review the case without a dynamical charged particle. For simplicity, we consider only SU(2) QCD $\rightarrow U(1)$ and use the differential forms on the lattice [10].

Consider a modified compact QED action of the form

$$Z_1 = \int_{-\pi}^{\pi} \mathcal{D}\theta \sum_{n \in \mathbb{Z}(c_2)} e^{-\frac{1}{4\pi^2} (\mathrm{d}\theta + 2\pi n, \Delta D(\mathrm{d}\theta + 2\pi n)) + i(QJ,\theta)}, \quad (1)$$

where the operator D is a general differential operator, Qis a charge of an external source and the current J takes ± 1 along the Wilson loop. Δ is the Laplacian on the lattice. Following Ref. [6] one can reduce the partition function (1) to that of the following monopole action

$$Z_{2} = \sum_{\substack{k \in \mathbb{Z}(*c_{1})\\\delta^{*}k = 0}} e^{-(*k,D^{*}k) + 2\pi i(*k,*N) - \pi^{2}Q^{2}(J,(\Delta^{2}D)^{-1}J)},$$
(2)

where $N \equiv \Delta^{-1} \delta^* S$ and S is an open surface spanned on the Wilson loop, i.e., $\delta S \equiv QJ$. Here and below the constant pre-factors are ignored. Numerically it is found [4] that the Abelian effective monopole action derived from SU(2) QCD can be described well in terms of two-point interactions alone in the long-range region as in (2).

It is also possible to transform the monopole partition function (2) into the string representation as follows [6]:

$$Z_3 = \sum_{\substack{\sigma \in \mathbb{Z}(c_2)\\\delta\sigma = QJ}} e^{-\pi^2(\sigma, (\Delta D)^{-1}\sigma)}, \qquad (3)$$

where the summation goes over the surfaces spanned (|Q|)times) on the external current J.

In the infrared limit the operator D is approximated well by Coulomb+self+nearest neighbor terms [4], that is, $D = \beta \Delta^{-1} + \alpha + \gamma \Delta$, where β , α and γ are renormalized coupling constants of the monopole action. They satisfy the relation $\beta \gg \alpha, \gamma$. Then the Wilson loop can be estimated from Eq.(3):

$$\langle W_Q(R,T)\rangle = \text{const.} e^{-\kappa |Q| RT + \cdots},$$
 (4)

where RT is the area of the minimal surface spanned on the contour J. There are |Q| such surfaces which must be parallel to each other [11] to maximize the contribution for $|Q| \ge 2$. This explains linearity in |Q| in Eq. (4). When $\beta \gg \alpha, \gamma$, the string tension is evaluated approximately as $\kappa = \pi^2/\beta$.

Now let us introduce a dynamical charged particle. A charged vector field corresponding to gluon fields and a fermion field corresponding to quark fields are rather complicated to deal with. Hence we consider first a charged scalar Higgs field in the London limit as a simple example. The radial part ρ_x of the scalar Higgs field $\Phi_x = \rho_x e^{i\vartheta_x}$ is frozen and the dynamical variable is the compact Higgs phase $\vartheta \in [-\pi, \pi)$ which carries the electric charge q. The Higgs field action is written in the Villain representation:

$$Z_4[\theta] = \int_{-\pi}^{\pi} \mathcal{D}\vartheta \sum_{l \in \mathbb{Z}(c_1)} e^{-(\mathrm{d}\vartheta + q\theta + 2\pi l, G(\mathrm{d}\vartheta + q\theta + 2\pi l))}, \quad (5)$$

where G is a local operator.

The integration of the phase of the Higgs field ϑ can be represented as a weighted sum of the Wilson loop over the closed charged currents following Ref. [12]:

$$Z_{4} = \int_{-\pi}^{\pi} \mathcal{D}\vartheta \int_{-\infty}^{\infty} \mathcal{D}F \sum_{l \in \mathbb{Z}(c_{1})} e^{-(F,(4G)^{-1}F) + i(F, \mathrm{d}\vartheta + q\theta + 2\pi l)}$$
$$= \int_{-\pi}^{\pi} \mathcal{D}\vartheta \sum_{j \in \mathbb{Z}(c_{1})} e^{-(j,(4G)^{-1}j) + i(j, \mathrm{d}\vartheta + q\theta)}$$
(6)

$$= \sum_{\substack{j \in \mathbb{Z}(c_1) \\ \delta j = 0}}^{j \in \mathbb{Z}(c_1)} e^{-(j,(4G)^{-1}j) + i(qj,\theta)},$$
(7)

where we consecutively introduce the Gaussian integration over the non-compact link variable F, apply the Poisson summation formula, $\sum_{l} e^{2\pi i (l,F)} = \sum_{j} \delta(j-F)$, integrate over the fields F and ϑ . The last integration gives the closeness constraint $\delta j = 0$.

Hence the Villain type compact QED with the charged scalar field is written as

$$Z_{5} = \sum_{\substack{j \in \mathbb{Z}(c_{1}) \\ \delta j = 0}} e^{-(j,(4G)^{-1}j)} \int_{-\pi}^{\pi} \mathcal{D}\theta \sum_{n \in \mathbb{Z}(c_{2})} (8)$$
$$e^{-\frac{1}{4\pi^{2}}(\mathrm{d}\theta + 2\pi n, \Delta D(\mathrm{d}\theta + 2\pi n)) + i(QJ + qj, \theta)}.$$

This expression can be reduced further to the monopoleelectric and the string-electric current models:

$$Z_{6} = \sum_{\substack{j \in \mathbb{Z}(c_{1}) \\ \delta j = 0}} \sum_{\substack{k \in \mathbb{Z}(^{*}c_{1}) \\ \delta^{*}k = 0}} e^{-(^{*}k,D^{*}k) + 2\pi i(^{*}k,^{*}N + ^{*}n)}$$
$$= \sum_{\substack{j \in \mathbb{Z}(c_{1}) \\ \delta j = 0}} \sum_{\substack{\sigma \in \mathbb{Z}(c_{2}) \\ \delta \sigma = QJ + qj}} e^{-(j,(4G)^{-1}j) - \pi^{2}(\sigma,(\Delta D)^{-1}(\sigma))}, (9)$$

where $*n \equiv \Delta^{-1}\delta^*s$ and s is a surface spanned q times on the dynamical current j, *i.e.*, $\delta s \equiv qj$.

From Eq.(9), we can see how the screening-confinement problem is solved. If $Q/q \notin \mathbb{Z}$, the static potential can not be screened completely. For example, in the case of Q = 1 and q = 2, the leading string tension is equivalent to the string tension without presence of the dynamical charges as in (4). In this case, j = 0 in the sum over j gives the leading term while other terms coming from non-zero j show stronger damping.

On the other hand, if $Q/q \equiv N \in \mathbb{Z}$ (as in the case of Q = 1 and q = 1) then the expectation value of the $R \times T$ Wilson loop is expanded as a perimeter term ($\propto (R+T)$) given by the action of the dynamical screening current with QJ + qj = 0 and the area-law term:

$$\langle W_Q(R,T)\rangle = c_0 e^{-mN^2(R+T)}$$
(10)
+ $c_1 e^{-\kappa q RT - m(N-1)^2(R+T)} + \cdots$.

These terms correspond to the screening currents, respectively, j = -NJ and j = (-N+1)J. The string breaking is seen at large distances. It is important to note that the area-law behavior is observed in the intermediate region even when the screening occurs.

One can easily show why the screening is better observed in the case of the Polyakov loop correlator rather than in the case of the Wilson loop quantum average as found recently [7]. Consider two Polyakov loops separated by a distance R corresponding to a pair of static quark and anti-quark in the case of periodic boundary condition in the time direction. The two leading terms in the average are:

$$\langle P_Q(0)P_{\bar{Q}}(R)\rangle = d_0 e^{-mN^2T} + d_1 e^{-\kappa qRT - m(N-1)^2T} + \cdots$$

A comparison of this expression with Eq.(10) gives that the area-law terms are the same for both averages. However, the perimeter terms are different: for the Wilson loop the perimeter term contains additional suppression factor $\sim e^{-m(2N-1)R}$ with respect to the area term. Thus at sufficiently large separations between the sources the perimeter term in the Wilson loop average may not be found numerically. As a result, the string breaking may not be observed in the Wilson loop even if the breaking can be seen in the Polyakov loop correlator. To derive numerically such an effective action (9) in terms of monopole and electric currents is very interesting. Integer monopoles can be defined following DeGrand-Toussaint[13] while definition of the integer electric currents in terms of original Abelian link fields seems to be impossible. Nevertheless, a given effective matter action as in (5), the action in terms of integer electric currents (9) can also be derived.

Now we consider the case of dynamical charged link variables which correspond to off-diagonal gluon fields after the Abelian projection of QCD. The SU(2) link variable can be parameterized as $U_{x,\mu} = c_{x,\mu}u_{x,\mu}$ where the matrix $c_{x,\mu}$ corresponds to the off-diagonal gluon while $u_{x,\mu}$ represents the diagonal photon contribution: $c_{x,\mu}^{11} = c_{x,\mu}^{22} = \cos \eta_{x,\mu}, c_{x,\mu}^{12} = c_{x,\mu}^{21*} = i \sin \eta_{x,\mu} e^{i\varphi_{x,\mu}},$ $u_{x,\mu}^{11} = u_{x,\mu}^{22*} = e^{i\theta_{x,\mu}}$ and $u_{x,\mu}^{12} = u_{x,\mu}^{21} = 0$. The link fields are restricted, $\eta \in [0, \pi/2], \theta, \varphi \in [-\pi, \pi)$.

The meaning of these fields can be understood after applying the U(1) gauge transformations, which are allowed in an Abelian projection. The field θ is the Abelian gauge field and φ is the vector field (phase of the offdiagonal gluon field) with the electric charge two, respectively: $\theta_{x,\mu} \to \theta_{x,\mu} + \xi_x - \xi_{x+\hat{\mu}}, \varphi_{x,\mu} \to \varphi_{x,\mu} + 2\xi_x$. In the continuum limit these transformations reduce to $\theta_{\mu}(x) \to \theta_{\mu}(x) - \partial_{\mu}\xi(x)$ and $\varphi_{\mu}(x) \to \varphi_{\mu}(x) + 2\xi(x)$, respectively. The variable $\eta_{x,\mu}$ is U(1) gauge invariant.

The SU(2) plaquette action $\operatorname{Tr} U_P$ in terms of the angles η , θ and φ can be written as a sum of three parts [14]. The first part is proportional to the compact QED action, $S^{\theta} = -\beta_{SU(2)} \cos d\theta \prod_{i=1}^{4} \cos \eta_i$, where $\beta_{SU(2)}$ is the SU(2) gauge coupling and the subscripts 1, ..., 4 correspond to the standard notations of the plaquette links: $1 \to \{x, x + \hat{\mu}\}, ..., 4 \to \{x, x + \hat{\nu}\}$. The second part, $S^{\theta\varphi}$, contains 6 terms representing the interaction of the matter field φ with the gauge field θ . All these terms are proportional to the products $\cos \eta_1 \cos \eta_2 \sin \eta_3 \sin \eta_4$ and the like. The last part of the action is given by $S^{\varphi} = -\beta_{SU(2)} \cos \tilde{d}(\varphi + \theta) \prod_{i=1}^{4} \sin \eta_i$, where $\tilde{d}\varphi = \varphi_1 - \varphi_2 + \varphi_3 - \varphi_4$.

The simplest way to demonstrate the adjoint string breaking is to use a mean-field approximation for the field η (for our purposes the fluctuations of this field are not essential). We set $\cos \eta_{x,\mu} \rightarrow \langle \cos \eta_{x,\mu} \rangle = c$ and $\sin \eta_{x,\mu} \rightarrow \langle \sin \eta_{x,\mu} \rangle = s$, respectively. Then the self-interactions of the gauge and matter fields can be written in a short form, $S^{\theta} = -\beta_{SU(2)}c^4 \cos d\theta$ and $S^{\varphi} = -\beta_{SU(2)}s^4 \cos \tilde{d}(\varphi + \theta)$, respectively.

The role of the actions $S^{\theta\varphi}$ and S^{φ} can be elucidated by noticing that they basically contains the following combinations of the fields: self-interaction of the gauge field, $d\theta$, the interaction of the gauge field with the off-diagonal field, $H_{x,\mu\nu} \equiv 2\theta_{x,\mu} + \varphi_{x,\nu} - \varphi_{x+\hat{\mu},\nu}$ and self-interaction of the off-diagonal field, $C_{x,\mu\nu} \equiv \varphi_{x,\mu} - \varphi_{x,\nu}$. Thus from the bare Wilson gauge action we get the following mean-field types of interactions involving the off-diagonal fields:

$$S^{\theta\varphi}_{x,\mu\nu} + S^{\varphi}_{x,\mu\nu} \longrightarrow \cos H_{x,\mu\nu} + \cos H_{x,\nu\mu} + \cos C_{x,\mu\nu}.$$

The tensor $H_{\mu\nu}$ is not antisymmetric contrary to $C_{\mu\nu}$. Therefore we need two terms with H at each plaquette.

A general effective action of charged link variables can be written in terms of two periodic actions:

$$\tilde{S}^{\theta\varphi}_{x,\mu\nu} = F_H(H) + F_H(H') + F_C(C), \qquad (11)$$

where the form $H \equiv H_{x,\mu\nu}$ $(H' \equiv H_{x,\nu\mu})$ corresponds to the positively (negatively) oriented plaquette interactions of the matter and gauge fields. Now let us consider the adjoint string breaking in the SU(2) gluodynamics using the following partition function:

$$Z_7 = \int_{-\pi}^{\pi} \mathcal{D}\theta \int_{-\pi}^{\pi} \mathcal{D}\varphi e^{-\sum_P \left(S_P^{\theta}(\theta) + \tilde{S}_P^{\theta\varphi}(\theta,\varphi)\right)} .$$
(12)

The screening of the external charge-2 (adjoint) Wilson loops is possible essentially due to the presence of the off-diagonal field action $\tilde{S}^{\theta\varphi}$. To show this we apply the duality transformation with respect to this action:

$$e^{-\sum_{P} \tilde{S}_{P}^{\theta\varphi}(\theta,\varphi)} = \sum_{\substack{n_{i} \in \mathbb{Z}(c_{2})\\i=1,2,3}} I_{n_{1}}^{H} I_{n_{2}}^{H} I_{n_{3}}^{C} e^{i(H,n_{1})+i(H',n_{2})+i(C,n_{3})} ,$$

where the weights $I_{n_i}^l$ correspond to the terms in Eq. (11).

The integration of Eq. (12) over φ gives us the sum over the particle trajectories with the charge 2:

$$\int_{-\pi}^{\pi} \mathcal{D}\varphi \, e^{-\sum_{P} \tilde{S}_{P}^{\theta\varphi}(\theta,\varphi)} = \sum_{\substack{j \in \mathbb{Z}(c_{1})\\\delta j = 0}} w(j) \, e^{2i(j,\theta)} \,, \qquad (13)$$

where w(j) is the weight corresponding to the ensemble of the Wilson loops $\{j\}$. The integration over the field φ provide us with constraints on the dynamical integer trajectories j: the trajectories must be closed, $\delta j = 0$, due to gauge invariance of both parts of Eq. (13).

The final step is to substitute the loop expansion (13) into the partition function (12) and to apply the arguments identical to those in the charge Q = 2 scalar particle considered earlier. Thus we get the screening of the charge-2 external charges (gluons) and, as a result, the breaking of the adjoint string.

Let us next discuss the case of charged fermions corresponding to quark fields. The fermion determinant can be rewritten as a sum of self-avoiding Wilson loops composed of charge q = 1 particle[15, 16]. Hence we get an action similar to (9) and the screening-confinement problem could be solved similarly.

But generally an effective fermionic action becomes more complicated and moreover it is desirable to adopt a form of action which makes numerical calculations possible. We employ a Polynomial Hybrid Monte Carlo Method [17]. Consider a polynomial $P_{N_{poly}}[z]$ with an even degree N_{poly} which approaches z^{-1} as N_{poly} increases. We get

$$\det[P_{N_{poly}}[D]] = \det\left[T_{N_{poly}}^{\dagger}[D]T_{N_{poly}}[D]\right]$$
(14)

where $T_{N_{poly}}[z]$ is a polynomial with a degree $N_{poly}/2$. Hence the fermionic determinant for N_f odd-number of flavors is written using a pseudo-fermion representation:

$$(\det[D])^{N_f} = \left[\det[DP_{N_{poly}}[D]]\right]^{N_f}$$
(15)

$$\times \int \mathcal{D}\phi^{\dagger} \mathcal{D}\phi \exp\left[-\phi^{\dagger}T^{\dagger}_{N_{poly}}[D]T_{N_{poly}}[D]\phi\right].$$

Since $T_{N_{poly}}[z]$ is a polynomial of degree $N_{poly}/2$, then

$$T^{\dagger}_{N_{poly}}[D]T_{N_{poly}}[D] = a_0 + a_1 M + a_1^* M^{\dagger} + \cdots, \quad (16)$$

where M is a hopping parameter term in D. Since the higher-order terms are irrelevant in the screening problem, we consider only the linear term with respect to M. Here we adopt the Wilson fermion action for M coupled to a U(1) compact gauge field as an example:

$$M = \kappa \sum_{\mu=1}^{4} \{ (1 - \gamma_{\mu}) \begin{pmatrix} e^{i\theta_{x,\mu}} & 0\\ 0 & e^{-i\theta_{x,\mu}} \end{pmatrix} \delta_{x+\mu,x'} + (1 + \gamma_{\mu}) \begin{pmatrix} e^{-i\theta_{x-\mu,\mu}} & 0\\ 0 & e^{i\theta_{x-\mu,\mu}} \end{pmatrix} \delta_{x-\mu,x'} \}$$
(17)

We separate the U(1) variant phase factor from the U(1) invariant term $\tilde{\phi}_{\alpha}(x)$ of the pseudo-fermion field $\phi_{\alpha}(x) = \tilde{\phi}_{\alpha}(x) \exp(i\chi_{x,\alpha}\tau_3)$ (α =Dirac indices). Regarding the field $\tilde{\phi}$ as a constant we get the following mean-field types of interactions from the Wilson-fermion action:

$$\sum_{\substack{\alpha,\mu,x\\\alpha,\beta,\mu,x}} \cos(\Delta_{\mu}\chi_{x,\alpha} + \theta_{x,\mu}),$$
$$\sum_{\substack{\chi,\beta,\mu,x\\\alpha,\beta,\mu,x}} \cos(\chi_{x+\mu,\beta} - \chi_{x,\alpha} + \theta_{x,\mu})\bar{\gamma}_{\alpha\beta}(\mu)$$

where $\bar{\gamma}(\mu)$ are realized Euclidean gamma matrices, *i.e.*, $\bar{\gamma}(\mu) = i\gamma_{\mu}$ for $\mu = 1, 3$ and $\bar{\gamma}(\mu) = \gamma_{\mu}$ for $\mu = 2, 4$.

As an effective fermionic action, it is natural to assume the following periodic pseudo-fermion action:

$$e^{-S_{f}(\theta,\chi)} = e^{F(\Delta_{\mu}\chi_{x,\alpha} + \theta_{x,\mu}) + G(\chi_{x+\mu,\beta} - \chi_{x,\alpha} + \theta_{x,\mu})\bar{\gamma}_{\alpha\beta}(\mu)}$$

$$= \sum_{j_{\alpha,\mu}(x)\in Z} \sum_{j_{\alpha\beta,\mu}(x)\in Z} \tilde{F}(j_{\alpha,\mu})\tilde{G}(j_{\alpha\beta,\mu}) \quad (18)$$

$$\times e^{ij_{\alpha,\mu}(x)(\Delta_{\mu}\chi_{x,\alpha} + \theta_{x,\mu})}$$

$$\times e^{ij_{\alpha\beta,\mu}(x)(\chi_{x+\mu,\beta} - \chi_{x,\alpha} + \theta_{x,\mu})\bar{\gamma}_{\alpha\beta}(\mu)}$$

where F and G are periodic functions. In Eq. (18) we have used the Fourier (duality) transformation. The integer current $j_{\mu}(x) \equiv \sum_{\alpha} j_{\alpha,\mu}(x) + \sum_{\alpha,\beta} j_{\alpha\beta,\mu}(x) \bar{\gamma}_{\alpha\beta}(\mu)$ is coupled to the field θ and carries unit electric charge. Due to the gauge invariance of the fermion determinant the integration of Eq. (18) over the phases χ gives us a conservation constraint for the currents, $\delta j = 0$. Thus

$$\int_{-\pi}^{\pi} \mathcal{D}\chi \, e^{-S_f(\theta,\chi)} = \sum_{\substack{j \in \mathbb{Z}(c_1)\\\delta j = 0}} w_F(j) \, e^{i(j,\theta)} \,, \tag{19}$$

where w_F is a weight for the electric particle trajectories.

With the gauge field action (1), we can see that the screening and the string breaking in full QCD also is successfully explained as done in the case of charged scalar particle in the framework of U(1) theory. It is again stressed that the area-law behavior is seen in the intermediate regions even in full QCD. Such an area-law behavior can be explained by the dual Meissner effect due to the monopole condensation. Confinement is important also in full QCD.

The authors are grateful to M. I. Polikarpov for useful discussions. M. N. Ch. is supported by JSPS fellowship P01023 and T. S. is supported by JSPS Grant-in-Aid for Scientific Research on Priority Areas 13135210.

- [1] G. 't Hooft, Nucl. Phys. B 190, 455 (1981).
- T. Suzuki and I. Yotsuyanagi, Phys. Rev. D42, 4257 (1990);
 T. Suzuki, in *Continuous Advances in QCD 1996* (World Scientific, 1997), p. 262.
- [3] H. Shiba and T. Suzuki, Phys. Lett. B351, 519 (1995).
- [4] S. Kato *et al.*, Nucl. Phys. B **520**, 323 (1998).
- [5] S. Fujimoto, S. Kato and T. Suzuki, Phys. Lett. B 476, 437 (2000).
- [6] M. N. Chernodub et al, Phys. Rev. D62, 094506 (2000).
- [7] V. G. Bornyakov et al, talk at Lattice 2002, Cambridge, USA, June, 2002; C. W. Bernard et al., Phys. Rev. D 64, 074509 (2001); A. Duncan, E. Eichten and H. Thacker, Phys. Rev. D 63, 111501 (2001); B. Bolder et al., Phys. Rev. D 63, 074504 (2001); H. D. Trottier, Phys. Rev. D 60, 034506 (1999); C. DeTar et al., Phys. Rev. D 59, 031501 (1999).
- [8] J. Ambjørn et al, JHEP 0002, 033 (2000)
- [9] G. 't Hooft, Nucl. Phys. B **138**, 1 (1978).
- [10] A. H. Guth, Phys. Rev. D 21, 2291 (1980); see also the review by M. N. Chernodub and M. I. Polikarpov, in *Confinement, Duality, and Nonperturbative Aspects of QCD*, Ed. by Pierre van Baal (Plenum, New York 1998), p. 387.
- [11] M.Zach, M.Faber, P.Skala, Nucl.Phys.B 529, 505 (1998).
- [12] M.B.Einhorn and R.Savit, Phys. Rev. D 17, 2583 (1978);
 19, 1198 (1979); M. N. Chernodub and M. I. Polikarpov, unpublished (1997).
- [13] T. A. DeGrand, D. Toussaint, Phys. Rev. D 22, 2478 (1980).
- [14] M. N. Chernodub, M. I. Polikarpov and A. I. Veselov, Phys. Lett. B 342, 303 (1995).
- [15] M. Karowski, R. Schrader and H. J. Thun, Comm. Math. Phys. 97, 5 (1985).
- [16] S. Hands and J. B. Kogut, Nucl. Phys. B 462, 291 (1996).
- [17] Ph. de Forcrand and T. Takaishi, Nucl. Phys. B(Proc. Suppl.) 53, 968 (1997); R. Frezzotti and K. Jansen, Phys. Lett. B402, 328 (1997); C. Alexandrou *et al.*, Phys. Rev. D60, 034504 (1999); S. Aoki *et al.*, hep-lat/0112051.