

**LATTICE QCD STUDY FOR
THE QUARK-GLUON MIXED CONDENSATE $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$**

TAKUMI DOI¹, NORIYOSHI ISHII², MAKOTO OKA¹ AND
HIDEO SUGANUMA¹

¹ *Dept. of Phys., Tokyo Institute of Technology, Meguro, Tokyo 152-8551, Japan*
E-mail: doi@th.phys.titech.ac.jp

² *Radiation Laboratory, RIKEN, Hirosawa 2-1, Wako, 351-0198, Japan*

We study the quark-gluon mixed condensate $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$, which is another chiral order parameter, using the $SU(3)_c$ lattice QCD with the Kogut-Susskind(KS) fermion at the quenched level. We generate 100 gauge configurations on the 16^4 lattice with $\beta = 6.0$, and measure $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ at 16 points in each gauge configuration for each current-quark mass of $m_q = 21, 36, 52$ MeV. From the 1600 data for each m_q , we find $m_0^2 \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle/\langle\bar{q}q\rangle \simeq 2.5$ GeV² at the lattice scale of $a^{-1} \simeq 2$ GeV in the chiral limit. The large value of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ suggests its importance in the operator product expansion(OPE) in QCD.

1. The importance of the quark-gluon mixed condensate

In order to understand the non-perturbative structure of the QCD vacuum, condensates are important quantities. Among various condensates, we emphasize the importance of the quark-gluon mixed condensate $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$. First, the mixed condensate represents a direct correlation between quarks and gluons in the QCD vacuum even at the qualitative level. Second, the mixed condensate is another chiral order parameter, which flips the chirality of the quark. Third, the mixed condensate plays an important role in various QCD sum rules, especially in the baryons¹, the light-heavy mesons² and the exotic mesons³. In the QCD sum rules, the value $m_0^2 \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle/\langle\bar{q}q\rangle \simeq 0.8 \pm 0.2$ GeV² has been proposed phenomenologically⁴. However, there was only one pioneering but preliminary lattice QCD work⁵, performed with very little statistics (only 5 data) using a small (8^4) and coarse lattice ($\beta = 5.7$). Therefore, we present the calculation for $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ in the $SU(3)_c$ lattice QCD with a larger (16^4) and finer ($\beta = 6.0$) lattice and with high statistics (1600 data), using the KS-fermion at the quenched level. With this high statistics, we perform re-

liable estimate for $m_0^2 \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle/\langle\bar{q}q\rangle$ at the lattice scale in the chiral limit.

2. Formalism and the Results of Lattice QCD

We calculate the condensates $\langle\bar{q}q\rangle$ and $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ using the $SU(3)_c$ lattice QCD with the KS-fermion at the quenched level. The KS-fermion preserves the explicit chiral symmetry for the quark mass $m = 0$. This is essential for our study because both of $\langle\bar{q}q\rangle$ and $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ work as the chiral order parameter. We generate 100 gauge configurations with the standard Wilson action at $\beta = 6.0$ on the 16^4 lattice. The lattice unit⁶ $a \simeq 0.10\text{fm}$ is obtained so as to reproduce the string tension $\sigma = 0.89\text{GeV}/\text{fm}$. We use $m = 21, 36, 52$ MeV (i.e. $ma = 0.0105, 0.0184, 0.0263$). We calculate the flavor-averaged condensates as

$$a^3\langle\bar{q}q\rangle = -\frac{1}{4}\sum_f \text{Tr} [\langle q^f(x)\bar{q}^f(x)\rangle], \quad (1)$$

$$a^5 g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle = -\frac{1}{4}\sum_f \sum_{\mu,\nu} \text{Tr} [\langle q^f(x)\bar{q}^f(x)\sigma_{\mu\nu}G_{\mu\nu}\rangle], \quad (2)$$

where $SU(4)_f$ quark-spinor field q , \bar{q} is converted into spinless Grassmann KS-fields χ , $\bar{\chi}$ and the gauge-link variable⁷. We adopt the clover-type definition of the gluon field strength $G_{\mu\nu}$ on the lattice as

$$G_{\mu\nu}^{\text{lat}}(s) = \frac{i}{16}\sum_A \sum_{\substack{s'=s,s-\mu, \\ s-\nu,s-\mu-\nu}} \lambda^A \text{Tr} [\lambda^A \{U_{\mu\nu}(s') - U_{\nu\mu}(s')\}], \quad (3)$$

which has no $O(a)$ discretization error. We measure the condensates on 16 different space-time points x in each configuration. For each m , we take the average over the 16 space-time points and 100 gauge configurations.

Figure 1 shows the bare condensates $\langle\bar{q}q\rangle$ and $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ against the quark mass ma . Note that the jackknife errors are almost negligible due to the high statistics of 1600 data for each m . We fit the data linearly and determine the condensates in the chiral limit. The obtained data are summarized in Table 1. We also check the finite volume artifact, by imposing the anti-periodic/periodic boundary conditions on $\chi, \bar{\chi}$. The two results with different boundary conditions almost coincide within about 1% deviation and we conclude that the physical volume $V \sim (1.6\text{ fm})^4$ in our simulations is large enough to avoid the finite volume artifact⁷.

The values of the condensates in the continuum limit are to be obtained through the renormalization, which, however, suffers from uncertainty of




Figure 1. The bare condensates $\langle \bar{q}q \rangle$ and $g\langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle$ plotted against the quark mass ma . The dashed lines denote the best linear extrapolations, and the cross symbols the values in the chiral limit. The jackknife errors are hidden in the circles.

Table 1. The numerical results of $\langle \bar{q}q \rangle$ and $g\langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle$ for various ma . The last column denotes their values in the chiral limit by the chiral extrapolation.

	$ma = 0.0263$	$ma = 0.0184$	$ma = 0.0105$	chiral limit
$a^3 \langle \bar{q}q \rangle$	-0.042397(16)	-0.032470(15)	-0.022124(16)	-0.008721(17)
$a^5 g \langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle$	-0.018820(15)	-0.014979(14)	-0.010884(14)	-0.005652(14)

the non-perturbative effect. As a more reliable quantity, we provide the ratio $m_0^2 \equiv g\langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle / \langle \bar{q}q \rangle$, which is free from the uncertainty from the wave function renormalization of the quark. Now, we present the result of m_0^2 using our bare results of lattice QCD as

$$m_0^2 \equiv g\langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle / \langle \bar{q}q \rangle \simeq 2.5 \text{ GeV}^2 \quad (\beta = 6.0 \text{ or } a^{-1} \simeq 2\text{GeV}). \quad (4)$$

We see that m_0^2 is rather large, which suggests the importance of the mixed condensate in OPE. Note that this bare result is determined very precisely⁷.

Finally, we again emphasize that the mixed condensate $g\langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle$ plays very important roles in various contexts in quark hadron physics. In particular, the thermal effect is interesting because the mixed condensate is another chiral order parameter. Considering the importance of finite-temperature QCD in the RHIC project, we are in progress for the study of the mixed condensate $g\langle \bar{q}\sigma_{\mu\nu}G_{\mu\nu}q \rangle$ at finite temperature⁸.

References

1. H.G. Dosch, M. Jamin and S. Narison, *Phys. Lett.* **B220**, 251 (1989).
2. H.G. Dosch and S. Narison, *Phys. Lett.* **B417**, 173 (1998)
3. J.I. Latorre, P. Pascual and S. Narison, *Z. Phys.* **C34**, 347 (1987).
4. V.M. Belyaev and B.L. Ioffe, *Sov. Phys. JETP* **56**, 493 (1982).
5. M. Kremer and G. Schierholz, *Phys. Lett.* **B194**, 283 (1987).
6. T.T. Takahashi et al., *Phys. Rev.* **D65**, 114509 (2002).
7. T. Doi, N. Ishii, M. Oka and H. Suganuma, hep-lat/0211039 (2002).
8. T. Doi, N. Ishii, M. Oka and H. Suganuma, hep-lat/0212006 (2002).