Lattice QCD Calculation of the Kaon B-parameter with the Wilson Quark Action

JLQCD Collaboration

S. Aoki^a, M. Fukugita^b, S. Hashimoto^c, N. Ishizuka^a, Y. Iwasaki^{a,d}, K. Kanaya^{a,d}, Y. Kuramashi^e, M. Okawa^e, A. Ukawa^a, T. Yoshié^{a,d}

> Institute of Physics, University of Tsukuba^a), Tsukuba, Ibaraki 305, Japan

Institute for Cosmic Ray Research, University of Tokyo^b, Tanashi, Tokyo 188, Japan

 $Computing Research Center^c$, High Energy Accelerator Research Organization(KEK), Tsukuba, Ibaraki 305, Japan

Center for Computational Physics, University of Tsukuba^d), Tsukuba, Ibaraki 305, Japan

Institute of Particle and Nuclear Studies^e), High Energy Accelerator Research Organization(KEK), Tsukuba, Ibaraki 305, Japan

Abstract

The kaon B parameter is calculated in quenched lattice QCD with the Wilson quark action. The mixing problem of the $\Delta s = 2$ four-quark operators is solved non-perturbatively with full use of chiral Ward identities, and this method enables us to construct the weak four-quark operators exhibiting good chiral behavior. We find $B_K(\text{NDR}, 2\text{GeV}) = 0.562(64)$ in the continuum limit, which agrees with the value obtained with the Kogut-Susskind quark action.

Reliable knowledge of the $K^0 - \bar{K}^0$ transition matrix element B_K is an indispensable ingredient for further advancement in CP violation phenomenology, and the evaluation of this B_K parameter has been one of the major targets for lattice QCD calculations. Indeed, lots of effort has been expended towards this end.The successful calculations of B_K so far achieved [[1, 2\]](#page-7-0) exclusively employ the Kogut-Susskind quark action that respects chiral $U(1)$ symmetry. Whereas the verification whether both Wilson and Kogut-Susskind quark actions yield the identical result for B_K is an important step to give a full credit to the lattice QCD calculation, the attempts made with the Wilson quark action have not yielded much success $[3, 4, 5]$ $[3, 4, 5]$ $[3, 4, 5]$ $[3, 4, 5]$ $[3, 4, 5]$ $[3, 4, 5]$: the use of the Wilson action with explicit breaking of chiral symmetry causes mixing among four-quark operators of different chiral structure, and ensuring the correct chiral behavior of the $\Delta s = 2$ operators is substantially more complicated than with the Kogut-Susskind action. Early studies have shown that the mixing problem is not adequately treated by perturbation theory, leading to an "incorrect answer" for the matrix element [\[3](#page-7-0)]. Attempts were then made to solve the mixing problem non-perturbatively with the aid of chiral perturbation theory[[4](#page-7-0)]. Unfortunately, they were not successful since the calculation contains large systematic uncertainties arising from higher order effects that survive even in the continuum limit. More recently a proposal is made[[5](#page-7-0)] to improve the chiral behavior of the $\Delta s = 2$ operator with the use of non-perturbative renormalization (NPR) [\[6](#page-7-0)]. The mixing coefficients derived in this approach, however, depend sharply on the momentum scale where the operator is to be evaluated, and this causes a subtlety in estimating the matrix element.

In this Letter we propose a non-perturbative method to solve the operator mixing problem with use of chiral Ward identities [[7](#page-7-0)]. This method fully incorporates the chiral properties of the Wilson action, yielding the $\Delta s = 2$ operator that shows good chiral behavior. No effective theories are invoked to estimate the matrix element. The final result we obtained is encouraging: it shows good agreement with the result with the Kogut-Susskind quark action. We shall also revisit the perturbative method.

Let us consider a set of weak operators in the continuum $\{\hat{O}_i\}$ which closes under chiral rotation $\delta^a \hat{O}_i = i c_{ij}^a \hat{O}_j$. These operators are given by linear combinations of a set of lattice operators $\{O_{\alpha}\}\$, as $\hat{O}_i = \sum_{\alpha} Z_{i\alpha} O_{\alpha}$. We choose the mixing coefficients $Z_{i\alpha}$ such that the Green functions of $\{\hat{O}_i\}$ with quarks in the external states satisfy the chiral Ward identity to $O(a)$. This identity can be derived in a standard manner[[7\]](#page-7-0) and takes the form

$$
-2\rho Z_A \langle \sum_x P^a(x)\hat{O}_i(0) \prod_k \tilde{\psi}(p_k) \rangle + c_{ij}^a \langle \hat{O}_j(0) \prod_k \tilde{\psi}(p_k) \rangle -i \sum_l \langle \hat{O}_i(0) \prod_{k \neq l} \tilde{\psi}(p_k) \delta^a \tilde{\psi}(p_l) \rangle + O(a) = 0,
$$
\n(1)

where p_k is the momentum of the external quark, Z_A and $\rho = (m - \delta m)/Z_A$ are constants to be determined from the Ward identities for the axial vector currents[[8](#page-7-0)], and P^a is the pseudoscalar density of flavor a. We note that the first term comes from the chiral variation of the Wilson quark action and the third represents the chiral rotation of the external fields.

The four-quark operator relevant for B_K is given by $\hat{O}_{VV+AA} = VV + AA$ where $V = \bar{s}\gamma_{\mu}d$ and $A = \bar{s}\gamma_{\mu}\gamma_5d$. Then, $\hat{O}_{VV+AA} = VV + AA$ and $\hat{O}_{VA} = VA$ form a minimal set of the operators that closes under $\lambda^3 = \text{diag}(1, -1, 0)$ chiral rotation. Taking account of CPS symmetry(note that we take $m_d = m_s$ in this article)[[3\]](#page-7-0). mixing of these operators is written $\hat{O}_{VV+AA}/2=\bar{Z}_{VV+AA} (O_0 + z_1O_1 + \cdots + z_4O_4)$ and $\hat{O}_{VA} = Z_{VA} \cdot z_5 O_5$, where the six lattice operators O_i are given by $O_0 =$ $(VV + AA)/2$, $O_1 = (SS + TT + PP)/2$, $O_2 = (SS - TT/3 + PP)/2$, $O_3 =$ $(VV - AA)/2 + (SS - PP)$, $O_4 = (VV - AA)/2 - (SS - PP)$ and $O_5 = VA$ with $S = \bar{s}d$, $P = \bar{s}\gamma_5 d$ and $T = \bar{s}[\gamma_\mu, \gamma_\nu]d/2$ $(\mu < \nu)$. These operators are in the Fierz eigenbasis which we find convenient when taking fermion contractions for evaluating the Green functions in (1).

We consider the four external quarks having an equal momentum p , and denote

by Γ_{VV+AA} and Γ_{VA} the sum of the Green functions on the left hand side of [\(1](#page-2-0)) with external quark legs amputated. Using the projection operator P_i for the Fierz eigenbasis corresponding to O_i , we write $\Gamma_{VV+AA}/Z_{VV+AA} = \Gamma_5 P_5$ and $\Gamma_{VA}/Z_{VA} = \Gamma_0 P_0 + \Gamma_1 P_1 + \cdots + \Gamma_4 P_4$. Writing $\hat{O}_{VV+AA,VA}$ in [\(1](#page-2-0)) in terms of lattice operators, we obtain six equations for the five coefficients z_1, \dots, z_5 :

$$
\Gamma_i = c_0^i + c_1^i z_1 + \dots + c_5^i z_5 = O(a), \quad i = 0, \dots, 5
$$
 (2)

This gives an overconstrained set of equations, and we may choose any five equations to exactly vanish to solve for z_i : the remaining equation should automatically be satisfied to $O(a)$. We choose four equations to be those for $i = 1, \dots, 4$, since O_1, \dots, O_4 do not appear in the continuum. The choice of the fifth equation, $i = 0$ or 5, is more arbitrary. We have confirmed that either $\Gamma_0 = 0$ or $\Gamma_5 = 0$ leads to a consistent result to $O(a)$ for z_1, \dots, z_4 in the region $pa \leq 1$. In the present analysis we choose $\Gamma_5 = 0$. The overall factor Z_{VV+AA} is determined by the NPR method[[6\]](#page-7-0). We convert the matrix elements on the lattice into those of the \overline{MS} scheme in the continuum using naive dimensional regularization (NDR) renormalized at the scale $\mu = 2 \text{GeV}[5]$ $\mu = 2 \text{GeV}[5]$ $\mu = 2 \text{GeV}[5]$:

$$
B_K(\text{NDR}, \mu) = \left(1 + \frac{\alpha_s(\mu)}{4\pi} \left(-4\text{log}\left(\frac{\mu}{p}\right) - \frac{14}{3} + 8\text{log}2\right)\right) \frac{\langle \bar{K}^0 | \hat{O}_{VV+AA} | K^0 \rangle}{\frac{8}{3} |\langle 0 | \hat{A} | K^0 \rangle|^2} \tag{3}
$$

where p denotes the momentum at which the mixing coeffecients are evaluated.

For comparative purpose we also calculate B_K with perturbative mixing coefficients, for which we use the one-loop expression in Ref. [\[9](#page-7-0)] after applying a finite correction in conversion to the NDR scheme together with the tadpole improvement with $\alpha_{\overline{MS}}(1/a)$.

Letus remark here that the equations obtained in the NPR method [[5\]](#page-7-0) corresponds to $\Gamma_i = 0$ for $i = 1, \dots, 4$ in which the contributions of the first and the third term in the Ward identity (1) are dropped, *i.e.*, the full chiral properties are not taken into account in the NPR approach.

Our calculations are made with the Wilson quark action and the plaquette action at $\beta = 5.9 - 6.5$ in quenched QCD. Table [1](#page-9-0) summarizes our run parameters. Gauge configurations are generated with the 5-hit pseudo heat-bath algorithm with $2000(\beta = 5.9 \text{ and } 6.1), 5000(\beta = 6.3) \text{ or } 8000(\beta = 6.5)$ sweep intervals apart. The physical size of lattice is chosen to be approximately constant at $La \approx 2.4$ fm where the lattice spacing is determined from $m_{\rho} = 770$ MeV. At each β four values of the hopping parameter are adopted such that the physical point for the K meson can be interpolated. We estimate $m_s a/2$ from $m_K/m_\rho = 0.648$ for degenerate d and s quark masses. Errors are estimated by the single elimination jackknife method for all measured quantities throughout this work.

Our calculations are carried out in two steps. We first calculate z_i and Z_{VV+AA} using the quark Green functions having finite space-time momenta. For this purpose quark propagators are solved in the Landau gauge for the point source located at the origin with the periodic boundary condition imposed on the lattice. We next extract B_K from the ratio $\langle \bar{K}^0(t = T) \hat{O}_{VV+AA}(t') K^0(t = 1) \rangle / \langle \bar{K}^0(t = 1) \rangle$ $T\hat{A}(t')\rangle/\langle\hat{A}(t')K^0(t=1)\rangle$, each Green function projected onto the zero spatial momentum, by fitting a plateau seen as a function of t' . For this calculation quark propagators are solved without gauge fixing employing the wall source placed at the edge where the Dirichlet boundary condition is imposed in the time direction. We obtain B_K at $m_s/2$ by quadratically interpolating the data at the four hopping parameters.

We plot in Fig. 1 a typical result for the mixing coefficients as a function of the external quark momenta. The plot shows, as desired, only weak dependence of z_i on momentum in the range $0.1 \leq p^2 a^2 \leq 1.0$. This enables us to evaluate the mixing coefficients with small errors at the scale $p^* \approx 2 \text{GeV}$, which always falls within the range of a plateau for our runs at $\beta = 5.9 - 6.5$. We remark that this weak scale dependence contrasts to the strong scale dependence obtained with the NPR method[[5](#page-7-0)].

In Fig. 2 we compare the mixing coefficients evaluated at the scale p^* (filled symbols) with the perturbative values obtained with $\alpha_{\overline{MS}}(1/a)$ (open symbols) as a function of lattice spacing. We remark that a large value of z_2 determined by the Ward identities sharply contrasts with the one-loop perturbative result $z_2 = 0$. For the other coefficients, the perturbative results agree with the non-perturbative ones in sign and rough orders of magnitude. They differ in quantitative details, however.

Let us examine the chiral property of the operator \hat{O}_{VV+AA} by calculating the ratio $\langle \bar{K}^0 | \hat{O}_{VV+AA} | K^0 \rangle / |\langle 0 | \hat{P} | K^0 \rangle|^2$, which vanishes at $m_q = 0$ in the continuum. In Fig. 3 we show the results at $m_q = 0$ obtained by a quadratic extrapolation of data in $m_q = (1/K - 1/K_c)/2$, where WI stands for our method using chiral Ward identities and PT for tadpole-improved one-loop perturbation theory (numbers are given in Table [2](#page-9-0)). The pseudoscalar density \hat{P} in the denominator is renormalized perturbatively for both cases. A significant improvement is clearly seen with use of the Ward identities, the ratio becoming consistent with zero at the lattice spacing $m_{\rho}a\leq 0.3(a\leq 0.08$ fm). In the perturbative approach, the chiral behavior is recovered only after extrapolation to the continuum limit, where we adopted a linear dependence on a expected for the Wilson quark action in the extrapolation shown in Fig. 3.

Our final results for $B_K(NDR, 2{\rm GeV})$ are presented in Fig. 4 as a function of lattice spacing (see Table 2 for numerical details). The method based on the Ward identity (WI) gives a value convergent from a lattice spacing of $m_{\rho}a \approx$ 0.3 ($B_K \sim 0.6 - 0.8$). The large error, however, hinders us from making an extrapolation to the continuum limit. Since the origin of the large error is traced to that of the mixing coefficients, we develop an alternative method, which we refer to as WI_{VS}, in which the denominator of the ratio for extracting B_K is estimated with the vacuum saturation of \hat{O}_{VV+AA} constructed by the WI method. In this case the fluctuations in the numerator are largely canceled by those in the

denominator, and the resulting error in B_K is substantially reduced as apparent in Fig. 4. The cost is that the correct chiral behavior of the denominator is not respected at a finite lattice spacing due to the contributions of the pseudoscalar matrix element besides the axial vector one. While WI and $W1_{VS}$ methods give different results at a finite lattice spacing, the discrepancy is expected to vanish in the continuum limit. A linear extrapolation in a of the WI_{VS} results yields $B_K(NDR, 2GeV) = 0.562(64)$, which we take as the best value in the present work. This value is consistent with a recent JLQCD result with the Kogut-Susskind action, $B_K(NDR, 2\text{GeV}) = 0.587(7)(17)[2]$ $B_K(NDR, 2\text{GeV}) = 0.587(7)(17)[2]$ $B_K(NDR, 2\text{GeV}) = 0.587(7)(17)[2]$.

Intriguing in Fig. 4 is that the perturbative calculation (PT), which gives completely "wrong value" at $a \neq 0$, also yields the correct result for B_K , when extrapolated to the continuum limit $a = 0$. This is a long extrapolation from negative to positive, but the linearly extrapolated value $B_K(NDR, 2GeV) = 0.639(76)$ is consistent with those obtained with the WI or W_{VS} method. We note that a long extrapolation may bring a large error in the extrapolated value.

Finally we mention possible sources of systematic errors in our results from quenching effects and uncertainties for Gribov copies in the Landau gauge. With the Kogut-Susskind quark action it has been observed that the error due to quenched approximation is small $[1, 10]$ $[1, 10]$. Whether this is supported by calculations with Wilson action we must defer to future studies. For the Gribov problem we only quote an earlier study [[11\]](#page-8-0) which suggests that ambiguities in the choice of the Gribov copies induce only small uncertainties comparable to typical statistical errors in current numerical simulations.

In conclusion our analysis for B_K demonstrate the effectiveness of the method using the chiral Ward identities for constructing the $\Delta s = 2$ operator with the correct chiral property. We have shown that both Wilson and Kogut-Susskind actions give virtually the identical answer for B_K in their continuum limit. We may hope that further improvement of our simulations leads to a precise determination of B_K with the Wilson quark action. The application of this method to B_B is also straightforward.

This work is supported by the Supercomputer Project (No.1) of High Energy Accelerator Research Organization(KEK), and also in part by the Grants-in-Aid of the Ministry of Education (Nos. 08640349, 08640350, 08640404, 08740189, 08740221).

References

- [1] G. W. Kilcup et al., Phys. Rev. Lett. 64 (1990) 25; N. Ishizuka et al., Phys. Rev. Lett. 71 (1993) 24; S. R. Sharpe, Nucl. Phys. B (Proc. Suppl.) 34 (1994) 403.
- [2] JLQCD Collaboration, S. Aoki et al., Nucl. Phys. B (Proc. Suppl.) 53 (1997) 341.
- [3] See, e.g., C. Bernard and A. Soni, Nucl. Phys. **B** (Proc. Suppl.) **9** (1989) 155.
- [4] M. B. Gavela et al., Nucl. Phys. B306 (1988) 677; C. Bernard and A. Soni, Nucl. Phys. B (Proc. Suppl.) 42 (1995) 391; R. Gupta, T. Bhattacharya and S. Sharpe, Phys. Rev. D55 (1997) 4036.
- [5] A. Donini et al., Phys. Lett. **B360** (1995) 83; M. Crisafulli et al., Phys. Lett. B369 (1996) 325; A. Donini et al., Nucl. Phys. B (Proc. Suppl.) 53 (1997) 883.
- [6] G. Martinelli *et al.*, Nucl. Phys. **B445** (1995) 81.
- [7] M. Bochicchio *et al.*, Nucl. Phys. **B262** (1985) 331.
- [8] L. Maiani and G. Martinelli, Phys. Lett. B178 (1986) 265.
- [9] G. Martinelli, Phys. Lett. B141 (1984) 395; C. Bernard, T. Draper and A. Soni, Phys. Rev. D36 (1987) 3224.
- [10] G. Kilcup, D. Pekurovsky and L. Venkataraman, Nucl. Phys. B (Proc. Suppl.) 53 (1997) 345.
- [11] M. L. Paciello, S. Petrarca, B. Taglienti and A. Vladikas, Phys. Lett. B341 (1994) 187.

Tables

	5.9	6.1	6.3	6.5
$L^3 \times T$	$\overline{24^3} \times 64$	$\overline{32^3} \times 64$	$\overline{40^3} \times 96$	$\overline{48^3} \times 96$
$\#\text{conf.}$	300	100	50	24
К	0.15862	0.15428	0.15131	0.14925
	0.15785	0.15381	0.15098	0.14901
	0.15708	0.15333	0.15066	0.14877
	0.15632	0.15287	0.15034	0.14853
K_c	0.15986(3)	0.15502(2)	0.15182(2)	0.14946(3)
a^{-1} (GeV)	1.95(5)	2.65(11)	3.41(20)	4.30(29)
$\alpha_{\overline{MS}}(1/a)$	0.1922	0.1739	0.1596	0.1480
$m_s a/2$	0.0294(14)	0.0198(16)	0.0144(17)	0.0107(16)
$p^{*2}a^2$	0.9595	0.5012	0.2988	0.2056

Table 1: Parameters of simulations.

Table 2: $\langle \bar{K}^0 | \hat{O}_{VV+AA} | K^0 \rangle / |\langle 0 | \hat{P} | K^0 \rangle|^2$ in the chiral limit and $B_K(NDR, 2 \text{GeV})$ for WI, WI[VS] and PT methods as a function of β .

	$\frac{\langle \bar K^0 \hat O_{VV+AA} K^0\rangle}{ \langle 0 \hat P K^0\rangle ^2}$ at $m_q=0$		$B_K(NDR, 2GeV)$		
	WІ	PT.	WI.	WI[VS]	PT
5.9	$-0.0200(39)$	$-0.0415(8)$	$+0.38(6)$	$+0.168(20)$	$-0.468(14)$
6.1	$-0.0068(55)$	$-0.0333(10)$	$+0.68(11)$	$+0.288(29)$	$-0.225(22)$
6.3	$-0.0017(74)$	$-0.0240(12)$	$+0.69(12)$	$+0.342(33)$	$-0.000(21)$
6.5	$+0.006(10)$	$-0.0188(17)$	$+0.72(18)$	$+0.360(52)$	$+0.156(40)$
$a=0$		$-0.0009(31)$		$+0.562(64)$	$+0.639(76)$

Figure Captions

- Fig. 1 Mixing coefficients z_1, \dots, z_4 plotted as a function of external momentum squared $(pa)^2$ for $K = 0.15034$ at $\beta = 6.3$. Vertical line corresponds to $p^* \approx 2 \text{ GeV}.$
- Fig. 2 Comparison of the mixing coefficients z_1, \dots, z_4 evaluated at $p^* \approx 2 \text{ GeV}$ using the Ward identity (WI; solid symbols) and perturbative (PT; open symbols) methods. The coefficients are plotted as a function of m_oa .
- Fig. 3 Test of the chiral behavior of $\langle \bar{K}^0|\hat{O}_{VV+AA}|K^0\rangle/|\langle 0|\hat{P}|K^0\rangle|^2$ at $m_q=0$ for the WI and PT methods. The operators are renormalized at 2 GeV in the NDR scheme. For both methods we use the same \hat{P} perturbatively corrected with the tadpole improvement. The solid line is a linear extrapolation to the continuum limit.
- Fig. 4 $K^0 \bar{K}^0$ matrix element $B_K(\text{NDR}, 2\text{GeV})$ plotted as a function of the lattice spacing for the WI, WI_{VS} and PT methods. The solid lines show linear extrapolations to the continuum limit.

