

Perturbative Renormalization of Improved Lattice Operators*

S. Capitani^a, M. Göckeler^b, R. Horsley^c, H. Perlt^d, P. Rakow^e, G. Schierholz^{a,e} and A. Schiller^d

^aDESY – Theory Group, Notkestrasse 85, D-22607 Hamburg, Germany

^bUniversität Regensburg, Institut für Theoretische Physik, D-93040 Regensburg, Germany

^cHumboldt-Universität, Institut für Physik, Invalidenstrasse 110, D-10115 Berlin, Germany

^dUniversität Leipzig, Institut für Theoretische Physik, Augustusplatz 10/11, D-04109 Leipzig, Germany

^eDESY-IfH Zeuthen, Platanenallee 6, D-15738 Zeuthen, Germany

We derive bases of improved operators for all bilinear quark currents up to spin two (including the operators measuring the first moment of DIS Structure Functions), and compute their one-loop renormalization constants for arbitrary coefficients of the improvement terms. We have thus control over $O(a)$ corrections, and for a suitable choice of improvement coefficients we are only left with errors of $O(a^2)$.

1. INTRODUCTION

In this talk we extend previous calculations of the renormalization constants of quark bilinear operators [1,2] using improved (Sheikholeslami-Wohlert [3]) fermions. Though the calculations are performed in one-loop perturbation theory, we quote results for arbitrary coefficients of the improved action and operators. If they are properly determined (for instance by imposing Ward identities [4]), then only $O(a^2)$ errors are left.

2. QUARK PROPAGATOR

Let us first consider the quark propagator

$$S^{-1} = i \not{p} + m + arp^2/2 - \Sigma^{\text{latt}},$$

where the bare mass is given by $ma = \frac{1}{2\kappa} - 4 - \frac{g^2}{16\pi^2} C_F \Sigma_0$. We write the self-energy as

$$\begin{aligned} \Sigma^{\text{latt}} &= \frac{g^2}{16\pi^2} C_F (i \not{p} \Sigma_1^{\text{latt}} + m \Sigma_2^{\text{latt}} + arp^2 \Sigma_3^{\text{latt}} \\ &+ armi \not{p} \Sigma_4^{\text{latt}} + arm^2 \Sigma_5^{\text{latt}} + O(a^2)). \end{aligned}$$

In the covariant gauge we obtain (for $r = 1$)²:

*Talk presented by S. Capitani at Lattice 97, Edinburgh.
² We denote the coefficient of the improvement term in the action by c_{sw} . We use the abbreviations $\eta = 1 - \alpha$ (where α is the usual gauge parameter) and $L(ap) = \log(a^2 p^2) + \gamma_E - F_0$. The quantities $\gamma_1 = 2$ and $\gamma_2 = 8$ are anomalous dimensions, and $F_0 = 4.369225$.

$$\begin{aligned} \Sigma_0^{\text{latt}} &= -51.4347 + 13.7331c_{sw} + 5.7151c_{sw}^2 \\ \Sigma_1^{\text{latt}} &= 15.6444 - 2.2489c_{sw} - 1.3973c_{sw}^2 \\ &\quad + \gamma_1(1 - \eta) \frac{1}{2} L(ap) + \eta \\ \Sigma_2^{\text{latt}} &= 9.0680 - 9.9868c_{sw} - 0.0169c_{sw}^2 \\ &\quad + (\gamma_2 - \gamma_1) \frac{1}{2} L(ap) + 2\eta \\ \Sigma_3^{\text{latt}} &= 7.0670 + 0.4857c_{sw} - 0.0817c_{sw}^2 \\ &\quad + 0.0719\eta + (-1 + 3c_{sw} - 2\eta) \frac{1}{2} L(ap) \\ \Sigma_4^{\text{latt}} &= -6.2029 - 1.4850c_{sw} + 1.2860c_{sw}^2 \\ &\quad - 0.1437\eta + (-5 - 3c_{sw} + 2\eta) \frac{1}{2} L(ap) \\ \Sigma_5^{\text{latt}} &= -13.4623 + 16.9857c_{sw} - 1.5234c_{sw}^2 \\ &\quad - 2.0719\eta + (-10 + 6c_{sw} + \eta) \frac{1}{2} L(ap). \end{aligned}$$

From this we derive the renormalization constant

$$\begin{aligned} Z_m &= 1 - \frac{g^2}{16\pi^2} C_F (6 \log(a\mu) \\ &\quad - 12.952 - 7.738c_{sw} + 1.380c_{sw}^2), \end{aligned}$$

and the critical κ

$$\kappa_c(g) = \frac{1}{2} \left(4 - \frac{g^2}{16\pi^2} C_F \Sigma_0 \right)^{-1}. \quad (1)$$

In Eq. (1) one has the choice of using $c_{sw} = 1$ or the actual value used in true simulations. In Fig. 1 we compare the various choices with the data, and we find that the latter choice, combined with tadpole improvement, agrees best. This justifies our procedure.




Figure 1. The dotted and dashed curves correspond to $c_{sw} = 1$, without and with tadpole improvement. The solid curve corresponds to c_{sw} as given by the Alpha Collaboration [5], plus tadpole improvement. This is compared with recent data.

The free propagator at $\kappa_c = \frac{1}{8r}$ is given by $S(p) = \frac{1}{i\not{p}} + \frac{ar}{2} + O(a^2)$. The $O(a)$ term is constant in momentum space, and there are deviations from the continuum propagator only at very short distances. We thus suggest our Ansatz for the interacting propagator as:

$$S(p) = \frac{Z_2}{1 + b_2 arm} \frac{1}{i\not{p} + m_R} + ar\lambda.$$

Imposing this structure for the propagator, we find for the values of the parameters (for $r = 1$):

$$\lambda = \frac{1}{2} \left(1 + \frac{g^2}{16\pi^2} C_F (2\Sigma_1 - 2\Sigma_3) \right)$$

$$Z_2 = \left(1 + \frac{g^2}{16\pi^2} C_F \Sigma_1 \right)$$

$$\begin{aligned} b_2 &= 1 + \frac{g^2}{16\pi^2} C_F (2\Sigma_1 - \Sigma_2 - 2\Sigma_3 - \Sigma_4) \\ m_R &= m Z_m (1 - b_m arm) \\ &= m \left(1 + \frac{g^2}{16\pi^2} C_F (\Sigma_1 - \Sigma_2) \right) \left(1 - \frac{arm}{2} - \frac{arm}{2} \frac{g^2}{16\pi^2} C_F (2\Sigma_1 - \Sigma_2 - 2\Sigma_3 - 2\Sigma_4 + 2\Sigma_5) \right). \end{aligned}$$

We see that λ can be a constant only if $c_{sw} = 1$:

$$\begin{aligned} 2\Sigma_1 - 2\Sigma_3 &= 17.1548 - 5.4691c_{sw} - 2.6311c_{sw}^2 \\ &\quad + 1.8562\eta + 6(1 - c_{sw}) \frac{1}{2} L(ap); \end{aligned}$$

for every other value of c_{sw} there are $a \log(ap)$ contributions to $S(p)$. The same happens for the parameters b_2 and b_m : both of them involve $a \log(ap)$ terms, that at $c_{sw} = 1$ cancel out.

Two possible expressions that remove $O(a)$ effects from the quark propagator are:

$$S_{\text{imp}}(p) \equiv (1 + b_2 arm)(S(p) - ar\lambda); \quad (2)$$

$$S^{-1}(p) = (1 + b_2 arm) S_{\text{imp}}^{-1}(p) \quad (3)$$

$$-ar\lambda S_{\text{imp}}^{-1}(p) S_{\text{imp}}^{-1}(p).$$

The improved propagators in these two definitions differ by terms of $O(a^2)$, but both are free of $O(a)$ effects. Eq. (3) is a non-linear equation which has to be solved iteratively, but it seems a better definition in practice than the first one. This is because Eq. (2) seems to have larger $O(a^2)$ effects, thus using Eq. (3) we can reach higher momenta.

3. BASES FOR IMPROVED OPERATORS

Some fundamental bases necessary to achieve full $O(a)$ improvement for point operators are:

$$(\bar{\psi}\psi)^{\text{imp}} = (1 + a b m) \bar{\psi}\psi - \frac{1}{2} ac_1 \bar{\psi} \overleftrightarrow{D} \psi$$

$$(\bar{\psi}\gamma_5\psi)^{\text{imp}} = (1 + a b m) \bar{\psi}\gamma_5\psi + \frac{1}{2} ac_1 \partial_\mu (\bar{\psi}\gamma_\mu\gamma_5\psi)$$

$$\begin{aligned} (\bar{\psi}\gamma_\mu\psi)^{\text{imp}} &= (1 + a b m) \bar{\psi}\gamma_\mu\psi - \frac{1}{2} ac_1 \bar{\psi} \overleftrightarrow{D}_\mu \psi \\ &\quad + i \frac{1}{2} ac_2 \partial_\lambda (\bar{\psi}\sigma_{\mu\lambda}\psi) \end{aligned}$$

$$(\bar{\psi}\gamma_\mu\gamma_5\psi)^{\text{imp}} = (1 + a b m)\bar{\psi}\gamma_\mu\gamma_5\psi - i\frac{1}{2}ac_1\bar{\psi}\sigma_{\mu\lambda}\gamma_5\overleftrightarrow{D}_\lambda\psi + \frac{1}{2}ac_2\partial_\mu(\bar{\psi}\gamma_5\psi),$$

where $\overleftrightarrow{D}_\mu = \overrightarrow{D}_\mu - \overleftarrow{D}_\mu$. For the tensor operator see Ref. [6]. The normalizations are chosen such that $c_i = 1 + O(g^2)$ (in this way, for $g = 0$ we realize tree-level improvement).

Some possible bases for the improvement of the one-link DIS operators $\mathcal{O}_{\mu\nu} = \bar{\psi}\gamma_\mu\overleftrightarrow{D}_\nu\psi$ and the polarized $\mathcal{O}_{\mu\nu}^5 = \bar{\psi}\gamma_\mu\gamma_5\overleftrightarrow{D}_\nu\psi$ are:

$$\begin{aligned} \mathcal{O}_{\mu\nu}^{\text{imp}} &= (1 + a b m)\bar{\psi}\gamma_\mu\overleftrightarrow{D}_\nu\psi \\ &- a c_1 g \bar{\psi}\sigma_{\mu\lambda}F_{\nu\lambda}^{\text{clover}}\psi - \frac{1}{4} a c_2 \bar{\psi}\{\overleftrightarrow{D}_\mu, \overleftrightarrow{D}_\nu\}\psi \\ &+ \frac{1}{2} a i c_3 \partial_\lambda(\bar{\psi}\sigma_{\mu\lambda}\overleftrightarrow{D}_\nu\psi); \end{aligned} \quad (4)$$

$$\begin{aligned} \mathcal{O}_{\mu\nu}^{5,\text{imp}} &= (1 + a b m)\bar{\psi}\gamma_\mu\gamma_5\overleftrightarrow{D}_\nu\psi \\ &- a i c_1 g \bar{\psi}\gamma_5 F_{\mu\nu}^{\text{clover}}\psi \\ &- \frac{1}{4} a i c_2 \bar{\psi}\sigma_{\mu\lambda}\gamma_5\{\overleftrightarrow{D}_\lambda, \overleftrightarrow{D}_\nu\}\psi \\ &+ \frac{1}{2} a i c_3 \partial_\mu(\bar{\psi}\gamma_5\overleftrightarrow{D}_\nu\psi). \end{aligned} \quad (5)$$

The relation $[\overleftrightarrow{D}_\mu, \overleftrightarrow{D}_\nu]^{\text{latt}} = 4 i g F_{\mu\nu}^{\text{clover}} + O(a^2)$ is useful to derive other bases.

If all these operators are inserted into forward matrix elements the surface terms $\partial_\mu(\bar{\psi}\mathcal{O}\psi)$ vanish due to momentum conservation. Using the equations of motion it is also possible to further reduce the number of improvement coefficients.

4. POINT OPERATORS

We present the general Z renormalization factors with the improvement coefficients c_{sw} (for the action) and c_i (for the operators) kept general. These factors will be essential when a determination of the improvement coefficients is done [4].

Writing (to order g^2) $\langle\mathcal{O}\rangle|_{g^2} = \mathcal{O}^{(1)} + a\mathcal{O}^{(2)}$, the calculation of the amputated matrix elements for $A_5 = \bar{\psi}\gamma_5\psi$ and $A_\mu = \bar{\psi}\gamma_\mu\gamma_5\psi$ gives

$$A_5^{(1)} = \frac{g^2}{16\pi^2}C_F\gamma_5(0.5750 + 3.4333c_{sw}^2$$

$$-2\eta + (-4 + \eta)L(ap))$$

$$A_5^{(2)} = 0$$

$$\begin{aligned} A_\mu^{(1)} &= \frac{g^2}{16\pi^2}C_F(\gamma_\mu\gamma_5(0.1519 - \eta - 19.3723c_1 \\ &+ 2.4967c_{sw} + 10.3167c_1c_{sw} - 0.8541c_{sw}^2 \\ &- 0.8846c_1c_{sw}^2 - (1 - \eta)L(ap)) - 2(1 - \eta)\frac{\not{p}\gamma_5\not{p}_\mu}{p^2} \end{aligned}$$

$$\begin{aligned} A_\mu^{(2)} &= \frac{ig^2}{16\pi^2}C_F\left(\frac{1}{2}(\not{p}\gamma_\mu\gamma_5 + \gamma_\mu\gamma_5\not{p})\right) \cdot (1.5323 \\ &+ 0.7322c_1 - 1.7181c_{sw} + 0.5430c_1c_{sw} \\ &+ 0.1302c_{sw}^2 + 0.0537c_1c_{sw}^2 - \eta(0.8563 \\ &- 4.0583c_1) + (1 - \eta)(1 + c_1)L(ap)). \end{aligned}$$

Similar results hold for the other bilinears [6]. In general the $O(a)$ terms do not contribute to the tree-level structure in the massless case.

To extract the Z factors we project forward matrix elements onto their tree-level structure: $\langle q(p)|\mathcal{O}(\mu)|q(p)\rangle = Z_\mathcal{O}Z_\psi^{-1}\langle q(p)|\mathcal{O}(a)|q(p)\rangle$, and $\langle q(p)|\mathcal{O}(\mu)|q(p)\rangle|_{p^2=\mu^2} = \langle q(p)|\mathcal{O}(a)|q(p)\rangle|_{p^2=\mu^2}^{\text{tree}}$, where Z_ψ is the wave function renormalization factor. The one-loop renormalization factor $Z_\mathcal{O}$ for a lattice operator \mathcal{O} can be cast into the form

$$Z_\mathcal{O}(a\mu, g) = 1 - \frac{g^2}{16\pi^2}C_F(\gamma_\mathcal{O}\log(a\mu) + B_\mathcal{O}),$$

where $\gamma_\mathcal{O}$ is its anomalous dimension, and $B_\mathcal{O}$ the finite part of $Z_\mathcal{O}$.

The Wilson coefficients are usually computed in the $\overline{\text{MS}}$ scheme; to convert the finite parts to this scheme one can use the relations $B_\mathcal{O}^{MS} = B_\mathcal{O} - B_\mathcal{O}^{\text{cont}}$ and $B_\mathcal{O}^{\overline{\text{MS}}} = B_\mathcal{O}^{MS} + \frac{\gamma_\mathcal{O}}{2}(\gamma_E - \log(4\pi))$, where $\gamma_\mathcal{O}$ and $B_\mathcal{O}^{\text{cont}}$, the finite contributions to the continuum renormalization factors $Z_\mathcal{O}^{\text{cont}}$, are:

| \mathcal{O} | $\gamma_\mathcal{O}$ | $B_\mathcal{O}^{\text{cont}}$ |
|----------------------------------|----------------------|---|
| 1, γ_5 | -6 | $5 + \frac{\gamma_\mathcal{O}}{2}(\gamma_E - \log(4\pi)) - \eta$ |
| $\gamma_\mu, \gamma_\mu\gamma_5$ | 0 | 0 |
| $\sigma_{\mu\nu}\gamma_5$ | 2 | $-1 + \frac{\gamma_\mathcal{O}}{2}(\gamma_E - \log(4\pi)) + \eta$ |

The results in the $\overline{\text{MS}}$ scheme for all point operators are then as follows:

$$\begin{aligned}
B_1^{\overline{MS}} &= 12.9524 - 19.1718 c_1 + 7.7379 c_{sw} \\
&\quad + 13.8007 c_1 c_{sw} - 1.3804 c_{sw}^2 - 3.5383 c_1 c_{sw}^2 \\
B_{\gamma_5}^{\overline{MS}} &= 22.5954 - 2.2487 c_{sw} + 2.0360 c_{sw}^2 \\
B_{\gamma_\mu}^{\overline{MS}} &= 20.6178 - 9.7864 c_1 - 4.7456 c_{sw} \\
&\quad + 3.4164 c_1 c_{sw} - 0.5432 c_{sw}^2 + 0.8846 c_1 c_{sw}^2 \\
B_{\gamma_\mu \gamma_5}^{\overline{MS}} &= 15.7963 - 19.3723 c_1 + 0.2478 c_{sw} \\
&\quad + 10.3167 c_1 c_{sw} - 2.2514 c_{sw}^2 - 0.8846 c_1 c_{sw}^2 \\
B_{\sigma_{\mu\nu} \gamma_5}^{\overline{MS}} &= 17.0181 - 16.2438 c_1 - 3.9133 c_{sw} \\
&\quad + 6.8553 c_1 c_{sw} - 1.9723 c_{sw}^2 + 0.5897 c_1 c_{sw}^2.
\end{aligned}$$

With the appropriate values of c_{sw} and c_i , all these operators are then fully $O(a)$ improved.

5. ONE-LINK OPERATORS

The many-link operators are essential in the OPE expansion of current-current correlators occurring in Structure Functions computations. Using Eq. (4) with $b = 0$ and $c_3 = 0$ for the unpolarized case, in one-loop perturbation theory for the representation $\tau_1^{(3)}$ ($\mathcal{O}_{\{44\}} - 1/3(\mathcal{O}_{\{11\}} + \mathcal{O}_{\{22\}} + \mathcal{O}_{\{33\}})$) we have:

$$\begin{aligned}
\mathcal{O}_{\mu\nu, \tau_1^{(3)}}^{\text{imp}} : B_{\mathcal{O}} &= -1.8826 - 3.9698 c_{sw} \\
&\quad - 1.0398 c_{sw}^2 + c_1 (5.9970 - 3.2685 c_{sw}) \\
&\quad - c_2 (6.6733 - 4.5371 c_{sw} - 0.4462 c_{sw}^2).
\end{aligned}$$

For the other representation, $\tau_3^{(6)}$ ($\mathcal{O}_{\{14\}}$), we find full agreement with the $O(a)$ improved result given in Ref. [1] for $c_{sw} = 1$ and $c_1 = c_2 = 1$.

For the polarized case, considering Eq. (5), the result for the representation $\tau_4^{(6)}$ ($\mathcal{O}_{\{14\}}^5$) is

$$\begin{aligned}
\mathcal{O}_{\mu\nu, \tau_4^{(6)}}^{5, \text{imp}} : B_{\mathcal{O}} &= -4.0988 - 1.3593 c_{sw} - 1.8926 c_{sw}^2 \\
&\quad - c_2 (27.5719 - 16.1193 c_{sw} + 0.7570 c_{sw}^2).
\end{aligned}$$

REFERENCES

1. S. Capitani and G. Rossi, Nucl. Phys. B433 (1995) 351.
2. M. Göckeler et al., Nucl. Phys. B472 (1996) 309; M. Göckeler et al., Nucl. Phys. B (Proc. Suppl.) 53 (1997) 896.
3. B. Sheikholeslami and R. Wohlert, Nucl. Phys. B259 (1985) 572.
4. P. Rakow, these proceedings; R. Horsley, these proceedings.
5. K. Jansen et al., Phys. Lett. B372 (1996) 275; M. Lüscher et al., Nucl. Phys. B491 (1997) 323.
6. S. Capitani et al., in preparation.