

Heavy Hybrids From NRQCD

Presented by S. Collins, UKQCD Collaboration^a *

^aDept. of Physics and Astronomy, Glasgow University, Glasgow, G12 8QQ, Scotland

We present results for the exotic 0^{+-} and 1^{-+} $b\bar{b}g$ hybrids at $\beta = 6.0$. A leading order NRQCD action is used to generate the heavy quark propagators. The optimisation of operators is discussed and we compare our results to the predictions of recent potential calculations.

1. INTRODUCTION

In hybrid mesons the gluon string between the $q\bar{q}$ pair is excited and contributes to the quantum numbers of the particle. These mesons are predictions of QCD which are not yet confirmed in experiment. More difficult to discriminate from quark model states than glueballs, it is particularly important to provide accurate predictions from lattice studies.

Most experimental candidates are for hybrids containing light quarks; however, the best system for a hybrid search may be $c\bar{c}$ or $b\bar{b}$, where there is a large gap between the lowest state and the $D\bar{D}$ and $B\bar{B}$ thresholds respectively. Theoretically, heavy hybrids are easier to study; on the lattice, NRQCD can be used to simulate heavy quarks and the results can be compared with those from hybrid potentials.

Here, we present results for $b\bar{b}g$ hybrids. The lattice hybrid operators are constructed from staples of gauge links, $\square-\square$, in combinations corresponding to the T_1^{+-} and T_1^{-+} representations of the cubic symmetry group, as described in Lacock et al. [1]. In early potential studies [2] it was found that this gluon excitation, corresponding to one unit of angular momentum about the $q\bar{q}$ axis, gives rise to the lowest hybrid meson states. Combined with $q\Gamma\bar{q}$ in a spin singlet (A_1^{-+}) and triplet (T_1^{-+}) the hybrid meson states

$$\begin{aligned} T_1^{+-} &: \quad 1^{--}, O^{-+}, \mathbf{1}^{-+}, 2^{-+} \\ T_1^{-+} &: \quad 1^{++}, \mathbf{O}^{+-}, 1^{+-}, \mathbf{2}^{+-} \end{aligned}$$

*In collaboration with C. Davies, Glasgow Univ.. Results provided by G. Bali, Southampton Univ. are also presented.

are accessible, where the exotic states are shown in bold. In the static limit these states are all degenerate; introduction of quark motion, using the lowest order NRQCD action, will create a splitting between the sets of states arising from the T_1^{+-} and T_1^{-+} gauge paths. Spin-spin and spin-orbit interactions are needed in order to obtain splittings between the individual states. These spin splittings, eg between the O^{-+} , 1^{-+} and 2^{-+} , will be ordered in the same way as for the non-hybrid 3P states and are likely to be of order 10 MeV. Since the statistical errors in our calculation are still at approximately 200 MeV we ignore the fine structure and use the lowest order NRQCD action including only the kinetic energy term. In this case there is no mixing between the hybrid and normal $b\bar{b}$ states.

The correlators corresponding to the hybrid operators were computed on 500 quenched $16^3 \times 48$ configurations at $\beta = 6.0$. A bare b quark mass, from $b\bar{b}$ spectroscopy, of $1.71a^{-1}$ was used in the NRQCD action. The hybrid signal is dominated by noise after only a few timeslices, and multiple spatial and temporal sources are required. It is essential to optimise the operators in terms of the $q\bar{q}$ separation, R , depth of the staple, D , and the coefficient, c , and level, n , of the fuzzing of the gauge links. We use the results of a study of the hybrid potential [3] on the same configurations to choose the optimal values of these parameters.

2. THE HYBRID POTENTIAL

The hybrid potential corresponding to the T_1 representation, E_u , was calculated in the standard way using operators of the form $\square-\square$. In

Table 1

$R \times D$	6×1	6×2	6×3	6×6
$A_{g.s.}$	0.90(2)	0.92(2)	0.92(2)	0.39(8)




Figure 1.

order to find the optimal operator, the depth D was varied at each R and the maximal overlap of the operator with the ground state, $A_{g.s.}$, was sought. Table 1 presents the results as a function of D for the example of $R = 6$. $A_{g.s.}$ is not very sensitive to D as long as D is small. A fairly uniform optimal overlap was found for a wide range of R , where we investigated up to $R = 9$.

Fuzzing the gauge links seemed to act as a smoothing function on the hybrid source and it was found sufficient to optimise the value of the spatial plaquette with respect to c and n in order to obtain their optimal values: roughly $c = 2.3$ and $n = 25$.

Using these optimal parameters for the gauge path, the hybrid potential shown in figure 1 was obtained; the ground state potential, A_{1g} , is also shown for comparison. The most striking feature of the potential is its flatness which suggests broad hybrid wavefunctions and densely packed states. To extract this information the potential was fitted to a functional form suggested by string models:

$$V = V_{g.s.} + (1 - e^{-cR}) \pi/R, \quad (1)$$

with parameter c . The hybrid energy levels, E_{nl} , and wavefunctions, u_{nl} , where $\psi = u_{nl} Y_m^{nl}/R$,

Table 2

n	1	2	3
$l = 1$	10.87	11.13	11.39
$l = 2$	11.02	11.29	11.55




Figure 2.

can then be extracted by solving the Schrödinger equation in the continuum. The lowest lying hybrid levels are given in table 2 in GeV and the corresponding radial wavefunctions are shown in figure 2. The scale was set using the average of the lattice spacings from the $b\bar{b}$ $P-S$ and $2S-1S$ splittings. Note that the first hybrid state is a P -state. The $1P$ $b\bar{b}$ wavefunction is shown for comparison in the figure.

The wavefunctions are broad, as expected. For a $L \sim 1.8$ fm lattice we can expect large finite volume errors for a ground state hybrid meson with $r_{r.m.s.} \sim 0.28$ fm. Figure 2 suggests a lattice hybrid operator with a $q\bar{q}$ separation of 0.5 fm or ~ 6 lattice units at $\beta = 6.0$ would minimise the overlap with the first excited state since the corresponding wavefunction has a node at this point. However, a more reliable method to extract the ground state energy is to perform multiple exponential fits to a matrix of correlation functions obtained from different $q\bar{q}$ separations, where the different values of R give rise to very different overlaps with the hybrid excited states in the off-diagonal elements of the matrix. For example, from figure 2 we see a suitable 2×2 matrix can




Figure 3.

be formed from $R = 3$ and $R = 7$.

3. HYBRIDS WITH PROPAGATING QUARKS

Considering the potential results we performed an initial calculation of the T_1^{+-} and T_1^{-+} hybrids using a staple size of $R = 3$ and $D = 2$ at both the source and the sink. Figure 3 presents the results for the effective mass of the T_1^{+-} hybrid obtained from 10,000 sources. There is a large contribution from excited states for this operator. However, a plateau seems to begin from timeslice 6 or 7. One-exponential fits in the region 6–10 are in agreement with two-exponential fits with $t_{min} = 3-5$. Considering all these fits, we find the ground state simulation energy to be consistent with 1.0(1), as indicated in the figure. This corresponds to a splitting with the 3S_1 $b\bar{b}$ state of 0.55(10) in lattice units. We also find a splitting of 0.1 – 0.3 with the T_1^{-+} states, which confirms that the 1^{-+} is the lowest exotic hybrid.

Converting our results into physical units we obtain 11.5(5) GeV for the T_1^{+-} hybrid state. This is compared with the lower lying $b\bar{b}$ spectrum obtained from an $O(Mv^4)$ NRQCD action [4] and also from a potential study, accurate to the same order in v [5] in figure 4. Our prediction lies 1σ above the threshold for $B\bar{B}$ production but 1σ below that for BB^{**} , which is predicted by flux tube models to be the dominant hybrid decay mode. A study by Manke et al. [6] using dif-




Figure 4.

ferent operators on the same configurations finds consistent results with the T_1^{+-} hybrid lying 1σ higher. Our results are also consistent with the predictions from the potential.

The position of the hybrid state with respect to the two thresholds is not settled. An improved determination of the hybrid state is needed and we propose to do this by computing a matrix of correlators between different $q\bar{q}$ separations. At the moment the large statistical errors dominate the uncertainty in the prediction. However, the central value may be shifted significantly by finite volume effects and to a lesser extent quenching and the systematic errors associated with NRQCD.

ACKNOWLEDGEMENTS

The computations were performed on the J90 at EPCC in Edinburgh as part of HPCI under EPSERC grant no. GR/K55745.

REFERENCES

1. P. Lacock et al., Phys. Rev. D55 (1997) 1548.
2. C. Michael et al., Nucl. Phys. B347 (1990) 854.
3. G. Bali, in preparation.
4. C. Davies et al., Phys. Rev. D50 (1993) 6963.
5. G. Bali et al., Phys. Rev. D56 (1997) 2566.
6. T. Manke, this publication.