The Light Quark Masses with the Wilson Quark Action using Chiral Ward Identities^{*}

JLQCD Collaboration

S. Aoki^a, M. Fukugita^b, S. Hashimoto^c, N. Ishizuka^{a,d}, Y. Iwasaki^{a,d}, K. Kanaya^{a,d} Y. Kuramashi^e, M. Okawa^e, A. Ukawa^a, T. Yoshié^{a,d}

^aInstitute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

^bInstitute for Cosmic Ray Research, University of Tokyo, Tanashi, Tokyo 188, Japan

^cComputing Research Center, High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan

^dCenter for Computational Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

^eInstitute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan

We present results for the light quark masses for the Wilson quark action obtained with the PCAC relation for the one-link extended axial vector current in quenched QCD at $\beta = 5.9 - 6.5$. This method leads to a remarkable improvement of scaling behavior of the light quark masses compared to the conventional method. We obtain $\overline{m}_l = 3.87(37)$ MeV for the averaged up and down quark mass and $\overline{m}_s = 97(9)$ MeV for the strange quark mass in the \overline{MS} scheme at $\mu = 2$ GeV.

1. Introduction

The chiral symmetry breaking term in the Wilson quark action causes large scaling violation effects of O(a) for physical quantities in numerical simulations of lattice QCD. A manifestation of this effect is that the light quark masses for the Wilson action show a strong a dependence at finite lattice spacings in contrast to a weak dependence for the case with the Kogut-Susskind (KS) action which retains U(1) chiral symmetry[1]. Uncertainties in a long extrapolation to the continuum limit needed for the Wilson case forms a part of the difficulty to settle the question whether the Wilson and KS quark actions give a consistent result in the continuum $\lim_{n \to \infty} \lim_{n \to \infty}$ leads us to reconsider the definition of quark mass for the Wilson action.

It is well known that the current quark mass defined by the PCAC relation for the Wilson action [3,4] differs by O(a) from the conventionally defined quark mass at finite lattice spacing. We examine this point in more detail and find that the former definition applied with the onelink separated axial vector current that naturally arises in chiral Ward identities [3] leads to a significantly improved scaling behavior for the light quark masses.

Our calculations are carried out as a part of our simulation for the kaon *B*-parameter[5] with the Wilson quark action in quenched QCD. The plaquette gauge action is employed, and data are collected at $\beta = 5.9 - 6.5$. Point source quark propagators calculated for four values of the hopping parameter at each β are used for this work. Errors are estimated by the single elimination jackknife procedure for all measured quantities.

2. Calculational method

The conventional (CV) definition of quark mass is given by $m_q^{\text{CV}}a = (1/K - 1/K_c^{\text{CV}})/2$ where

^{*}presented by Y. Kuramashi



Figure 1. m_{π}^2 and ρ as a function of 1/(2K) at $\beta = 6.3$.

the critical hopping parameter $K_c^{\rm CV}$ is determined by extrapolating the pseudo scalar meson mass squared m_{PS}^2 linearly to $m_{PS}^2 = 0$ in the inverse hopping parameter 1/(2K). To extract the averaged up and down quark mass $m_l^{\rm CV} = (m_u^{\rm CV} + m_d^{\rm CV})/2$ and the strange quark mass $m_s^{\rm CV}$ we assume the following quark mass dependence for m_{PS} and the vector meson mass m_V ,

$$m_{PS}^2 = C_{PS}(m_{q1}^{\rm CV} + m_{q2}^{\rm CV})/2$$
 (1)

$$m_V = m_V^0 + C_V (m_{q1}^{\rm CV} + m_{q2}^{\rm CV})/2.$$
 (2)

The lattice spacing a is fixed with m_V^0 using $m_\rho = 770$ MeV. We employ the hadron mass ratio $m_\pi/m_\rho = 0.1783$ to determine the averaged up and down quark mass m_l^{CV} . The strange quark mass m_s^{CV} is estimated in two ways using $m_K/m_\rho = 0.644$ and $m_\phi/m_\rho = 1.323$. We convert m_q^{CV} into $\overline{m}_q^{\text{CV}}$ defined in the $\overline{\text{MS}}$ scheme at the scale $\mu = 1/a$ using the tadpole-improved perturbative mass renormalization factor $Z_m(\mu a = 1)$ evaluated with $\alpha_{\overline{\text{MS}}}(1/a)$.

The Ward identity (WI) method requires a calculation of the ρ parameter defined by the PCAC relation[3];

$$2\rho(K) = \frac{\langle 0|\nabla_{\mu}A_{\mu}^{\text{ext},a}|\pi^{a}(\mathbf{p}=0)\rangle}{\langle 0|P^{a}|\pi^{a}(\mathbf{p}=0)\rangle},$$
(3)

We take the one-link extended axial vector current $A^{\text{ext},a}_{\mu}$ since it is this current



Figure 2. Averaged up and down quark mass in the $\overline{\text{MS}}$ scheme at $\mu = 2 \text{GeV}$ as a function of a.

which naturally arises in the Ward identities. We extract the ρ parameter from the ratio $\langle \nabla_4 A_4^{\text{ext}}(t) P(1) \rangle / \langle P(t) P(1) \rangle / 4 + \langle \nabla_4 A_4^{\text{ext}}(T-t+2) P(1) \rangle / \langle P(T-t+2) P(1) \rangle / 4$, each two-point function projected to the zero spatial momentum, by fitting a plateau as a function of t.

In Fig. 1 a representative result for the ρ parameter is plotted as a function of 1/(2K) together with m_{π}^2 . We observe a clear linear behavior both for ρ and m_{π}^2 . The critical hopping parameter K_c^{WI} extracted from a linear extrapolation for ρ is slightly different from K_c^{CV} for m_{π}^2 (see Table 1 for numerical details). We ascribe the discrepancy to uncertainties in the extrapolations of ρ and m_{π}^2 , because the critical hopping parameter obtained with the two definitions should agree at each β . To avoid this problem, we define the bare quark mass for the WI method $m_q^{\text{WI}}(1/a)$ by

$$m_q^{\rm WI}(1/a) = \frac{\rho(K)}{(1/K - 1/K_c^{\rm WI})/2} m_q^{\rm CV}(1/a).$$
(4)

We convert quark masses defined on the lattice into those in the $\overline{\text{MS}}$ scheme at the scale $\mu = 1/a$ using the tadpole-improved perturbative renormalization factor $Z_{A^{\text{ext}}}(\mu a = 1)/Z_P(\mu a = 1)$ evaluated with $\alpha_{\overline{\text{MS}}}(1/a)$.

We quote final results for $\overline{m}_q^{\text{CV,WI}}$ at the scale $\mu = 2 \text{GeV}$ which are obtained by a two-loop renormalization group running from $\mu = 1/a[6]$.

Critical hopping parameters and the light quark masses in the MS scheme at $\mu = 2 \text{GeV}$.										
	input	$\beta = 5.9$	6.1	6.3	6.5	a = 0				
$K_c^{\rm CV}$		0.15986(3)	0.15502(2)	0.15182(2)	0.14946(3)					
$K_c^{\rm WI}$		0.15976(1)	0.15497(1)	0.15179(1)	0.14948(1)					
$\overline{m}_l^{\rm CV}$ (MeV)	$m_{\pi}/m_{ ho}$	5.54(14)	5.14(22)	4.89(30)	4.62(41)	3.96(49)				
$\overline{m}_l^{\text{WI}}$ (MeV)	$m_{\pi}/m_{ ho}$	3.97(10)	3.94(17)	3.97(23)	3.86(32)	3.87(37)				
$\overline{m}_s^{\rm CV}$ (MeV)	$m_K/m_{ ho}$	139(3)	129(5)	123(8)	116(10)	100(12)				
$\overline{m}_{s}^{\mathrm{WI}}$ (MeV)	$m_K/m_{ ho}$	100(3)	99(4)	100(6)	97(8)	97(9)				
$\overline{m}_s^{\rm CV}$ (MeV)	$m_{\phi}/m_{ ho}$	184(13)	149(17)	151(24)	120(23)					
$\overline{m}_s^{\rm WI} \ ({\rm MeV})$	$m_{\phi}/m_{ ho}$	132(9)	115(13)	122(20)	99(19)					

				-	
	1 + 1 1: 1.	+	: +1 MIC	·	
Unitical nonphing parameters	and the nor	t onark masses	in the MS	scheme at	$\mu = 2 \tau e v$
eritical hopping parameters	, and the light	guarn masses	111 0110 1010	bonionio au	p 2001

3. Results for light quark masses

Table 1

In Table 1 we summarize the results for the averaged up and down quark mass \overline{m}_l . The lattice spacing dependence of \overline{m}_l is shown in Fig. 2. The results with the WI method show a remarkably flat behavior. This allows a reliable extrapolation to the continuum limit by a linear function in a, with which we find $\overline{m}_l^{\text{WI}} = 3.87(37) \text{MeV}$ at a = 0. In contrast a large scaling violation effect is seen for the results with the CV method. While a linear extrapolation yields a consistent result $\overline{m}_{a}^{\text{CV}} = 3.96(49) \text{MeV}$ in the continuum limit, systematic uncertainties due to a long extrapolation are quite large. Finding a theoretical explanation as to why finite a corrections are so small for quark masses with the WI method would be an interesting problem [7].

In Fig. 2 we also plot the KS results [8] for comparison. They show small scaling violations and are systematically smaller than the results for the WI method. However, we cannot conclude at this stage whether the Wilson results are inconsistent with the KS results. The one-loop correction in the mass renormalization factor for the KS quark action is 50% - 100% depending on the lattice spacing for the results in Fig. 2, which suggests that higher order corrections might be large.

Our results for the strange quark mass \overline{m}_s determined from m_K and m_{ϕ} are listed in Table 1. We find that results with the WI method are quite flat also in this case.

Estimates of \overline{m}_s using m_K are expected to satisfy $\overline{m}_s \approx 25\overline{m}_l$ for each β if we assume the functional form (1), since then $m_K^2/m_{\pi}^2 =$ $(\overline{m}_l + \overline{m}_s)/(2\overline{m}_l) \approx 13$. Making linear extrapolations in *a* for the results of the WI and the CV method we find a mutually consistent result in the continuum limit: $\overline{m}_s^{\text{WI}} = 97(9)$ MeV and $\overline{m}_s^{\text{CV}} = 100(12)$ MeV.

For the alternative determination of \overline{m}_s with the use of m_{ϕ} , the discrepancy between the WI and the CV results reduces toward the continuum limit. However, large errors originating from those of the vector meson mass hinder us from reliably extrapolating the results to the continuum limit.

This work is supported by the Supercomputer Project (No. 97-15) of High Energy Accelerator Research Organization (KEK), and also in part by the Grants-in-Aid of the Ministry of Education (Nos. 08640349, 08640350, 08640404, 09246206, 09304029, 09740226).

REFERENCES

- A. Ukawa, Nucl. Phys. B (Proc. Suppl.) 30 (1993) 3.
- R. Gupta and T. Bhattacharya, Phys. Rev. D55 (1997) 7203.
- M. Bochicchio *et al.*, Nucl. Phys. **B262** (1985) 331.
- 4. S. Itoh *et al.*, Nucl. Phys. **B274** (1986) 33.
- JLQCD Collaboration, S. Aoki *et al.*, heplat/9705035.
- C. R. Allton *et al.*, Nucl. Phys. **B431** (1994) 667.
- 7. Y. Kuramashi, in progress.
- JLQCD Collaboration, S. Aoki *et al.*, Nucl. Phys. B (Proc. Suppl.) 53 (1997) 209.