## A New Approach of Fermion Field on Lattice

Bo Feng<sup>a</sup>, \*Jian<br/>ming Li<sup>a,b</sup>, Xingchang Song<sup>a,b</sup>

a Department of Physics, Peking University, Beijing 100871, China

b Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China

## Abstract

A new approach to formulate the fermion field on lattice is introduced by proposing a new Dirac operator on lattice. This approach can eliminate the Fermion doubling problem, preserve the chiral symmetry and get the same dispersion relation for both Fermion and Boson fields. Then the Weinberg-Salam model on lattice may be formulated in this approach.

PACS number: 11.15.Ha

<sup>\*</sup>Email address: lijm@itp.ac.cn

In lattice field theory, it is well-known that the naive discretization of Fermion field suffers from the problem of Fermion doubling, while the scalar field doesn't. The studies of Fermion species doubling problem persists a long duration since Wilson [1] [2]. To cure from this Wilson [1] [2] added a new term, so-called the "Wilson term", to the lattice Fermion Lagrangian. This term kills the superfluous components of the Fermion field but breaks the chiral symmetry and leads to different dispersion relations for Fermions and Bosons. Kogut and Susskind [3] [4] proposed the "stagger model", but it needs four generations of Fermions at least, and deals with the Fermions and Bosons on different footing. Besides these two popular prescriptions, Drell et al [5] developed another approach which preserves the chiral symmetry and at the same time correctly counts the number of Fermion states. Their crucial point was to introduce a lattice gradient operator which couples all lattice sites along a given direction instead of coupling only nearest-neighbor sites. However, no convincing approach which can fit all the physical phenomena exists so far [6], such as we don't know how to reformulate the Weinberg-Salam model on lattice.

In this paper, to overcome those difficulties we propose to adopt a new approach in which a new Dirac operator is introduced. The square of this Dirac operator is the D'Alembertian operator, which recovers the relation in continue field theory. By using this approach, we can eliminate the Fermion doubling problem and at the same time preserve the chiral symmetry. Besides these, we can get the same dispersion relation for both Fermions and Bosons.

First let us take a look upon how the Fermion-doubling problem arises. For free scalar field  $\phi(x)$ , the lattice action in the Euclidean space is (for more detail, see [7])

$$S(\phi) = \sum_{n} \{ \frac{a^2}{2} \sum_{\mu} (\phi_{n+\mu} - \phi_n)^2 + \frac{a^4}{2} m^2 \phi_n^2 \}.$$
 (1)

With the help of displacement operator  $R_{\mu}$ , which is defined as

$$R_{\mu}\phi_n = \phi_{n+\mu},\tag{2}$$

the free scalar action (1) may be written as

$$S(\phi) = \sum_{n} \{ \frac{a^2}{2} \sum_{\mu} [(\partial_{\mu} \phi_n]^2 + \frac{a^4}{2} M^2 \phi_n^2 \}$$

$$= \sum_{n} \{ \frac{a^2}{2} \phi^{\dagger} \Box \phi_n + \frac{a^4}{2} M^2 \phi_n^2 \},$$
(3)

where the discrete derivative  $\partial_{\mu} = \frac{R_{\mu}-1}{a}$  and D'Alembertian operator  $\Box = \sum_{\mu} \frac{R_{\mu}+R_{-\mu}-2}{a^2}$ . Using the formula

$$\phi_n = \left[\int_{\frac{\pi}{a}}^{\frac{\pi}{a}} \frac{dk^4}{(2\pi)^4} e^{ikna} \phi(k)\right],\tag{4}$$

we can transfer (1) into the momentum space form

$$S(\phi) = \frac{1}{2} \int \frac{d^4k}{(2\pi)^4} \phi(-k) \left[\sum_{\mu} \frac{4}{a^2} \sin^2(\frac{ak_{\mu}}{2}) + M^2\right] \phi(k).$$
(5)

The dispersion relation read from this equation is

$$S_{\phi}(k) = \sum_{\mu} \frac{4}{a^2} \sin^2(\frac{ak_{\mu}}{2}) + M^2.$$
 (6)

The naive dicretization of free fermion field is usually taken as

$$S(\psi) = \sum_{n} \{ \frac{a^{3}}{2} \sum_{\mu} \bar{\psi}_{n} (\overrightarrow{\mathcal{P}} - \overleftarrow{\mathcal{P}}) \psi_{n} + M a^{4} \bar{\psi}_{n} \psi_{n} \}$$
$$= \sum_{n} \{ \frac{a^{3}}{2} \sum_{\mu} \bar{\psi}_{n} \gamma_{\mu} (R_{\mu} - R_{-\mu}) \psi_{n} + M a^{4} \bar{\psi}_{n} \psi_{n} \}$$
(7)

where  $\mathcal{D} = \gamma_{\mu} \partial_{\mu}$  is the Dirac operator. Transferring it into momentum space we get

$$S(\psi) = \int \frac{d^4k}{(2\pi)^4} \bar{\psi}(-k) [i\sum_{\mu} \gamma_{\mu} \frac{\sin(ak_{\mu})}{a} + M] \psi(k).$$
(8)

From this equation we see the dispersion relation

$$S_{\psi}(k) = \sum_{\mu} \frac{1}{a^2} \sin^2(ak_{\mu}) + M^2$$
(9)

is different from (6), and the doubling of Fermionic degrees of freedom appears.

In above formulas, two key operators on lattice were introduced. The first is the Dirac operator  $\mathcal{P}$ , the second is the D'Alembertian operator  $\Box$ , which domain the dynamics of fields. Now we may raise a question, why the lattice description of scalar field is successful, while the fermion field fails? One possible way to answer this question is to regard the lattice definition of D'Alembertian operator is "good" but that of Dirac operator is "bad".

So we advocated a new point that all the disaster of lattice Fermions come from the "bad" definition of Dirac operator. Therefore, to overcome the Fermionic difficulties we propose a new lattice Dirac operator, which may shed the new light on studying lattice Fermion fields.

Consulting the definition of the translation operator in the momentum space<sup>1</sup>.

$$R_{\mu}\phi_{n} = R_{\mu} \left[ \int_{\frac{-\pi}{a}}^{\frac{\pi}{a}} \frac{dk^{4}}{(2\pi)^{4}} e^{ikna}\phi(k) \right]$$
  
=  $\phi_{n+\mu} = \int_{\frac{-\pi}{a}}^{\frac{\pi}{a}} \frac{dk^{4}}{(2\pi)^{4}} e^{ikna}\psi(k)e^{ik\mu a}$  (10)

we can define an operator  $R_{\frac{\mu}{2}}$ , corresponding to the "half-spacing translation", as

$$R_{\frac{\mu}{2}}\psi_{n} := \int_{\frac{-\pi}{a}}^{\frac{\pi}{a}} \frac{dk^{4}}{(2\pi)^{4}} e^{ikna}\psi(k)e^{\frac{ik\mu a}{2}};$$
  
$$R_{-\frac{\mu}{2}}\psi_{n} := \int_{\frac{-\pi}{a}}^{\frac{\pi}{a}} \frac{dk^{4}}{(2\pi)^{4}} e^{ikna}\psi(k)e^{-\frac{ik\mu a}{2}}.$$
 (11)

From the definition (11) we can also see that  $R_{\frac{\mu}{2}}$  and  $R_{-\frac{\mu}{2}}$  are not "local operators". When acting on some functions of lattice sites, they not only concern with the nearest neighbor sites but also all the sites in direction  $\mu$  on the lattice. However, it is evident that the combination  $\frac{R_{\frac{\mu}{2}}-R_{-\frac{\mu}{2}}}{a}$  will approximate to  $\partial_{\mu}$  when the lattice spacing  $a \to 0$ . So we can call the combination  $\frac{R_{\frac{\mu}{2}}-R_{-\frac{\mu}{2}}}{a}$  the " quasi-local operator". The quasi-local feature is the price for getting the above mentioned nice characters.

In this approach, the key point is to introduce the auxiliary operator  $R_{\frac{\mu}{2}}$ . Actually, physical fields  $\phi$  and  $\psi$  distribute on the integer sites and the elementary translation operator is  $R_{\mu}$ . To discuss the physical meaning of what we really did, it is desirable to represent the auxiliary operator  $R_{\frac{\mu}{2}}$  in terms of physical operator  $R_{\mu}$ .

By the help of definition (11), it is easy to show

$$R_{\frac{\mu}{2}}\psi_{n} = \frac{a}{2\pi} \sum_{m=-\infty}^{\infty} \int_{\frac{-\pi}{a}}^{\frac{\pi}{a}} dk e^{ik \cdot (n-m+\frac{1}{2})a} \psi_{m}$$
$$= \frac{1}{\pi} \sum_{m=-\infty}^{\infty} \frac{(-1)^{n-m}}{n-m+\frac{1}{2}} R_{\mu}^{m-n} \psi_{n}.$$
(12)

 $<sup>^{1}</sup>$ The similar operator was also introduced in some papers at different angle [8,9]

So in terms of  $R_{\mu}$ , the auxiliary operator  $R_{\frac{\mu}{2}}$  may be written as

$$R_{\frac{\mu}{2}} = \frac{1}{\pi} \sum_{m=-\infty}^{\infty} \frac{(-1)^m}{m + \frac{1}{2}} R^m_{-\mu}.$$
(13)

Similarly,

$$R_{-\frac{\mu}{2}} = -\frac{1}{\pi} \sum_{m=-\infty}^{\infty} \frac{(-1)^m}{m - \frac{1}{2}} R^m_{-\mu}.$$
 (14)

From the above formulas, we see that the operators  $R_{\frac{\mu}{2}}$  and  $R_{-\frac{\mu}{2}}$  are well defined globally, although they were introduced in the first Brillouin zone.

Using the expressions of operators  $R_{\frac{\mu}{2}}$ ,  $R_{-\frac{\mu}{2}}$  in equations (13) and (14) we may prove the following relations hold,

$$R_{\frac{\mu}{2}}^2 = R_{\mu},\tag{15}$$

$$R_{\frac{\mu}{2}}R_{-\frac{\mu}{2}} = R_{-\frac{\mu}{2}}R_{\frac{\mu}{2}} = I.$$
 (16)

If the Dirac operator is redefined as

$$\mathcal{D} = \gamma_{\mu} \left(\frac{R_{\frac{\mu}{2}} - R_{\frac{\mu}{2}}}{a}\right),\tag{17}$$

by the help of equation (13) and (14), it may be written as

$$\mathcal{D} = \frac{2}{\pi a} \sum_{m=1}^{\infty} \frac{(-1)^m m}{m^2 - \frac{1}{4}} \sum_{\mu} \gamma_{\mu} (R^m_{-\mu} - R^m_{\mu}).$$
(18)

It is interesting to find that in this case we may prove the following equation exits

$$\Box = \mathcal{D}^2, \tag{19}$$

which recover the same relation of continuum field theory. In continuous limit  $a \to 0$ ,  $\Box \to \sum_{\mu=1}^{4} \partial_{\mu}^{2}$ , because the square root of D'Alembertian operator is unique in continuum theory, so by the help of equation (19), we show that the continuous limit of the new Dirac operator is just the ordinary Dirac operator.

Substitute the modified Dirac operator into (7), we may get the free fermion action as

$$S(\psi) = \sum_{n} \{a^{3} \frac{2}{\pi} \sum_{m=1}^{\infty} \frac{(-1)^{m} m}{m^{2} - \frac{1}{4}} \sum_{\mu} \bar{\psi}_{n} \gamma_{\mu} (R^{m}_{-\mu} - R^{m}_{\mu}) \psi_{n} + M a^{4} \bar{\psi}_{n} \psi_{n} \}.$$
(20)

which shows how the fermion field on non-neighbour sites correlated each other. When m is taken to 1, (20) reduce to the form of naive discretization action (7). Transform (20) into its momentum space form, we have

$$S(\psi) = \int \frac{d^4k}{(2\pi)^4} \bar{\psi}(-k) \left[\sum_{\mu} \gamma_{\mu}(-\frac{4i}{\pi}) \sum_{m=1}^{\infty} \frac{(-1)^m m \sin(mak_{\mu})}{(m^2 - \frac{1}{4})a} + M\right] \psi(k)$$
(21)

$$= \int \frac{d^4k}{(2\pi)^4} \bar{\psi}(-k) [2i\sum_{\mu} \gamma_{\mu} \frac{\Delta(ak_{\mu})}{a} + M] \psi(k), \qquad (22)$$

where function  $\Delta(x)$  is defined as

$$\Delta(x) = \begin{cases} \sin(\frac{x}{2} - p\pi), & (2p - 1)\pi < x < (2p + 1)\pi \\ 0, & x = (2p + 1)\pi \end{cases}$$
(23)

and may be illustrated by following figure,



From this we see immediately that there is no double-counting anymore and the Fermion field has the same dispersion relation as the one for scalar field (6) one on the boundaries of the Brillouin zone  $\Delta(ak_{\mu}) = 0$ . When M = 0 this new formula (20) will preserve the chiral symmetry.

The next work is to construct the gauge invariant action in this frame. Under gauge transformation of lattice theory,  $R_{\mu}$  will be changed into its covariant form as

$$R_{\mu}\psi_{n} \longrightarrow U_{n,\mu}R_{\mu}\psi_{n}$$

$$R_{-\mu}\psi_{n} \longrightarrow U_{n-\mu,\mu}^{\dagger}R_{\mu}\psi_{n}$$
(24)

where  $U_{n,\mu}$  are the known link variables. Then we have

$$R^{m}_{\mu}\psi_{n} \longrightarrow (U_{n,\mu}R_{\mu})^{m}\psi_{n}$$

$$R^{m}_{-\mu}\psi_{n} \longrightarrow (U^{\dagger}_{n-\mu,\mu}R_{-\mu})^{m}\psi_{n}$$
(25)

Expanding the exponential, one gets

$$(U_{n,\mu}R_{\mu})^{m} = U_{n,\mu}U_{n+\mu,\mu}\cdots U_{n+(m-1)\mu,\mu}R_{\mu}^{m}$$
$$(U_{n-\mu,\mu}^{\dagger}R_{-\mu})^{m} = U_{n-\mu,\mu}^{\dagger}U_{n-2\mu,\mu}^{\dagger}\cdots U_{n-m\mu,\mu}R_{-\mu}^{m}$$
(26)

Now we can build a gauge invariant toy model including a Fermion field  $\psi$ , scalar field  $\phi$  and gauge field  $U_{n,\mu}$ , where  $U_{n,\mu}$  is the link variable of gauge group G. The action of the toy model may be written out as following

$$S = S_{Fermion} + S_{Scalar} + S_{gauge} \tag{27}$$

where

$$S_{Fermion} = \sum_{n} \{ a^{3} \frac{2}{\pi} \sum_{m=1}^{\infty} \frac{(-1)^{m} m}{m^{2} - \frac{1}{4}} \sum_{\mu} \bar{\psi}_{n} \gamma_{\mu} [(U_{n,\mu}^{\dagger} R_{-\mu})^{m} - (U_{n,\mu} R_{\mu})^{m})] \psi_{n} + \lambda (\bar{\psi}_{n}^{L} \phi_{n} \psi_{n}^{R} + \bar{\psi}_{n}^{R} \phi_{n}^{\dagger} \psi_{n}^{L}) \},$$

$$S_{Scalar} = \sum_{n} \{ \frac{a^{2}}{2} \phi^{\dagger} (U_{n,\mu} R_{\mu} + U_{n,\mu}^{\dagger} R_{-\mu} - 2] \phi_{n} ]^{2} + \frac{a^{4}}{2} m^{2} \phi_{n}^{2} \}, \qquad (28)$$

$$S_{Gauge} = \frac{1}{g^2} \sum_{Plaqu} [1 - \frac{1}{2}(U_P + U_P^{\dagger})],$$

where  $\psi^L$  and  $\psi^R$  are the left and right hand Fermion respectively.

With all the previous points in mind, a construction for the  $SU(2) \times U(1)$  electroweak theory on lattice is now available. Consider just the leptonic first generation section: electron and electron neutrino and define the fields as follows:

$$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}, \ R = e_R$$
$$U_{n,\mu}^L = U_{n,\mu} V_{\mu}, U_{n,\mu}^R = V'_{n,\mu}$$
(29)

where  $U_{n,\mu} \exp^{iga\tau \cdot A_{n,\mu}}$  is SU(2) gauge field and  $V_{n,\mu} = \exp^{-\frac{1}{2}ig'aB_{n,\mu}}, V'_{n,\mu} = \exp^{-ig'aB_{n,\mu}}$  are U(1) gauge field. Because the action of scalar field and gauge field parts are easy to write out, we only give the action of fermionic part here,

$$S_{Fermion} = \sum_{n} \{ a^{3} \frac{2}{\pi} \sum_{m=1}^{\infty} \frac{(-1)^{m} m}{m^{2} - \frac{1}{4}} \sum_{\mu} \bar{L}_{n} \gamma_{\mu} [ (U_{n,\mu}^{L} R_{-\mu})^{m} - (U_{n,\mu}^{L} R_{\mu})^{m}) L_{n} + \sum_{n} \{ a^{3} \frac{2}{\pi} \sum_{m=1}^{\infty} \frac{(-1)^{m} m}{m^{2} - \frac{1}{4}} \sum_{\mu} \bar{R}_{n} \gamma_{\mu} [ (G_{n,\mu}^{R} R_{-\mu})^{m} - (U_{n,\mu}^{R} R_{\mu})^{m}) R_{n} + \lambda a^{4} (\bar{L}_{n} \phi_{n} R_{n} + \bar{R}_{n} \phi_{n}^{\dagger} L_{n}) \},$$
(30)

where  $\lambda$  is Yukawa coupling constant and  $\phi = \begin{pmatrix} \phi_+ \\ \phi_0 \end{pmatrix}$  is the Higgs doublet.

In summary, in this paper we have proposed to introduce the Dirac operator  $\mathcal{P} = \frac{2}{\pi a} \sum_{m=1}^{\infty} \frac{(-1)^m m}{m^2 - \frac{1}{4}} (R^m_{-\mu} - R^m_{\mu})$  in lattice theory, which shares the property  $\mathcal{P}^2 = \Box$ . By doing so, we can eliminate the Fermion-doubling problem, get the same dispersion relation for both Boson and Fermion fields and preserve the chiral symmetry. We can also construct the gauge theory in the new frame. Because of these good characters, we can discuss chiral model which is difficult to touch upon in normal lattice theory and we completed the construction of Weinber-Salam Electro-Weak model on lattice. Modifying the Dirac operator is not new in lattice theory, actually, the introduction of Wilson fermion may also be regarded as another kind of modifying method in some sense.

The authors would like to thank Dr. Zenkin and Dr.A.Polychronakos for their useful discussions and Jianming Li also want to give thanks to Professors H.Y Guo, Ke Wu and C.P. Sun for their stimulate discussions and help.

This work is supported in part by the National PanDeng (Clime Up) Plan and the Chinese National Science Foundation and in part by the Chinese Postdoctoral Foundation.

## REFERENCES

- [1] K.G. Wilson, Phys. Rev. D **10**(1974), 2456.
- [2] K.G. Wilson, "New phenomena in subnuclear physics", Part A, ed. by A. Zichichi, 1977 (Plenum Press New York).
- [3] J. Kogut and L. Susskind, Phys. Rev. D 11(1975), 395;
  T. Banks, J. Kogut and L. Susskind, Phys. Rev. D 13(1976), 1043.
- [4] L. Susskind, Phys. Rev. D 16(1977), 3031.
- [5] S. Drell, et al., Phys. Rev. D 14(1976), 1627.
- [6] N.D. Petcher, Nucl. Phys. B**30**(Proc. Suppl.) 50 (1993).
- [7] Ta-Pei Cheng and Ling-Fong Li, "Gauge theory of elementary particle physicd", Clarendon Press, Oxford 1984.
- [8] S.V.Zenkin, Phys.Lett.B **366**(1996),261.
- [9] Jayme Vaz, Advances in Applied Clifford Algebras 7(1997), 37; The Dirac operator over abelian finite group, hep-th/9706189.