# Resolution of the Landau pole problem in QED<sup>\*</sup>

M. Göckeler<sup>a</sup>, R. Horsley<sup>b</sup>, V. Linke<sup>c</sup>, P. Rakow<sup>d</sup>, G. Schierholz<sup>d,e</sup> and H. Stüben<sup>f</sup>

<sup>a</sup>Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg

<sup>b</sup>Institut für Physik, Humboldt Universität zu Berlin, D-10115 Berlin

<sup>c</sup>Institut für Theoretische Physik, Freie Universität Berlin, D-14195 Berlin

<sup>d</sup>Deutsches Elektronen Synchrotron DESY, Institut für Hochenergiephysik und HLRZ, D-15735 Zeuthen

<sup>e</sup>Deutsches Elektronen Synchrotron DESY, D-22603 Hamburg

<sup>f</sup>Konrad-Zuse-Zentrum für Informationstechnik Berlin, D-14195 Berlin

We present new numerical results for the renormalized mass and coupling in non-compact lattice QED with staggered fermions. Implications for the continuum limit and the role of the Landau pole are discussed.

## 1. INTRODUCTION

In the 1950s Landau investigated the relation between the bare charge e and the renormalized charge  $e_R$  in QED. He found [1]

$$\frac{1}{e_R^2} - \frac{1}{e^2} = \frac{N_f}{6\pi^2} \ln \frac{\Lambda}{m_R} \,, \tag{1}$$

where  $\Lambda$  is the momentum cutoff,  $m_R$  is the renormalized mass of the electron and  $N_f$  is the number of flavours (for staggered fermions  $N_f = 4$ ). It is well known that (1) implies two potential problems when  $\Lambda \to \infty$ :

- When e is fixed the theory has a *trivial* continuum limit, i.e.,  $e_R \rightarrow 0$ .
- When  $e_R > 0$  is fixed *e* becomes singular at  $\Lambda_{\text{Landau}} = m_R \exp(6\pi^2/N_f e_R^2)$ . This singularity is called the *Landau pole*.

These problems do not affect the phenomenological success of QED because in practical perturbative calculations the cutoff can be chosen to be large compared with experimental energies. Finding a solution to them is of fundamental theoretical interest and requires non-perturbative methods. We have therefore extensively studied QED on the lattice using non-compact gauge-fields and dynamical staggered fermions [2,3].

In this talk we report on new measurements of the renormalized mass and charge. This was started in [2] and is now extended to lattices of size 16<sup>4</sup> and bare masses am down to 0.005 (*a* denotes the lattice spacing). Using our measurements we have determined functions  $am_R(e^2, am)$  and  $e_R^2(e^2, am)$ . These functions imply that the theory is trivial. They also give a resolution of the Landau pole problem.

#### 2. THE RENORMALIZED MASS

The renormalized mass was obtained from fits to the fermion propagator as explained in [2]. To get  $am_R$  as a function of am and e we need to know the equation of state which relates the bare parameters to the chiral condensate  $\sigma \equiv a^3 \langle \bar{\chi} \chi \rangle$ . In [3] we have found that the  $\sigma$  data obey a mean field equation of state with logarithmic corrections

$$am = A_0 \frac{\sigma^3}{\ln^{p_0}(1/\sigma)} + A_1 \left(\frac{1}{e^2} - \frac{1}{e_c^2}\right) \frac{\sigma}{\ln^{p_1}(1/\sigma)} .(2)$$

We fitted this expression to  $\sigma$  data that were extrapolated to infinite lattice size and obtained  $1/e_c^2 = 0.19040(9), A_0 = 1.798(5),$  $p_0 = 0.324(15), A_1 = 6.76(3), p_1 = 0.485(7),$ 

<sup>\*</sup>Talk presented by H. Stüben



Figure 1. Sketch of the mapping  $(e^2, am) \leftrightarrow (e_R^2, am_R)$ .

 $\chi^2$ /d.o.f = 7.6.

We have observed that  $\sigma$  can be well described by a polynomial in  $am_R$ 

 $\sigma = A_1 a m_R + A_3 a^3 m_R^3 + A_5 a^5 m_R^5 + A_7 a^7 m_R^7 (3)$ 

where the first parameter  $A_1 \equiv 0.6197$  can be taken from perturbation theory and a fit to data from  $12^4$  and  $16^4$  lattices gave  $A_3 = -0.321(5)$ ,  $A_5 = +0.169(13)$ ,  $A_7 = -0.040(7)$ ,  $\chi^2/\text{d.o.f.} =$ 2.1. Because the results of both lattice sizes fall on a universal curve we conclude that the polynomial (3) is also valid on an infinite lattice.

#### 3. THE RENORMALIZED CHARGE

The determination of the renormalized charge has been improved since [2]. The method can only be sketched here. It consists of making a global fit to *all* gauge field propagators D(k) that we have measured

$$\frac{1}{e^2 D(k)} - \frac{1}{e^2} = -\Pi(k, am_R, L) \tag{4}$$

where L is the linear lattice size. The *ansatz* for the fit function  $\Pi$  was taken from [4] to be

$$\Pi = U - \frac{V}{U}\ln(1 - e^2 U) \tag{5}$$

where U is given by 1-loop lattice perturbation theory (see [2]) and where we have set

$$V(k, am_R, L) = v_0 + v_1 U(k, am_R, L).$$
(6)

We then find  $e_R^2(e^2, am_R)$  using  $e_R^2 = Z_3 e^2$  and  $Z_3 = \lim_{k \to 0} \lim_{L \to \infty} D(k)$  from

$$\frac{1}{e_R^2} - \frac{1}{e^2} = -\Pi(0, am_R, \infty) \,. \tag{7}$$

A simultaneous fit of (4) to the gauge field propagators at our 52 values  $(e^2, am, L)$  gave  $v_0 =$  $-0.00207(2), v_1 = -0.0328(7), \chi^2/\text{d.o.f.} = 1.7.$ Since  $U(0, am_R, \infty) \approx (N_f/6\pi^2)(-0.31 + \ln am_R)$ we only find small corrections to the old result (1).

#### 4. DISCUSSION

We can now discuss the mapping  $(e^2, am) \leftrightarrow (e_R^2, am_R)$ . A global qualitative view of this mapping is shown in Figure 1, while a quantitative plot of  $(1/e^2, am) \leftrightarrow (1/e_R^2, am_R)$  is shown in Figure 2.

In both figures accessible regions are plotted in grey. The whole plane of bare parameters is accessible but this plane is mapped only onto a part of the plane of the renormalized parameters.



Figure 2. Quantitative picture of the mapping  $(1/e^2, am) \leftrightarrow (1/e_R^2, am_R)$ . In the plane of the bare parameters lines of constant  $1/e_R^2$  are shown indicating the renormalization group flow of  $e_R^2$ . In the plane of the renormalized parameters the lines of constant am are shown.

The border of the accessible region is shown as a thick line. It is the image of the corresponding thick line on the am = 0 line starting/ending at the critical value of the coupling constant.

On the line  $am_R = 0$  in Figure 1 the only accessible point is the origin. This reflects triviality. In Figure 2 triviality is expressed by the fact that no line of (finite) constant  $1/e_R^2$  flows into the critical point.

The dotted line in both figures is the position of the Landau pole, i.e., the line of pairs  $(e_R^2, am_R)$ with  $1/e^2 = 0$  from (7) with  $\Pi \equiv U$ . This line is well separated from the border for all finite  $e_R^2$ .

# 5. CONCLUSIONS

From the presented analysis we conclude that non-compact lattice QED with staggered fermions has a trivial continuum limit. In addition our analysis implies a resolution of the Landau problem. The resolution is that for given e and am the theory does not allow arbitrary choices of  $am_R$ . Instead through chiral symmetry breaking the theory itself provides a minimal lattice spacing or maximal cutoff which is below  $\Lambda_{\text{Landau}}$ . Non-perturbatively the function  $\Pi$  is very close to what one finds in perturbation theory, but there is no Landau pole problem because the pole always lies in the forbidden region.

### ACKNOWLEDGEMENTS

Our new measurements were done on the CRAY T3D at the Konrad-Zuse-Zentrum Berlin. Financial support by the Deutsche Forschungsgemeinschaft is gratefully acknowledged.

# REFERENCES

- L. D. Landau, in: Niels Bohr and the Development of Physics, ed. W. Pauli, Pergamon Press, London, 1955.
- M. Göckeler, R. Horsley, P. Rakow, G. Schierholz and R. Sommer, Nucl. Phys. B371 (1992) 713.
- M. Göckeler, R. Horsley, V. Linke, P. Rakow, G. Schierholz and H. Stüben, Nucl. Phys. B487 (1997) 313.
- 4. D.V. Shirkov, Nucl. Phys. B371 (1992) 467.