

Spacetime Fluctuations and the Cosmological Constant Problem

J. W. Moffat

*Department of Physics, University of Toronto, Toronto, Ontario M5S 1A7,
Canada*

Abstract

Spacetime geometry is treated as a fluctuating, stochastic quantum system allowing an effective quantum gravity solution to the cosmological constant problem. A Fokker-Planck equation for the probability density of spacetime metric fluctuations is solved for an FRW universe and the most probable value for the metric scale factor R is for a very small cosmological constant.

e-mail: moffat@medb.physics.utoronto.ca

1 Introduction

The cosmological constant problem continues to defy all attempts to solve it in a natural way that avoids any fine-tuning of parameters [1, 2, 3, 4]. Recent investigations of this problem have focused on bulk-brane scenarios in which the standard model fields are confined to a $(3+1)$ brane and gravity and moduli fields propagate in the higher-dimensional bulk. The Randall-Sundrum models and various extensions of the Rubakov-Shaposhnikov model have shown that a fine-tuning of brane tension constants and the bulk cosmological constant cannot be avoided [5]. Moreover the ‘self-tuning’ models have also been confronted with fine-tuning problems related to the existence of naked singularities. These attempted solutions to the problem are based on classical or semi-classical models and the issue of quantum gravity and other quantum field theory vacuum corrections remains to be resolved. It is not yet understood how string theory and D-brane theory can solve the cosmological constant problem [2]. The supersymmetry solution to the cosmological constant problem is prevented from being applicable, because supersymmetry must be broken for energies below 1 TeV, and the fine-tuning of vacuum

contributions to the cosmological constant occurs already at the standard model energy scale, $E_{SM} \sim 10^2$ GeV, and lower energies.

An early attempt by Hawking [6, 7] to solve the cosmological constant problem was based on quantum cosmology and the existence of a four-index tensor in four-dimensional spacetime. The wave function of the universe led to a functional integral in Euclidean space with a probability distribution for the wave function of the universe that favours a vanishingly small cosmological constant. Coleman's wormhole solution is also based on the probabilistic interpretation that the universe with the smallest cosmological constant is the most probable [8]. However, as in the case of Hawking's proposal, it suffers from the difficulty of interpreting Euclidean space gravity and other technical problems. The anthropic principle solution [9] is another attempt to resolve the problem, but it does not explain how to solve the vanishingly small cosmological constant given the current theory of particle physics and gravitation and its parameters.

Because we do not yet possess a satisfactory quantum gravity theory, the problem of how to treat the linkage between general relativity (GR) and quantum field theory remains somewhat in the dark. The standard semi-classical approach, in which we write Einstein's field equations as

$$G_{\mu\nu} - \Lambda_0 g_{\mu\nu} = 8\pi G \langle T_{\mu\nu} \rangle, \quad (1)$$

where $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$, Λ_0 is the 'bare' Einstein cosmological constant and $\langle T_{\mu\nu} \rangle$ denotes the expectation value of the stress-energy tensor treated as a quantum operator, is basically inconsistent, for the left-hand side of the equation can only be reconciled with the right-hand side by making the gravitational field change in a discontinuous, acausal manner, a behavior unacceptable from the classical point of view.

An 'effective' quantum gravity interpretation of spacetime can be developed, in which the metric tensor $g_{\mu\nu}$ is treated as a stochastic random variable, which is defined by a set of possible values and a probability distribution $P(g)$ over this set with $P(g) \geq 0$ [10, 11]. A $(3+1)$ foliation of spacetime in GR is combined with the canonical formalism to derive a Langevin equation for the conjugate canonical momentum, which is treated as a random variable. In this formalism, both the gravitational field and the matter or vacuum are treated in a probabilistic fashion, so that we have

$$\langle G_{\mu\nu} \rangle - \Lambda_0 \langle g_{\mu\nu} \rangle = 8\pi G \langle T_{\mu\nu} \rangle. \quad (2)$$

Thus, there is now no conflict between the two sides of the equation.

By converting the Friedmann equation for a Friedmann-Robertson-Walker (FRW) cosmology into a first-order stochastic differential equation, we can calculate the approximate stationary probability density for the cosmological scale factor R , and we find that the most probable value for R in our universe occurs for a very small cosmological constant.

2 Stochastic Differential Equation for the Expansion of the Universe

Wheeler and his co-workers [12] suggested that quantum fluctuations in the metric tensor of spacetime, $g_{\mu\nu}$, should occur at the order of the Planck length: $L_P = (\hbar G/c^3)^{1/2} = 1.6 \times 10^{-33}$ cm. These fluctuations are of order

$$\Delta g \sim \frac{L_P}{L}, \quad (3)$$

where L is the spatial dimension of the region in a local Lorentz frame of reference. These fluctuations are expected to be significant at the Planck scale $\sim 10^{-33}$ cm.

Let us consider a Friedmann-Robertson-Walker (FRW) spacetime with the metric

$$ds^2 = dt^2 - R^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right], \quad (4)$$

where $k = 0, 1, -1$ corresponding to a flat, closed and open universe, respectively. The Einstein field equations are ($c=1$):

$$\dot{R}^2 + k = \frac{8\pi G}{3} \rho R^2 + \frac{\Lambda_0}{3} R^2, \quad (5)$$

$$2R\ddot{R} + \dot{R}^2 + k = -8\pi G p R^2 + \Lambda_0 R^2, \quad (6)$$

where $\dot{R} = \partial R / \partial t$.

From Eq.(5), we can obtain the first order equation

$$\dot{R} = \left(\frac{8\pi G}{3} \rho R^2 + \frac{\Lambda_0}{3} R^2 - k \right)^{1/2}, \quad (7)$$

where we have chosen the positive square root for an expanding universe. Including geometrical quantum randomness in the gravitational equations, leads to the Langevin stochastic equation for the cosmic scale factor R [13, 14]:

$$dR_t = f(R_t)dt + \sigma \xi_t h(R_t)dt, \quad (8)$$

where R_t denotes the random scale factor, σ denotes the intensity of the quantum gravity fluctuations and ξ_t is Gaussian fluctuations with $\langle \xi_t \rangle = 0$. We can take into account the fluctuations of the spacetime geometry, in an effective way, by replacing the Planck length $L_P = \sqrt{G}$ by a stationary random variable [10]:

$$L_{tP} \equiv \sqrt{G_t} = \sqrt{G} + \sigma \xi_t, \quad (9)$$

where \sqrt{G} denotes the mean value.

The extreme irregularity of Gaussian white noise means that the time derivative of R_t is not well defined. However, the standard theory of stochastic processes can handle this difficulty by defining the equivalent integral equation

$$R_t = R_0 + \int_{t_0}^t f(R_y)dy + \sigma \int_{t_0}^t h(R_y)\xi_y dy. \quad (10)$$

Two definitions of the stochastic integral $\int h(R_y)\xi_y dy$ have been given by Ito [15] and Stratonovich [16]. Both the Ito and the Stratonovich definitions of the integral lead to a consistent calculus. The transition probability density $p(R, t)$ of the scale factor diffusion process R_t satisfies the Fokker-Planck equation (FPE):

$$\partial_t p(R, t) = -\partial_R f(R, t)p(R, t) + \frac{\sigma^2}{2} \partial_R^2 [h^2(R, t)p(R, t)]. \quad (11)$$

An important and useful solution of the FPEs can be obtained for stationary random metric processes. We expect that a quantum gravitational system subjected to metric fluctuations for a sufficiently long time will settle down to a stationary behavior. This means that as time approaches infinity, the system will reach a state for which the probability density $p_S(R)$ has a shape that does not change with time i.e. the probability to find the system in the neighborhood of a particular state becomes time independent. However, the sample paths $R_t(\omega)$ will in general not approach a steady-state value $R_s(\omega)$, so that the state of the system still fluctuates. However, these fluctuations are such that R_t and $R_{t+\tau}$ have the same probability density, $p_S(R)$.

The stationary solution $p_S(R)$ of the FPE satisfies

$$\partial_t p(R, t|R_i, t_i) + \partial_R J(R, t|R_i, t_i) = 0, \quad (12)$$

where R_i and t_i denote the initial values of R and t , respectively, and J is the probability flow current

$$J(R, t|R_i, t_i) = f(R)p(R, t|R_i, t_i) - \frac{\sigma^2}{2} \partial_R [h^2(R)p(R, t|R_i, t_i)]. \quad (13)$$

The stationary FPE is then given by

$$\partial_R J_S(R) = 0, \quad (14)$$

which implies that $J_S(R) = \text{constant}$ for $R \in [R_1, R_2]$. In the stationary case, there are no sources or sinks in the universe for the probability. We have the equation for the probability density $p_S(R)$:

$$-f(R)p_S(R) + \frac{\sigma^2}{2} \partial_R h^2(R)p_S(R) = -J. \quad (15)$$

We define the auxiliary function $u(R) = h^2(R)p_S(R)$ and obtain the solution to the stationary FPE:

$$\begin{aligned} p_S(R) &= \left(\frac{N}{h^2(R)} \right) \exp\left(\frac{2}{\sigma^2} \int_{R_i}^R \frac{f(x)}{h^2(x)} dx \right) \\ &\quad - \left(\frac{2}{\sigma^2 h^2(R)} \right) J \int_{R_i}^R \exp\left(\frac{2}{\sigma^2} \int_y^R \frac{f(x)}{h^2(x)} dx \right) dy. \end{aligned} \quad (16)$$

We have $J = 0$ when there is no flow of probability out of the universe. By using the Ito prescription, we get

$$p_S(R) = \left(\frac{N}{h^2(R)} \right) \exp\left(\frac{2}{\sigma^2} \int_{R_i}^R \frac{f(x)}{h^2(x)} dx \right). \quad (17)$$

In this case, the normalization constant N is given by

$$N^{-1} = \int_{R_1}^{R_2} \frac{dR}{h^2(R)} \exp\left(\frac{2}{\sigma^2} \int_{R_i}^R \frac{f(x)}{h^2(x)} dx \right) < \infty. \quad (18)$$

3 Metric Fluctuations and the Cosmological Constant

Let us now apply our effective, stochastic quantum gravity formalism to the cosmological constant problem. We shall assume that matter is dominated by the vacuum density ρ_{vac} with the equation of state: $\rho = -p = \rho_{\text{vac}} = \text{constant}$ and that the universe is spatially flat $k = 0$. Then, we have from Eq. (7):

$$dR = \left(\frac{8\pi}{3}\right)^{1/2} \sqrt{G\bar{\rho}_{\text{vac}}^{-1/2}} R dt, \quad (19)$$

where

$$\bar{\rho}_{\text{vac}} = \rho_{\text{vac}} + \frac{\Lambda_0}{8\pi G} = \frac{\Lambda}{8\pi G}. \quad (20)$$

We obtain for R the classical de Sitter solution

$$R = R_i \exp\left(\sqrt{\frac{\Lambda}{3}} t\right), \quad (21)$$

where R_i is the initial value of R at $t = 0$.

By replacing R and \sqrt{G} in Eq.(19) by the random variables R_t and $\sqrt{G_t}$, we obtain the stochastic differential equation (8) with

$$f(R_t) = \left(\frac{8\pi}{3}\right)^{1/2} \sqrt{G_t} \bar{\rho}_{\text{vac}}^{-1/2} R_t, \quad (22)$$

$$h(R_t) = \left(\frac{8\pi}{3}\right)^{1/2} \bar{\rho}_{\text{vac}}^{-1/2} R_t. \quad (23)$$

The approximate stationary probability density is given by

$$\begin{aligned} p_S(R) &= N\left(\frac{3GR_i^2}{\Lambda R^2}\right) \exp\left[\left(\frac{2G}{\sigma^2}\right) \sqrt{\frac{3}{\Lambda}} \ln(R/R_i)\right] \\ &= N\left(\frac{3G}{\Lambda}\right) \left(\frac{R}{R_i}\right)^{\beta-2}, \end{aligned} \quad (24)$$

where $\beta = \left(\frac{2G}{\sigma^2}\right) \sqrt{\frac{3}{\Lambda}}$ and we have used Eq.(20) to define Λ .

We arrive at the result that the maximum probability for the scale factor R , for a fixed value of the fluctuation intensity σ and $R/R_i > 1$, occurs for a very small cosmological constant Λ . We illustrate this result in Fig 1., where we have plotted the stationary probability density p_S as a function of R and Λ for an arbitrary normalization and for decreasing values of Λ .




Fig. 1. Three-dimensional plot of stationary probability density p_S versus R and Λ for arbitrary normalization and dimensionless units.

4 Discussion of Results and Conclusions

It is important to recognize that the cosmological constant Λ is the true effective cosmological constant, that would be measured in gravitational phenomena at long ranges. It includes all effects of fields and any effects due to phase transitions, such as those that occur at the electroweak energy scale, 10^2 Gev, and the QCD energy scale. We see that p_S peaks sharply as Λ decreases for fixed values of the fluctuation intensity σ and for $R/R_i > 1$. Thus, we conclude that the most likely value of R for all possible universes occurs for a very small cosmological constant.

The intensity of the vacuum fluctuations of spacetime are expected to be strong at the Planck energy $\sim 10^{19}$ Gev, where quantum gravitational effects begin to dominate. However, this is the energy scale at which most models of inflation begin to inflate [17]. Although the quantum gravity fluctuations lead to a small probable value of the cosmological constant in the early expansion of the universe, this contradicts the usual expectation that it is huge during the inflation period. This just underscores the cosmological constant

problem faced by scalar field, potential-driven models of inflation, namely, the need for a very large cosmological constant initially in the universe, and the subsequent observationally small bound on the constant in the present universe:

$$|\Lambda| \leq 10^{-56} \text{ cm}^{-2} \sim 10^{-84} \text{ GeV}^2. \quad (25)$$

For the vacuum energy, the bound is

$$\rho_{\text{vac}} \leq 10^{-29} \text{ g/cm}^3 \sim 10^{-47} \text{ GeV}^4 \sim 10^{-9} \text{ ergs/cm}^3. \quad (26)$$

In our scenario, this would presuppose that to uphold a large value of the cosmological constant (vacuum energy) would require that spacetime is very smooth at the beginning of the universe with $\sigma \sim 0$ in contradiction with our intuitive expectations that quantum gravity effects should be very strong at the Planck energy scale. If the vacuum energy in the universe is truly constant, as is demanded for ordinary matter by the Bianchi identity: $\nabla_\nu T^{\mu\nu} \equiv 0$ and $\nabla_\nu g^{\mu\nu} = 0$ in Einstein's field equations, then when this constant is required to be zero at the Planck energy scale, it should remain zero for the rest of the universe's evolution. This scenario would avoid the embarrassing situation, in which we have to somehow guarantee that Λ relaxes to zero (or almost zero) whenever there is a new phase transition as the universe evolves in time. If there exists exotic 'quintessence' matter [18], then a time dependent 'cosmological constant' could exist and the aforementioned difficulty would prevail. Moreover, we are also faced in this case with the difficulty of explaining the coincidence of the quintessence dark energy density with the present value of the matter energy density (including cold dark matter). At present, the acceleration of the expansion of the universe detected by supernovae data does not exclude a fit to all the current observational data with a constant Λ .

Our explanation of the smallness of the cosmological constant is based on an effective quantum gravity, stochastic argument, in which a probability density is peaked at a vanishingly small cosmological constant. The result is similar in spirit to the Baum, Hawking and Coleman explanations [6, 7, 8] (for a detailed critique of these papers, see ref. [1]). These proposals are based on a solution of the wave function of the universe Ψ as an Euclidean path integral

$$\Psi \propto \int [dg][d\Phi] \exp(-S[g, \Phi]), \quad (27)$$

where S is the Einstein-Hilbert gravity action including matter terms, and the integral is over all Euclidean 4-metrics g and matter fields Φ defined on a

4-manifold. Hawking uses a three index field $B_{\mu\nu\sigma}$ and Coleman uses worm-holes as topological structures to obtain an effective cosmological constant. There have been several important criticisms of the underlying formalism, including the problem of having to use a complex transformation to an Euclidean spacetime geometry, in which the Euclidean action S is unbounded from below. In our effective quantum gravity formalism, we have not introduced any new fields beyond the standard Einstein-Hilbert action, but we have based our probability analysis on a phenomenological stochastic formalism, using a solution of the Fokker-Planck equation to obtain our probability distribution. This formalism is well-founded in many applications to fluctuating physical phenomena, and we suppose that it also applies to metric spacetime fluctuations at the Planck energy scale.

The quantum gravity, stochastic fluctuations can be considered as the fluctuations of micro-objects, e.g. strings or D-branes, which are expected to dominate spacetime physics at the Planck energy. In this way, our stochastic, effective quantum gravity formalism can be considered as a phenomenological picture of how string and D-brane theory could resolve the cosmological constant problem. However, the problem of supersymmetry breaking of string and D-brane theory would have to be fully understood in such a scenario.

Acknowledgments

I thank Michael Clayton for helpful discussions. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

References

- [1] S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989).
- [2] E. Witten, *The cosmological constant problem from the point of view of string theory*, talk in: *Dark Matter 2000*, Marina del Rey, California, February 2000. hep-ph/0002297.
- [3] N. Straumann, *On the mystery of the cosmic vacuum energy density*, astro-ph/0009386.

- [4] S. E. Rugh and H. Zinkernagel, *The quantum vacuum and the cosmological constant problem*, hep-th/0012253.
- [5] V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. **B125**, 139 (1983); L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999). hep-th/9906064; Arkani-Hamed, S. Dimopoulos, N. Kaloper, and R. Sundrum, Phys. Lett. **480**, 193 (2000). hep-th/0001197; S. Kachru, M. Schulz, and E. Silverstein, Phys. Rev. **D62**, 045021 (2000). hep-th/0001206; S. Foerste, Z. Lalak, S. Lavignac, and H. P. Nilles, Phys. Lett. **B481**, 360 (2000). hep-th/0002164; C. Csaki, J. Erlich, C. Grojean, and T. Hollowood, Nucl. Phys. **B584**, 359 (2000). hep-th/0004133; H. Collins and B. Holdom, hep-th/0009127; C. Csaki, J. Erlich, and C. Grojean, hep-th/0012143.
- [6] E. Baum, Phys. Lett. **133B**, 185 (1983).
- [7] S. W. Hawking, **134B**, 403, (1984).
- [8] S. Coleman, Nucl. Phys. **B310**, 643 (1988).
- [9] S. Weinberg, Phys. Rev. Lett. **59**, 2607 (1987); S. M. Bar, J. F. Donoghue, and D. Seckel, Phys. Rev. **D57**, 5480; S. Weinberg, *The cosmological constant problems*, talk in: *Dark Matter 2000*, Marina del Rey, California, February 2000. astro-ph/0005265.
- [10] J. W. Moffat, Phys. Rev. **56**, 6264 (1997).gr-qc/9610067.
- [11] In the preceding reference [9], Eq. (165) should be replaced by
- $$p_S(R) \sim C \left(\frac{3R^2}{8\pi\rho_{0,r}} \right) \exp \left[\left(\frac{2}{\sigma^2} \right) \left(\frac{3G}{32\pi} \right)^{1/2} \left(\frac{R^2}{\rho_{0,r}^{1/2}} \right) \right]. \quad (28)$$
- The conclusions regarding the smearing of the big bang singularity remain the same.
- [12] J. A. Wheeler, *Relativity, Groups and Topology*, lectures delivered at Les Houches, 1963, edited by B. DeWitt and C. DeWitt, Gordon and Breach, New York, p 317; C. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, J. Freeman, San Francisco, 1973.
- [13] B. Øksendal, *Stochastic Differential Equations, an Introduction with Applications*, Springer, Berlin, 1985.

- [14] W. Horsthemke and R. Lefever, *Noise-Induced Transitions, Theory and Applications in Physics, Chemistry, and Biology*, Springer-Verlag, Berlin, 1984.
- [15] K. Ito, Proc. Imp. Acad. **20**, 519 (1944).
- [16] L. Stratonovich, *Conditional Markov Processes and their Applications to the Theory of Optimal Control*, Elsevier, New York, 1968.
- [17] R. H. Brandenberger, *A status review of inflationary cosmology*, hep-ph/0101119.
- [18] S. Perlmutter et al. Astrophys. J. **517**, 565 (1999); A. G. Riess, et al. Astron. J. **116**, 1009 (1998); P. M. Garnavich, et al. Astrophys. J. **509**, 74 (1998); R. R. Caldwell, R. David, and P. J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998); M. S. Turner and M. White, Phys. Rev. **D56**, R1139 (1997).