QCD sum rules for ρ mesons in nuclear matter

S. Mallik^{*}

Saha Institute of Nuclear Physics, 1/AF, Bidhannagar, Calcutta 700 064, India

Andreas Nyffeler[†]

Centre de Physique Théorique, CNRS Luminy, Case 907, F-13288 Marseille Cedex 9, France

(22 May 2001)

We investigate QCD sum rules for vector currents in the rho meson channel in the nuclear medium. For increased sensitivity, we subtract out the vacuum contributions. With a saturation scheme often considered in the literature, we find these "subtracted" sum rules to be unstable. It indicates the importance of other contributions neglected in the saturation scheme. These include more Landau singularities at low energy, additional operators in the medium, and possibly the in-medium width of the rho meson.

PACS number(s): 11.55.Hx, 12.38.Lg, 24.85.+p, 14.40.Cs

I. INTRODUCTION

It was quite some time ago that the original vacuum QCD sum rules [1] were first extended to a medium at finite temperature and density [2,3]. Since then much work has been devoted to extracting results from them. Typically one tries to predict the medium dependence of the parameters of particles and resonances in the low energy region. However, these in-medium sum rules have several new features. These must be included quantitatively in the saturation scheme for reliable predictions.

One such feature, recognized soon after their initial formulation, is that the in-medium sum rules have contributions from new operators over those already present in the vacuum sum rules [4,5]. The velocity four-vector of the medium allows one to construct additional Lorentz scalar operators. (In the rest frame of the medium they are simply the time components of Lorentz tensor operators.) At higher dimensions there arises a multitude of such operators.

The other features relate to the hadron spectral function. A communicating single particle state may acquire a large width (in addition to a possible shift in mass) due to its scattering from the particles in the medium, so that it may not be possible to treat it in the narrow width approximation as in the vacuum. Also multiparticle states in the medium may give rise to several nearby singularities of the Landau type.

In this work we reexamine the sum rules in the ρ -meson channel in the nuclear medium at zero temperature. Our saturation scheme is similar to one mostly employed in the literature, which includes only some of these features [6–8]. In the operator product expansion we include properly all the operators up to dimension 4 and only the four-quark operators at dimension 6. In the spectral function we take the ρ meson in the narrow width approximation and the $N\bar{N}$ state.¹

A complete or "full" in-medium sum rule is actually the corresponding vacuum sum rule, perturbed by small, density dependent terms. Such a "full" sum rule, when evaluated, is guaranteed to be stable, more or less, irrespective of these small contributions, simply because the vacuum sum rule is so. To get rid of this insensitivity, we subtract out the vacuum sum rule from the "full" one, retaining on both sides only the terms to first order in the density.

Though well-known, these sum rules are derived here to bring out some technical points. We start with a broader framework, namely the ensemble average of the correlation function of currents at finite temperature and chemical potential. The special cases of sum rules at finite temperature and at finite density may then be recovered from the general formulation by setting respectively the chemical potential and the temperature to zero. We derive the mixing of two operators of dimension 4, namely the quark and the gluon parts of the energy momentum tensor, a result known in the context of deep inelastic scattering. We also present a field theoretic derivation for the $N\bar{N}$ contribution to the spectral function.

^{*}Electronic address: mallik@tnp.saha.ernet.in

[†]Electronic address: nyffeler@cpt.univ-mrs.fr

¹Actually there could be further density independent contributions in the continuum, but they are eliminated in the subtracted sum rules we shall be considering.

On evaluation we find that the sum rules as saturated above do not yield stable results, as we have already reported recently [9]. This instability points to the importance of other contributions we did not include, namely, the Lorentz nonscalar operators at dimension 6 and the Landau singularities besides the one from the $N\bar{N}$ state. The neglect of the (large) in-medium width of the ρ -meson also possibly adds to this instability [10,11].

In Sec. II we introduce the kinematic decomposition of the two point function of vector currents. In Sec. III a simple derivation of the operator product expansion is presented in coordinate space. Then in Sec. IV we construct the spectral representation from ρ and $N\bar{N}$ intermediate states at finite density. The sum rules are written and evaluated in Sec. V. We conclude in Sec. VI.

II. KINEMATICS

Consider the ensemble average of the time ordered (T) product of two currents

$$\mathcal{T}^{ab}_{\mu\nu}(q) = i \int d^4x e^{iq \cdot x} \left\langle T \left\{ V^a_\mu(x) V^b_\nu(0) \right\} \right\rangle.$$
(2.1)

Here $V^a_{\mu}(x)$ is the vector current (in the ρ -meson channel) in QCD,

$$V^{a}_{\mu}(x) = \bar{q}(x)\gamma_{\mu}\frac{\tau^{a}}{2}q(x), \qquad (2.2)$$

q(x) being the field of the u and d quark doublet and τ^a the Pauli matrices. The ensemble average of an operator O is denoted by $\langle O \rangle$,

$$\langle O \rangle = \operatorname{Tr} \left[e^{-\beta (H-\mu N)} O \right] / \operatorname{Tr} \left[e^{-\beta (H-\mu N)} \right], \qquad (2.3)$$

where H is the QCD Hamiltonian, β is the inverse of the temperature T, and Tr denotes the trace over any complete set of states. Keeping our application in mind, we have introduced the nucleon chemical potential μ with the corresponding number operator N.

For explicit calculations one generally chooses the rest frame of the medium, where the temperature is defined. This breakdown of Lorentz invariance may be restored if we let the medium have an arbitrary four-velocity u_{μ} [12]. [In the matter rest frame $u_{\mu} = (1, 0, 0, 0)$.] The time and space components of q_{μ} are then raised to the Lorentz scalars, $w = u \cdot q$ and $\bar{q} = \sqrt{w^2 - q^2}$.

There are two independent, conserved kinematic covariants [13], in which the two point function from Eq. (2.1) may be decomposed. In choosing their forms we must note that the dynamical singularities extend up to $q^2 = 0$. We thus have to ensure that the kinematic covariants do not have any singularity in this region, particularly at $q^2 = 0$. The covariants $P_{\mu\nu}$ and $Q_{\mu\nu}$ defined by

$$P_{\mu\nu} = -g_{\mu\nu} + \frac{q_{\mu}q_{\nu}}{q^2} - \frac{q^2}{\bar{q}^2}\tilde{u}_{\mu}\tilde{u}_{\nu}, \qquad Q_{\mu\nu} = \frac{q^4}{\bar{q}^2}\tilde{u}_{\mu}\tilde{u}_{\nu}, \tag{2.4}$$

where $\tilde{u}_{\mu} = u_{\mu} - wq_{\mu}/q^2$, are indeed so, as can be seen by inspecting their components for space and time indices. The invariant decomposition of $\mathcal{T}^{ab}_{\mu\nu}$ is then given by

$$\mathcal{T}^{ab}_{\mu\nu}(q) = \delta^{ab}(Q_{\mu\nu}T_l + P_{\mu\nu}T_t), \qquad (2.5)$$

where the invariant amplitudes $T_{l,t}$ are functions of q^2 and w.

The kinematic covariants are still not regular: they have a dependence on the direction of the three-vector \vec{q} even as $|\vec{q}| \to 0$. Thus the spatial components of $\mathcal{T}^{ab}_{\mu\nu}$ in this limit become

$$\mathcal{T}_{ij}^{ab}(q) = \delta^{ab} [\delta_{ij} T_t - n_i n_j (T_t - q_0^2 T_l)], \qquad (2.6)$$

where n_i is the *i*th component of the unit vector along \vec{q} . To eliminate this direction dependence, we must require that [2]

$$T_t(q_0, |\vec{q}| = 0) = q_0^2 T_l(q_0, |\vec{q}| = 0).$$
(2.7)

III. OPERATOR PRODUCT EXPANSION

Here we outline the derivation of the short distance expansion of the product of two current operators in Eq. (2.1). As already stated, the availability of the velocity four-vector u^{μ} of the medium allows one to construct new, independent Lorentz scalar operators. Thus up to dimension 4 we have, in addition to the old ones **1**, $\bar{q}q = \bar{u}u + \bar{d}d$, $G^2 \equiv (\alpha_s/\pi)G^a_{\mu\nu}G^{\mu\nu a}$, appearing in the vacuum expectation value, two new ones $\Theta^q \equiv u^{\mu}\Theta^q_{\mu\nu}u^{\nu}$ and $\Theta^g \equiv u^{\mu}\Theta^g_{\mu\nu}u^{\nu}$ for the ensemble average. Here $G^c_{\mu\nu}$, c = 1, ..., 8 are the gluon field strengths and $\alpha_s = g_s^2/4\pi$, g_s being the QCD coupling constant. $\Theta^{q}_{\mu\nu}$ are the quark and the gluon parts of the traceless energy momentum tensor

$$\Theta^{q}_{\mu\nu} = \bar{q}i\gamma_{\mu}D_{\nu}q - \frac{\hat{m}}{4}g_{\mu\nu}\bar{q}q,$$

$$\Theta^{g}_{\mu\nu} = -G^{c}_{\mu\lambda}G^{\ \lambda c}_{\nu} + \frac{1}{4}g_{\mu\nu}G^{c}_{\alpha\beta}G^{\alpha\beta c},$$
(3.1)

where \hat{m} is the quark mass in the limit of SU(2) symmetry.

We now obtain the (singular) coefficients of the operators in coordinate space [14–17]. In the free field theory these are obtained from the Wick decomposition of the operator product. If we treat the gauge fields as external, the same form of decomposition also holds for the interacting theory

$$T \{ V^{a}_{\mu}(x) V^{a}_{\nu}(0) \} = \operatorname{tr}[\gamma_{\mu} S(x,0) \gamma_{\nu} S(0,x) \tau^{a} \tau^{b}/4] -i\bar{q}(0) \gamma_{\nu} S(0,x) (\tau^{b} \tau^{a}/4) q(x) - i\bar{q}(x) \gamma_{\mu} S(x,0) \gamma_{\nu} (\tau^{a} \tau^{b}/4) q(0) + \operatorname{a regular term.}$$
(3.2)

The trace in the first term is over all the indices, namely spin, flavor and color of the quark field. The quark propagator S(x, 0) is now in the presence of the (matrix valued) gauge potential $A_{\mu}(x)$,

$$\{-i\gamma^{\mu}[\partial_{\mu} - igA_{\mu}(x)] + \hat{m}\} S(x,0) = \delta^{4}(x).$$
(3.3)

An important technical point here is that quantities can be put in a gauge covariant form from the beginning, if we use the Fock-Schwinger gauge, in which, $x^{\mu}A_{\mu}(x) = 0$. In this gauge

$$A_{\mu}(x) = (1/2)x^{\alpha}G_{\alpha\mu}(0) + \cdots,$$

$$q(x) = q(0) + x^{\mu}D_{\mu}q(0) + \cdots.$$
(3.4)

Equation (3.3) may be solved for the quark propagator in a series of increasing number of gluon field strengths.

The unit and the quark operators 1, $\bar{q}q$, and Θ^q reside in the first and the next two terms in Eq. (3.2), respectively. To calculate their coefficients, it suffices to use the free quark propagator

$$S_0(x,0) = \frac{1}{4\pi^2} \left(-\frac{2\not x}{(x^2 - i\epsilon)^2} - \frac{i\hat{m}}{x^2 - i\epsilon} \right), \tag{3.5}$$

where we retain only the leading singular term in \hat{m} . The quark operators in Eq. (3.2) can then be projected on to the scalar operators by using the formulae

$$\langle \bar{q}_A^j q_B^k \rangle = \frac{1}{8} \delta^{jk} (\delta_{BA} \langle \bar{q}q \rangle + (\psi)_{BA} \langle \bar{q}\psi q \rangle), \tag{3.6}$$

$$\langle \bar{q}_{A}^{j} i(D_{\mu}q)_{B}^{k} \rangle = \frac{1}{2} \delta^{kj} \left\{ \frac{1}{4} \delta_{BA} u_{\mu} \hat{m} \langle \bar{q} \psi q \rangle + (\gamma_{\mu})_{BA} \left(\frac{\hat{m}}{16} \langle \bar{q}q \rangle - \frac{1}{12} \langle \Theta^{q} \rangle \right) + u_{\mu}(\psi)_{BA} \frac{1}{3} \langle \Theta^{q} \rangle \right\}.$$

$$(3.7)$$

The flavor and spinor indices of the quark field have been denoted by (j, k) and (A, B), respectively. A sum over color indices on both sides is understood. These projection formulae are easily obtained by expanding the quark bilinear matrix in terms of the complete set of Dirac matrices, noting parity conservation, and using the free equation of motion for the quark field. (Chirality forbids $\langle \bar{q}\psi q \rangle$ to appear in the vector-vector correlation function.)

The two-gluon operators G^2 and Θ^g arise only from the first term in Eq. (3.2). To get their coefficients we have to work out the quark propagator up to two gluons. The general fourth rank tensor with two gluon fields may be projected on to the scalar operators by

$$\langle \operatorname{tr}^{c} G_{\alpha\beta} G_{\lambda\sigma} \rangle = \frac{1}{6} (g_{\alpha\lambda} g_{\beta\sigma} - g_{\alpha\sigma} g_{\beta\lambda}) \left(\frac{1}{4} \langle G^{c}_{\mu\nu} G^{\mu\nu c} \rangle + \langle \Theta^{g} \rangle \right) - \frac{1}{3} (u_{\alpha} u_{\lambda} g_{\beta\sigma} - u_{\alpha} u_{\sigma} g_{\beta\lambda} - u_{\beta} u_{\lambda} g_{\alpha\sigma} + u_{\beta} u_{\sigma} g_{\alpha\lambda}) \langle \Theta^{g} \rangle,$$

$$(3.8)$$

 tr^c indicating trace over the color matrices. It turns out that if we insert this expression into the two gluon terms present in S, they do not contribute to G^2 . Thus the calculation of its coefficient reduces to inserting S up to the one-gluon term

$$S(x,0) = S_0(x,0) - \frac{ig_s \gamma^{\mu} \not{x} \gamma^{\nu} G_{\mu\nu}}{16\pi^2 (x^2 - i\epsilon)},$$
(3.9)

in the first term of Eq. (3.2) and then using Eq. (3.8). However the two-gluon terms in S do contribute to Θ^g . But it is not necessary to calculate this coefficient. As we show below, the renormalization group improvement automatically brings it in.

Having indicated the procedure to calculate the coefficients in coordinate space, we take the Fourier transform and write the result in momentum space $(Q^2 \equiv -q^2)$,

$$i \int d^4 x e^{iq \cdot x} T\left\{ V^a_\mu(x) V^b_\nu(0) \right\} = \delta^{ab} \left[Q_{\mu\nu} \left\{ -\frac{1}{8\pi^2} \ln(Q^2/\mu^2) \mathbf{1} + \frac{1}{Q^4} \left(\frac{1}{2} \hat{m} \bar{q} q + \frac{1}{24} G^2 + \frac{2}{3} \Theta^q \right) \right\} + P_{\mu\nu} \left\{ \frac{Q^2}{8\pi^2} \ln(Q^2/\mu^2) \mathbf{1} - \frac{1}{Q^2} \left(\frac{1}{2} \hat{m} \bar{q} q + \frac{1}{24} G^2 + \frac{2}{3} \left(1 - \frac{2\bar{q}^2}{Q^2} \right) \Theta^q \right) \right\} \right].$$
(3.10)

Here μ is the renormalization scale, which we take to be 1 GeV as the natural scale to normalize the operators. But this scale is not convenient for the coefficients. These, calculated above only to the lowest order in α_s , may contain powers of $\ln(Q^2/\mu^2)$ in higher orders. To get rid of these large logarithms, ² we must shift the renormalization point for the coefficients from μ^2 to Q^2 .

Now the left hand side of Eq. (3.10) is independent of the renormalization scale. So are all the operators $\mathbf{1}, \hat{m}\bar{q}q, G^2$ and hence their coefficients. But it is not so for Θ^q and $\Theta^{g,3}$ Indeed, it is well-known in the context of deep inelastic scattering that Θ^q and Θ^g mix under a change of scale [18]. Let us denote the contribution of these operators generally as

$$C_q \mid_{\mu} \Theta^q \mid_{\mu} + C_g \mid_{\mu} \Theta^g \mid_{\mu} \equiv \sum_i C_i \mid_{\mu} \Theta^i \mid_{\mu},$$
(3.11)

where only the dependence on the scale μ is shown explicitly and the index *i* runs over *q* and *g*. The coefficients C_i satisfy a coupled set of renormalization group equations having the solution

$$C_{i}|_{\mu} = C_{j}|_{Q} \left(e^{-\int_{0}^{t} \gamma(t')dt'} \right)_{ji}, \qquad (3.12)$$

where $t = \frac{1}{2} \ln(Q^2/\mu^2)$ and γ is the 2 \otimes 2 anomalous dimension matrix. In the basis we are using it is

$$\gamma = -\frac{g_s^2}{(4\pi)^2} \begin{pmatrix} -\frac{64}{9} & \frac{4}{3}n_f \\ \frac{64}{9} & -\frac{4}{3}n_f \end{pmatrix},$$
(3.13)

where $n_f(=2)$ is the number of flavors. The exponentiation in Eq. (3.12) can be easily worked out by a spectral decomposition of the matrix $\gamma = \sum_i \lambda_i P_i$, where λ_i and P_i are the eigenvalues and the corresponding projection operators [19]. The eigenvalues are

$$\lambda_1 = 0, \quad \lambda_2 = \frac{g_s^2}{(4\pi)^2} \frac{4}{3} \left(\frac{16}{3} + n_f \right), \tag{3.14}$$

²The logarithm in the coefficient of the unit operator in Eq. (3.10) is of different origin. See the first footnote in Sec. 4.6 of Ref. [1].

³If we had calculated the coefficient of Θ^{g} , we would have found it to be proportional to $\alpha_{s} \ln(Q^{2}/\mu^{2})$. It signals mixing with other operators.

and the projection operators can be built out of the (right) eigenvectors and left eigenvectors. We thus finally get

$$\sum_{i} C_{i} \mid_{\mu} \Theta_{i} \mid_{\mu} = \frac{1}{\left(\frac{16}{3} + n_{f}\right)} \left\{ \left(n_{f} C_{q} + \frac{16}{3} C_{g} \right) \Big|_{Q} \Theta + \lambda(Q^{2}) (C_{q} - C_{g}) \mid_{Q} \left(\frac{16}{3} \Theta^{q} - n_{f} \Theta^{g} \right) \Big|_{Q} \right\},$$
(3.15)

where $\Theta = \Theta^q + \Theta^g$ and the anomalous dimension of the second operator resides in $\lambda(Q^2)$,

$$\lambda(Q^2) = \left(\frac{\alpha_s(\mu^2)}{\alpha_s(Q^2)}\right)^{-d/2b}, \qquad d = \frac{4}{3}\left(\frac{16}{3} + n_f\right), \qquad b = 11 - \frac{2}{3}n_f. \tag{3.16}$$

We will use $\alpha_s(\mu^2 = 1 \text{ GeV}^2) = 0.4$ in our numerical evaluations below.

Once the renormalization group improvement is made, we can replace the coefficient functions by their lowest order values. These may be read off from Eq. (3.10) for the amplitude T_l , for example, as $C_f \mid_Q = 2/(3Q^4)$, $C_g \mid_Q = 0$. Thus the renormalization group improved result for T_l reads (with $n_f = 2$)

$$T_l = -\frac{1}{8\pi^2} \ln(Q^2/\mu^2) + \frac{1}{Q^4} \left(\frac{1}{2} \hat{m} \langle \bar{q}q \rangle + \frac{\langle G^2 \rangle}{24} + \frac{2}{11} \left\{ \langle \Theta \rangle + \lambda(Q^2) \left\langle \frac{8}{3} \Theta^q - \Theta^g \right\rangle \right\} \right). \tag{3.17}$$

Notice that if $\lambda(Q^2)$ is set equal to unity, it collapses to the previous result in Eq. (3.10). The amplitude T_t is also given by the same expression, except for an overall factor of $-Q^2$ and a factor of $(1 - 2\bar{q}^2/Q^2)$ multiplying the term with Θ 's.

Let us summarize the effect of mixing of Θ^q and Θ^g under the renormalization group. It introduces the multiplicative factor, $\lambda(Q^2)$, which is nonperturbative in the sense that it is a sum over all the leading logarithms in the perturbation expansion. It is true that the coefficient of Θ^g is smaller than Θ^q by a factor of g_s^2 . We have left out this smaller coefficient, as did the earlier authors, but retained the nonperturbative factor. Numerically the factor is significant for Q^2 away from $\mu^2 = 1 \text{ GeV}^2$.

At dimension 6 we include only the Lorentz scalar operators in vacuum, namely the four-quark operators [1]. In the approximation of ground state saturation, its contribution reduces to the familiar expression involving $\langle \bar{q}q \rangle^2$.

IV. SPECTRAL REPRESENTATION

QCD sum rules make contact with the physical states through the spectral representation of the two point function. For the vacuum amplitudes, these are the Källen-Lehmann representations in q^2 , the only variable available there. At finite temperature and chemical potential, it is convenient to use the Landau representation, which is a spectral decomposition in q_0 at fixed $|\vec{q}|$ [20]. For the invariant amplitudes they are

$$T_{l,t}(q_0, |\vec{q}|) = \frac{1}{\pi} \int_{-\infty}^{+\infty} dq'_0 \frac{\mathrm{Im}T_{l,t}(q'_0, |\vec{q}|)}{q'_0 - q_0 - i\epsilon} \tanh(\beta q'_0/2),$$
(4.1)

up to subtraction terms. The imaginary parts from different intermediate states can be written as phase space integrals over the matrix elements of currents. But we shall not write these expressions directly. Instead, we first calculate the full amplitudes with these intermediate states in the real time formulation of field theory at finite temperature and density [21] and then read off their imaginary parts. Since we have to calculate the amplitudes at the tree level, only the 11-component of the propagators will be needed.

We first consider the ρ -meson pole, see Fig. 1(a), which would dominate the absorptive part of the vacuum correlation function.



FIG. 1. Contributions to the spectral function from (a) the ρ meson and (b) the $N\bar{N}$ state.

In the vacuum, its coupling to the current $V^a_{\mu}(x)$ is given by

$$\langle 0|V^a_\mu(0)|\rho^b\rangle = \delta^{ab}F_\rho m_\rho \epsilon_\mu, \qquad (4.2)$$

where ϵ_{μ} and m_{ρ} are the polarization vector and the mass of the ρ . Experimentally $F_{\rho} = 153.5$ MeV. This matrix element suggests that, as far as the ρ -meson contribution is concerned, we may write the operator relation

$$V^{a}_{\mu}(x) = m_{\rho} F_{\rho} \rho^{a}_{\mu}(x). \tag{4.3}$$

In the medium such a relation is generally modified in two ways. First, the parameters m_{ρ} and F_{ρ} must be replaced by their effective values in the medium, to be denoted below by a star. Secondly, the space and the time components have different constants of proportionality. However, the latter modification takes place in higher orders in density and temperature [22]. As we work to the lowest order, the operator relation (4.3) changes in the medium simply to

$$V^{a}_{\mu}(x) = m^{\star}_{\rho} F^{\star}_{\rho} \rho^{a}_{\mu}(x).$$
(4.4)

Then the ρ -meson contribution to the correlation function is given essentially by its propagator in the medium. As before, we get

$$\begin{pmatrix} \operatorname{Im} T_l(q_0, |\vec{q}|^2) \\ \operatorname{Im} T_t(q_0, |\vec{q}|^2) \end{pmatrix} = \begin{pmatrix} 1 \\ m_{\rho}^{\star 2} \end{pmatrix} \pi F_{\rho}^{\star 2} \delta\left(q_0^2 - (|\vec{q}|^2 + m_{\rho}^{\star 2})\right) \coth(\beta q_0/2).$$

$$(4.5)$$

In the medium there are other equally important contributions. These are from the 2π and, if the medium is nuclear, also the $N\bar{N}$ intermediate state, see Fig. 1(b). As we restrict in this work to sum rules at finite nuclear density but at zero temperature, the 2π contribution would be small and close to its vacuum value. The $N\bar{N}$ contribution in vacuum would also be small, beginning with a cut at the threshold $q_0^2 = 4m_N^2 + |\vec{q}|^2$. However, in the nuclear medium the current also interacts with the real nucleons to give rise to a short cut around the origin, $-|\vec{q}| < q_0 < +|\vec{q}|$. Below we evaluate this contribution.

The contribution of the nucleon isodoublet, spinor field $N_A^i(x)$, i = 1, 2, A = 1, ..., 4, to the vector current is

$$V^{a}_{\mu}(x) = \bar{N}(x)\gamma_{\mu}\frac{\tau^{a}}{2}N(x).$$
(4.6)

The $N\bar{N}$ contribution to the two point function is then

$$\mathcal{T}^{ab}_{\mu\nu}(q) = \frac{i}{2} \delta^{ab} \int d^4x e^{iq \cdot x} \mathrm{tr}[\gamma_{\mu} \mathcal{S}(x) \gamma_{\nu} \mathcal{S}(-x)], \qquad (4.7)$$

where the trace is over the gamma matrices and $i\mathcal{S}(x)$ is the 11-component of the nucleon propagator in the medium $\langle T\left\{N_A^i(x)\bar{N}_B^j(y)\right\}\rangle = i\delta^{ij}\mathcal{S}_{AB}(x-y)$, with

$$i\mathcal{S}(x-y) = \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot (x-y)} (\not\!\!\!k + m_N) \left\{ \frac{i}{k^2 - m_N^2 + i\epsilon} - 2\pi\delta(k^2 - m_N^2) \{n^-\theta(k_0) + n^+\theta(-k_0)\} \right\},\tag{4.8}$$

the distribution functions n^{\mp} for the nucleon and the anti-nucleon being given by $n^{\mp}(\omega) = (e^{\beta(\omega \mp \mu)} + 1)^{-1}$ with $\omega = \sqrt{\vec{k}^2 + m_N^2}$.

Our aim is to integrate out the time component of the loop momentum, when the imaginary part of $\mathcal{T}_{\mu\nu}^{ab}(q)$ can be read off from the resulting expression. However, the above expression for the propagator is not convenient for this purpose because of the presence of $\theta(\pm k_0)$. So we carry out the k_0 integration in the propagator itself, getting

$$i\mathcal{S}(x) = \int \frac{d^3k}{(2\pi)^3 2k_0} \left[\theta(x^0) \left\{ (\not\!\!k + m_N)(1 - n^-)e^{-ik\cdot x} - (-\not\!\!k + m_N)n^+ e^{ik\cdot x} \right\} + \theta(-x^0) \left\{ -(\not\!\!k + m_N)n^- e^{-ik\cdot x} + (-\not\!\!k + m_N)(1 - n^+)e^{ik\cdot x} \right\} \right].$$
(4.9)

Here the time component of k_{μ} is understood to be given by ω . We can now integrate over x^{μ} in Eq. (4.7) to get δ -functions in three-momentum and energy denominators in the time components. After some algebra we get for the imaginary part

$$\operatorname{Im}\mathcal{T}^{ab}_{\mu\nu}(q) = -\frac{\pi}{2}\delta^{ab}\operatorname{coth}(\beta q_0/2)\{I_{\mu\nu}(q) + I_{\mu\nu}(-q)\},\tag{4.10}$$

where

$$I_{\mu\nu}(q) = \int \frac{d^3k}{(2\pi)^3} \frac{\Lambda_{\mu\nu}}{4\omega_1\omega_2} [(1 - n_1^- - n_2^+)\delta(q_0 - \omega_1 - \omega_2) - (n_2^- - n_1^-)\delta(q_0 - \omega_1 + \omega_2)], \qquad (4.11)$$

with

$$\Lambda_{\mu\nu} \equiv \operatorname{tr}\{\gamma_{\mu}(\not\!\!k + m_{N})\gamma_{\nu}(\not\!\!k - \not\!\!q + m_{N})\},\$$

= 4[2k_{\mu}k_{\nu} - (k_{\mu}q_{\nu} + k_{\nu}q_{\mu}) - g_{\mu\nu}\{k \cdot (k - q) - m_{N}^{2}\}]. (4.12)

Here the subscripts on n^{\mp} serve to indicate that their arguments are $\omega_1 = \sqrt{\vec{k}^2 + m_N^2}$ and $\omega_2 = \sqrt{(\vec{k} - \vec{q})^2 + m_N^2}$, respectively. The first and the second terms in Eq. (4.11) are nonvanishing in the timelike $(q^2 > 4m_N^2)$ and spacelike $(q^2 < 0)$ regions, respectively. Working out the angular integration, it becomes

$$I_{\mu\nu}(q) = \frac{1}{4|\vec{q}|} \int_{\omega_{-}}^{\omega_{+}} \frac{d\omega_{1}}{(2\pi)^{2}} \Lambda_{\mu\nu} (1 - n_{1}^{-} - n_{2}^{+}) \theta(q^{2} - 4m_{N}^{2}) - \frac{1}{4|\vec{q}|} \int_{\omega_{+}}^{\infty} \frac{d\omega_{1}}{(2\pi)^{2}} \Lambda_{\mu\nu} (n_{2}^{-} - n_{1}^{-}) \theta(-q^{2}),$$
(4.13)

where the integration limits are given by $\omega_{\pm} = [q_0 \pm |\vec{q}| v(q^2)]/2$, with $v(q^2) = \sqrt{1 - 4m_N^2/q^2}$.

To project the tensor $I_{\mu\nu}$ on the imaginary parts of the invariant amplitudes it is convenient to form the scalars I^{μ}_{μ} and $u^{\mu}I_{\mu\nu}u^{\nu}$ and relate them to the invariant amplitudes. Changing the variable ω_1 to x given by $\omega_1 = \frac{1}{2}(q_0 + |\vec{q}|x)$, we finally get for the contribution from the $N\bar{N}$ state

$$\begin{pmatrix} \operatorname{Im}T_{l}(q_{0}, |\vec{q}|^{2}) \\ \operatorname{Im}T_{t}(q_{0}, |\vec{q}|^{2}) \end{pmatrix} = -\frac{\pi}{2} \operatorname{coth}(\beta q_{0}/2) \left[-\frac{v(3-v^{2})}{24\pi^{2}} \begin{pmatrix} 1 \\ q^{2} \end{pmatrix} + \begin{pmatrix} I_{l}^{+} \\ I_{t}^{+} \end{pmatrix} \right], \quad \text{for } q^{2} > 4m_{N}^{2},$$

$$(4.14)$$

with

$$\begin{pmatrix} I_l^+ \\ I_t^+ \end{pmatrix} = \frac{-1}{32\pi^2} \int_{-v}^{v} dx \begin{pmatrix} 2(x^2 - 1) \\ -q^2(2 - v^2 + x^2) \end{pmatrix} \{ n((q_0 + |\vec{q}|x)/2 - \mu) - n((q_0 - |\vec{q}|x)/2 + \mu) \},$$
(4.15)

and

$$\begin{pmatrix} \operatorname{Im} T_l(q_0, |\vec{q}|^2) \\ \operatorname{Im} T_t(q_0, |\vec{q}|^2) \end{pmatrix} = -\frac{\pi}{2} \operatorname{coth}(\beta q_0/2) \begin{pmatrix} I_l \\ I_t^- \end{pmatrix}, \quad \text{for } q^2 \le 0,$$

$$(4.16)$$

with

$$\begin{pmatrix} I_l^-\\ I_t^- \end{pmatrix} = \frac{-1}{32\pi^2} \int_v^\infty dx \begin{pmatrix} 2(x^2 - 1)\\ -q^2(2 - v^2 + x^2) \end{pmatrix} \{ n((-q_0 + |\vec{q}|x)/2 - \mu) - n((q_0 + |\vec{q}|x)/2 - \mu) \}.$$
(4.17)

The superscripts (\pm) on $I_{l,t}$ denote timelike and spacelike q_{μ} , respectively. The distribution function n appearing in Eqs. (4.15) and (4.17) is defined by $n(\varepsilon) = (e^{\beta \varepsilon} + 1)^{-1}$.

Still higher continuum contributions, at least their leading parts, are density independent, as is shown by the coefficient of the unit operator. They will drop out in our subtraction process.

V. SUM RULES

We now go to the spacelike region in q_{μ} , $(Q_0^2 = -q_0^2 > 0)$ at fixed $|\vec{q}|$ and take the Borel transform of both the spectral representation and the operator product expansion with respect to Q_0^2 . The sum rules are obtained by equating these transforms at sufficiently high M^2 . For T_l it is

$$F_{\rho}^{\star 2} e^{-m_{\rho}^{\star 2}/M^{2}} + \frac{1}{48\pi^{2}} \int_{4m_{N}^{2}}^{\infty} dq_{0}^{2} e^{-q_{0}^{2}/M^{2}} v^{3}(q_{0}^{2} + |\vec{q}|^{2}) + e^{|\vec{q}|^{2}/M^{2}} \left(\int_{4m_{N}^{2} + |\vec{q}|^{2}}^{\infty} dq_{0}^{2} e^{-q_{0}^{2}/M^{2}} I_{l}^{+}(q_{0}, |\vec{q}|) + \int_{0}^{|\vec{q}|^{2}} dq_{0}^{2} e^{-q_{0}^{2}/M^{2}} I_{l}^{-}(q_{0}, |\vec{q}|) \right) = \frac{M^{2}}{8\pi^{2}} + \frac{\langle O \rangle}{M^{2}} - \frac{\langle O' \rangle}{2M^{4}},$$
(5.1)

where $\langle O \rangle$ and $\langle O' \rangle$ represent the contributions of all dimension 4 operators and of dimension 6, scalar operators in vacuum, respectively,

$$\langle O \rangle = \frac{1}{2} \hat{m} \langle \bar{q}q \rangle + \frac{\langle G^2 \rangle}{24} + \frac{2}{11} \left\{ \langle \Theta \rangle + \lambda (M^2) \left\langle \frac{8}{3} \Theta^q - \Theta^g \right\rangle \right\}, \quad \langle O' \rangle = \frac{7g_s^2}{81} \langle \bar{q}q \rangle^2.$$
(5.2)

The amplitude T_t satisfies a similar sum rule.

=

Considerable simplification results if we take $|\vec{q}| \rightarrow 0$. This limit may be taken inside the integrals in Eq. (5.1), except for the last one, where the integrand diverges, while the range of integration shrinks to zero. A careful evaluation gives a result resembling a pole at $q^2 = 0.4$ Then the above sum rule for T_l becomes

$$F_{\rho}^{\star 2} e^{-m_{\rho}^{\star 2}/M^{2}} + \frac{1}{48\pi^{2}} \int_{4m_{N}^{2}}^{\infty} dq_{0}^{2} v_{0}(3-v_{0}^{2}) \left[n(q_{0}/2-\mu) + e^{-q_{0}^{2}/M^{2}} \{1-n(q_{0}/2-\mu) - n(q_{0}/2+\mu)\} \right]$$

$$= \frac{M^{2}}{8\pi^{2}} + \frac{\langle O \rangle}{M^{2}} - \frac{\langle O' \rangle}{2M^{4}},$$
(5.3)

while the sum rule for T_t in this limit may similarly be found to be

$$m_{\rho}^{\star 2} F_{\rho}^{\star 2} e^{-m_{\rho}^{\star 2}/M^{2}} + \frac{1}{48\pi^{2}} \int_{4m_{N}^{2}}^{\infty} dq_{0}^{2} q_{0}^{2} v_{0} (3 - v_{0}^{2}) e^{-q_{0}^{2}/M^{2}} \{1 - n(q_{0}/2 - \mu) - n(q_{0}/2 + \mu)\}$$

$$= \frac{M^{4}}{8\pi^{2}} - \langle O \rangle + \frac{\langle O' \rangle}{M^{2}}, \qquad (5.4)$$

where $v_0 = (1 - 4m_N^2/q_0^2)^{1/2}$. Note that the contribution of the short cut is nonvanishing in this limit only in T_l .

As they stand, the above sum rules cannot be well saturated, since the 2π and the continuum contributions on the spectral side are missing. However, if we subtract the corresponding vacuum sum rules, the continuum contribution will practically drop out. We now go to the limit of zero temperature, when the 2π contribution reduces to zero. With $\mu > 0$, the nucleon and the antinucleon distribution functions reduce in this limit to a θ -function and zero, respectively. We thus finally arrive at the following sum rules at finite nuclear density:

$$\overline{F_{\rho}^{\star 2} e^{-m_{\rho}^{\star 2}/M^{2}}} + \frac{1}{24\pi^{2}} \int_{4m_{N}^{2}}^{4\mu^{2}} ds (1 - e^{-s/M^{2}}) \sqrt{1 - 4m_{N}^{2}/s} (1 + 2m_{N}^{2}/s) = \frac{\overline{\langle O \rangle}}{M^{2}} - \frac{\overline{\langle O' \rangle}}{2M^{4}}, \tag{5.5}$$

$$\overline{m_{\rho}^{\star 2} F_{\rho}^{\star 2} e^{-m_{\rho}^{\star 2}/M^{2}}} - \frac{1}{24\pi^{2}} \int_{4m_{N}^{2}}^{4\mu^{2}} ds \, s \, e^{-s/M^{2}} \sqrt{1 - 4m_{N}^{2}/s} (1 + 2m_{N}^{2}/s) = -\overline{\langle O \rangle} + \frac{\overline{\langle O' \rangle}}{M^{2}}, \tag{5.6}$$

where the bar over a quantity denotes subtraction of its vacuum value, e.g., $\overline{\langle O \rangle} = \langle O \rangle - \langle 0 | O | 0 \rangle$.

The spectral integrals in the sum rules can be expanded in powers of the Fermi momentum $p_F = \sqrt{\mu^2 - m_N^2}$, or the nucleon number density [4]

$$\bar{n} = 4 \int \frac{d^3 p}{(2\pi)^3} \theta(p_F - |\vec{p}|) = \frac{2p_F^3}{3\pi^2}.$$
(5.7)

The operator matrix elements can also be expanded in powers of the nucleon number density. With the normalization of the one nucleon state $\langle p, \alpha | p', \alpha' \rangle = (2\pi)^3 2p_0 \delta_{\alpha,\alpha'} \delta^3(\vec{p} - \vec{p'}), \alpha, \alpha'$ denoting the spin and isospin degrees of freedom, we have the expansion for any operator R up to first order as

⁴The situation is similar to that at finite temperature [2]. See also Ref. [23].

$$\langle R \rangle = \langle 0|R|0\rangle + \int \frac{d^3p}{(2\pi)^3 2p_0} \sum_{\alpha} \langle p, \alpha|R|p, \alpha\rangle \theta(p_F - |\vec{p}|).$$
(5.8)

Denoting the constant, averaged matrix element by $\langle p|R|p\rangle$, we get

$$\langle R \rangle = \langle 0|R|0 \rangle + \frac{\langle p|R|p \rangle}{2m_N} \bar{n}.$$
(5.9)

Thus

$$\langle \hat{m}\bar{q}q \rangle = \langle 0|\hat{m}\bar{q}q|0\rangle + \sigma\bar{n},\tag{5.10}$$

where the σ matrix element is defined by $\sigma = \langle p | \hat{m}(\bar{u}u + \bar{d}d) | p \rangle / (2m_N) \simeq 45$ MeV [24]. The two-gluon operator is determined by the trace anomaly

$$\Theta^{\mu}_{\mu} = -\frac{9}{8} \frac{\alpha_s}{\pi} G_{\mu\nu} G^{\mu\nu} + \hat{m}\bar{q}q.$$
 (5.11)

We then get

$$\langle G^2 \rangle = \langle 0 | G^2 | 0 \rangle - \frac{8}{9} (m_N - \sigma) \bar{n}.$$
(5.12)

To determine the quark and the gluon energy densities in the medium, we need their nucleon matrix elements. More generally one may write the traceless $\Theta_{\mu\nu}^{q,g}$ as

$$\langle p|\Theta_{\mu\nu}^{q,g}|p\rangle = 2A^{q,g}\left(p_{\mu}p_{\nu} - \frac{1}{4}g_{\mu\nu}p^{2}\right).$$
(5.13)

The factor 2 ensures that the constants satisfy $\sum_f A^f + A^g = 1$. These constants individually may be obtained from fits with the data on deep inelastic scattering. Based on the parametrizations for the quark and the gluon distribution functions in Ref. [25], these are found at $Q^2 = 1 \text{ GeV}^2$ to be

$$A^q = 0.62, \qquad A^g = 0.35. \tag{5.14}$$

Collecting the above results we get

$$\overline{\langle O \rangle} = C\bar{n}, \quad C = \frac{\sigma}{2} - \frac{1}{27}(m_N - \sigma) + \frac{3}{22}m_N \left\{ A^q + A^g + \lambda(M^2) \left(\frac{8}{3}A^q - A^g\right) \right\}, \tag{5.15}$$

$$\overline{\langle O' \rangle} = D\bar{n}, \quad D = \frac{56}{81} \pi \alpha_s \langle 0 | \bar{q}q | 0 \rangle \frac{\sigma}{\hat{m}}, \tag{5.16}$$

where the quark condensate in vacuum is given by $\langle 0|\bar{q}q|0\rangle/2 = \langle 0|\bar{u}u|0\rangle = -(225 \text{ MeV})^3$. Furthermore, we will use $\hat{m} = 7 \text{ MeV}$ below.

The sum rules now determine the nuclear density dependence of the parameters of the ρ meson. To first order in the number density, we may expand these as

$$m_{\rho}^{\star} = m_{\rho} \left(1 + a \frac{\bar{n}}{\bar{n}_s} \right), \qquad F_{\rho}^{\star} = F_{\rho} \left(1 + b \frac{\bar{n}}{\bar{n}_s} \right), \tag{5.17}$$

where \bar{n}_s is the saturation nucleon number density, $\bar{n}_s = (110 \text{ MeV})^3$. From the sum rules (5.5) and (5.6) we then obtain the following expressions for the coefficients a and b:

$$a = -\frac{\bar{n}_s}{2F_{\rho}^2} e^{m_{\rho}^2/M^2} \left[C\left(\frac{1}{m_{\rho}^2} + \frac{1}{M^2}\right) - \frac{D}{M^2} \left(\frac{1}{m_{\rho}^2} + \frac{1}{2M^2}\right) - \frac{1}{4m_N} - \left(\frac{m_N}{m_{\rho}^2} - \frac{1}{4m_N}\right) e^{-4m_N^2/M^2} \right],\tag{5.18}$$

$$b = -\frac{\bar{n}_s}{2F_{\rho}^2} e^{m_{\rho}^2/M^2} \left[C\frac{m_{\rho}^2}{M^4} - \frac{D}{2M^4} \left(1 + \frac{m_{\rho}^2}{M^2} \right) + \frac{1}{4m_N} \left(1 - \frac{m_{\rho}^2}{M^2} \right) - \left(\frac{m_N}{M^2} + \frac{1}{4m_N} \left(1 - \frac{m_{\rho}^2}{M^2} \right) \right) e^{-4m_N^2/M^2} \right].$$
(5.19)



FIG. 2. Coefficients of \bar{n}/\bar{n}_s in the expansion of m_{ρ}^* and F_{ρ}^* from Eq. (5.17) as function of the Borel parameter M.

A numerical evaluation of a and b is shown in Fig. 2. It will be noted that there is no sign of constancy of a and b in any region of M^2 ; rather, they increase rapidly and monotonically throughout, in particular a. We conclude that the sum rules as saturated above are inconsistent and cannot give any reliable information about the density dependence of m_{ρ} and F_{ρ} .

VI. DISCUSSION

We have studied QCD sum rules in the nuclear medium in the ρ -meson channel using a saturation scheme considered earlier by a number of authors [6–8]. The sum rules we actually evaluate consist of terms, all proportional to the nuclear density. We find that the results are not stable in any range of variation of the Borel parameter. We conclude that the sum rules with the present saturation scheme become inconsistent, in apparent contradiction with the evaluation by these authors.

The resolution of this contradiction lies in the fact that the "full" sum rules considered by these authors are dominated by vacuum contributions, the density dependent terms serving as small perturbations. As stated already in the Introduction, their stability follows essentially from that of the vacuum sum rules. The variation of some free parameters such as s_0 , the beginning of the continuum, in these works would further contribute to this apparent stability.

It would appear from earlier works that QCD sum rules can accommodate different saturation schemes, each one giving rise to a new set of results [11]. By working with a more sensitive version of the sum rules, we show that the situation may not actually be so: a simple saturation scheme, such as here, is rejected and a more realistic one is called for.

Let us finally discuss the neglected contributions, whose inclusion should restore the stability of these sum rules. At dimension 6, five more operators would have contributed. These contributions can be evaluated using the data on deep inelastic scattering [26]. To get an idea of the convergence of the series of operators, we note that the contributions of a family of similar operators, differing only in the number of Lorentz indices, form a series in powers of (m_N/M) with decreasing coefficients, given by the corresponding moments of the structure functions.

In the spectral function, it is not just the NN state, which can give rise to the Landau cut in the low q^2 region. Any other state $N\bar{N}^*$, where N^* is a resonance communicating with the ρN channel, can contribute a cut for $q^2 \leq (m_{N^*} - m_N)^2$. The two resonances $\Delta(1232)$ and N(1440) would presumably contribute significantly. To include these contributions we need, however, know the current- $N\bar{N}^*$ couplings. Also the increased width of the ρ meson in the medium is another source of altered contributions to the sum rules. We note that a study of such "subtracted" sum rules for ρ mesons at finite temperature showed no such instabilities as mentioned above [27].

ACKNOWLEDGMENTS

We thank H. Leutwyler and P. Minkowski for helpful discussions. We also acknowledge the kind hospitality at the Institute for Theoretical Physics, University of Berne, where we started this work. This work was supported in part by Schweizerischer Nationalfonds.

- M.A. Shifman, A.I. Vainshtein, and V.I. Zakharov, Nucl. Phys. B147, 385 (1979); B147, 448 (1979); for a collection of original papers and comments, see *Vacuum Structure and QCD Sum Rules*, edited by M.A. Shifman (North Holland, Amsterdam, 1992); see also S. Narison, *QCD Spectral Sum Rules* (World Scientific, Singapore, 1989).
- [2] A.I. Bochkarev and M.E. Shaposhnikov, Nucl. Phys. **B268**, 220 (1986).
- [3] E.G. Drukarev and E.M. Levin, JETP Lett. 48, 338 (1988).
- [4] X. Jin, T.D. Cohen, R.J. Furnstahl, and D.K. Griegel, Phys. Rev. C 47, 2882 (1993).
- [5] E.V. Shuryak, Rev. Mod. Phys. 65, 1, (1993).
- [6] T. Hatsuda and S.H. Lee, Phys. Rev. C 46, R34 (1992); T. Hatsuda, S.H. Lee, and H. Shiomi, *ibid.* 52, 3364 (1995).
- [7] X. Jin and D.B. Leinweber, Phys. Rev. C 52, 3344 (1995).
- [8] T.D. Cohen, R.D. Furnstahl, D.K. Griegel, and X. Jin, Prog. Part. Nucl. Phys. 35, 221 (1995).
- [9] A. Nyffeler, Centre de Physique Théorique Marseille Report No. CPT-2000/P.4071, presented at the Conference on Strong and Electroweak Matter, Marseille, 2000, hep-ph/0010329 (to appear in the proceedings).
- [10] F. Klingl, N. Kaiser, and W. Weise, Nucl. Phys. A624, 527 (1997).
- [11] S. Leupold, W. Peters, and U. Mosel, Nucl. Phys. A628, 311 (1998).
- [12] H.A. Weldon, Phys. Rev. D 26, 1394 (1982)
- [13] J.I. Kapusta and E.V. Shuryak, Phys. Rev. D 49, 4694 (1994).
- [14] H. Fritzsch and H. Leutwyler, Phys. Rev. D 10, 1624 (1974).
- [15] W. Hubschmid and S. Mallik, Nucl. Phys. **B207**, 29 (1982).
- [16] S. Mallik, Phys. Lett. B **416**, 373 (1998).
- [17] For an earlier summary of the coordinate method, see V.A. Novikov, M.A. Shifman, A.T. Vainshtein, and V. Zakharov, Fortschr. Phys. 32, 600 (1984).
- [18] See, for example, M.E. Peskin and D.V. Schroeder, An Introduction to Quantum Field Theory (Addison-Wesley, Reading, MA, 1995).
- [19] D.J. Gross, in Methods of Field theory, edited by R. Balian and J. Zinn-Justin (North Holland, Amsterdam, 1976).
- [20] L.D. Landau, Sov. Phys. JETP 7, 182 (1958), reproduced in *Collected Papers of L.D. Landau*, edited by D. Ter Haar (Pergamon, Oxford, 1965).
- [21] A.J. Niemi and G.W. Semenoff, Ann. Phys. (N.Y.) 152, 105 (1984); see also N.P. Landsman and Ch.G.van Weert, Phys. Rep. 145, 141 (1987).
- [22] H. Leutwyler, Phys. Rev. D 49, 3033 (1994); R.D. Pisarski and M. Tytgat, *ibid.* 54, 2989 (1996); D. Toublan, *ibid.* 56, 5629 (1997).
- [23] S. Mallik and K. Mukherjee, Phys. Rev. D 58, 096011 (1998).
- [24] J. Gasser, H. Leutwyler, and M.E. Sainio, Phys. Lett. B 253, 252 (1991).
- [25] A.D. Martin, R.G. Roberts, W.J. Stirling, and R.S. Thorne, Eur. Phys. J. C 4, 463 (1998).
- [26] S. Leupold and U. Mosel, Phys. Rev. C 58, 2939 (1998).
- [27] S. Mallik and K. Mukherjee, Phys. Rev. D 61, 116007 (2000).