Non-exponential decays and nonextensivity

G.Wilk^{1*}and Z.Włodarczyk² †

¹The Andrzej Sołtan Institute for Nuclear Studies Hoża 69; 00-689 Warsaw, Poland
²Institute of Physics, Świętokrzyska Academy Konopnickiej 15; 25-405 Kielce, Poland

November 25, 2018

Abstract

We demonstrate that non-exponential decays of unstable systems can be understood as yet another example of nonextensivity encountered in many physical systems and as such can be characterized by the nonextensivity parameter q.

PACS numbers: 05.40.Fb 24.60.-k 05.10.Gg *Keywords:* Nonextensive statistics, Lévy distributions, Thermal models

^{*}e-mail: wilk@fuw.edu.pl

[†]e-mail: wlod@pu.kielce.pl

It is known that for the unstable system, for which the probability γ to decay in a unit of time is constant, its survival time probability is given by simple exponential law

$$P(t) = \exp(-\gamma t). \tag{1}$$

This means that our system is not really aging but with passing time shows its persistency to decay. On the other hand, it is also widely known [1] that spontaneous decays of quantum-mechanical unstable systems cannot be described by the pure exponential law (neither for short nor for long times), i.e., one observes that for them $P(t) > \exp(-\gamma t)$ and, in practice, P(t) exhibits the power-like behaviour of the type [1]¹

$$P(t) \propto t^{-\delta}.$$
 (2)

We would like to concentrate in this letter on the observation that there is enormous variety of different physical phenomena which do exhibit a power-like behaviour instead of naively expected exponential one. They can be described in a most economical and adequate way in terms of Tsallis nonextensive statistics [4] by introducing only one new parameter q, known as the nonextensivity parameter. The common feature of these phenomena is that they are characterized either by long-range interactions and/or long-range microscopic memories or by space-time (and/or phase-space) (multi)fractal structure of the underlying physical process [4], or by intrinsic fluctuations of some parameters present in the system under consideration [5]². In this approach one parameter power-like formula replaces the usual exponential (Boltzmann-Gibbs) one³:

$$L_q(t) = \frac{2-q}{\tau} \left[1 - (1-q)\frac{t}{\tau} \right]^{\frac{1}{1-q}} \quad \stackrel{q \to 1}{\Longrightarrow} \quad L_{q=1}(t) = \frac{1}{\tau} \cdot \exp\left[-\frac{t}{\tau}\right]. \tag{3}$$

The question we would like to address is whether non-exponentiality of decays mentioned above can be expressed in the same approach, by deviation of the suitably defined parametr q from unity.

In [5] we have demonstrated that parameter q (and power-like distribution it characterizes) occurs whenever parameter $1/\tau$ of the original exponential distribution fluctuates according to gamma distribution. In this case q is given by the

¹The ability to observe such nonexponential decays [2] and interest in quantum version of the Zeno paradox [3] resulted in the increase of interest in the problem of unstable quantum systems.

²This last point, which we shall explore here, is receiving growing attention recently [7]. ³We received bills to strong at this point the fact that x_{1} (2) is the speed solution of a simple

³We would like to stress at this point the fact that eq. (3) is the exact solution of a simple nonlinear ordinary differential equation of the type: $\frac{dy}{dt} = \frac{-y^q}{\tau}$ with $y(t) = \frac{L(t)}{L(0)}$, as discussed in detail in cf. [6].

relative variance of such fluctuations. We shall demonstrate now, using random matrix approach [8] that, indeed, the non-exponential decays can emerge in a natural way from the possible fluctuations of parameter $\gamma = 1/\tau$ in exponential distribution (1). To this end let us start with discussion of some consequences of quantum chaos, which emerge from applying random matrix hamiltonians, elements of which follow some random probability distribution [8]. Namely, let us consider situation where we have N intercoupled states with couplings between them described by hamiltonian given in terms of $N \times N$ hermitian (and rotatiotionally and time reversal invariant) matrix (with real matrix elements): $H = [H_{ij}]_{N \times N}$, $(H_{ij} = H_{ji})$. Suppose that one is interested in ensemble of such random matrices with P(H) being a probability of a given matrix H. There is a number of propositions how to define P(H) correspoding to different models of randomness of matrix elements used [9]. However, in the limit of large matrices (large N) which are hermitian and real, all of them lead to results analogous to those obtained by assuming gaussian ensemble⁴. This can be seen as follows. Let P(H) be such that: (i) - elements H_{ij} are uncorrelated, i.e.,

$$P(H) = \prod_{i=1}^{N} \prod_{j \ge 1}^{N} P(H_{ij}),$$
(4)

(*ii*) - distribution (4) is invariant in respect to arbitrary unitary transformation of the basic states (because initial basic states $|i\rangle$ used to define H_{ij} have been chosen in arbitrary way). Let us now look for distribution $P(C_i)$ of the amplitudes C_i for eigenvectors of H assuming that our hermitean ensemble is invariant in respect to the orthogonal change of the basic vectors. Such ensemble is distributed in the probability space with some density $P(C_1, \ldots, C_N)$ with $C_1^2 + C_2^2 + C_3^2 + \cdots + C_N^2 = 1$ and joint distribution function for N amplitudes has the form

$$P(C_1, \dots, C_N) = \frac{2}{\Omega_N} \delta\left(1 - \sum_{i=1}^N C_i^2\right)$$
(5)

(where normalization constant Ω_N is equal to the volume of N-dimensional sphere of unit radius and factor 2 originates from the fact that both positive and negative C_i enter $P(C_1, \ldots, C_N)$). Integrating (5) over all but one coordinates C_i gives the following distribution $P(C_i)$ of amplitude C_i

$$P(C_i) = \frac{\Omega_{N-1}}{\Omega_N} \left(1 - C_i^2 \right)^{\frac{N-3}{2}} = \frac{\Gamma\left(\frac{N+2}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{N+1}{2}\right)} \left(1 - C_i^2 \right)^{\frac{N-3}{2}}, \tag{6}$$

 $^{^{4}}$ The gaussian ensembles were first discussed by E.P.Wigner [10]. For analysis of the gaussian ensemble considered here see [11].

which in the limit of large N has the following form ⁵

$$P(C_i) = \sqrt{\frac{N}{2\pi}} \exp\left(-\frac{N}{2}C_i^2\right) = \frac{1}{\sqrt{2\pi \langle C_i^2 \rangle}} \exp\left(-\frac{C_i^2}{2 \langle C_i^2 \rangle}\right),\tag{7}$$

where

$$\left\langle C_i^2 \right\rangle = \frac{1}{N}$$
 and $\frac{\left\langle (C_i^2)^2 \right\rangle - \left\langle C_i^2 \right\rangle^2}{\left\langle C_i^2 \right\rangle^2} = 2.$ (8)

However, for our purpose we are really interested in decay width γ_i , which is proportional to the probability C_i^2 of finding single component $|i\rangle$ in the composite state. Its distribution is given by

$$P(\gamma_i)d\gamma_i = 2 P(C_i) |dC_i|, \qquad (9)$$

where factor 2 reflects the fact that both positive and negative values of C_i contribute to the same value of γ_i . Because $|dC_i| = \left|\frac{dC_i}{d\gamma_i}\right| d\gamma_i$ one gets following distribution of such widths:

$$P(\gamma_i)d\gamma_i = 2P(C_i) \left| \frac{d\gamma_i}{d\gamma_i/dC_i} \right| = \frac{1}{\sqrt{2\pi\gamma_i < \gamma_i > 2}} \exp\left(-\frac{\gamma_i}{2 < \gamma_i > 2}\right) d\gamma_i \qquad (10)$$

characteristic function of which is $\phi(z) = (1 - i 2 < \gamma_i > z)^{-1/2}$. In the case where contribution to the width γ comes from many channels, distribution we are looking for is obtained by summing all contributions from the particular channels of widths γ_i . For ν different independent channels with the same mean values $\gamma_0 = <\gamma_i >$ the corresponding characteristic function will be $\phi_{\nu}(z) = (1 - i 2\gamma_0 z)^{-\nu/2}$ (with $<\gamma > = \nu\gamma_0$), i.e., the corresponding distribution function will be gamma function of the form

$$P_{\nu}(\gamma)d\gamma = \frac{1}{\Gamma\left(\frac{\nu}{2}\right)} \left(\frac{\nu}{2<\gamma>}\right) \left(\frac{\nu\gamma}{2<\gamma>}\right)^{\frac{\nu}{2}-1} \exp\left(-\frac{\nu\gamma}{2<\gamma>}\right).$$
(11)

This completes our proof that, as a results of composition of contributions from the ν channels, we obtain distribution of the widths of the decays $P_{\nu}(\gamma)$ in the form of gamma distribution, fluctuations of which are characterized by the relative variance of the form

$$\frac{\langle (\gamma - \langle \gamma \rangle)^2 \rangle}{\langle \gamma \rangle^2} = \frac{2}{\nu}$$
(12)

and they decrease with growing ν^{6} .

⁵By using Stirling formula for (N >> 1) leading to $\ln P(C_i) \simeq \frac{1}{2} \ln \frac{N}{2\pi} + \frac{N}{2} \ln (1 - C_i^2)$, and noticing that for large N we have $C_i^2 \ll 1$.

⁶The (11) is for the integer values of ν encoutered here identical to the χ^2 distribution obtained in [7].

We have therefore demonstrated that, indeed (at least in the above example), the fluctuations of the life times $\tau = 1/\gamma$ around mean value $\tau_0 = 1/\langle \gamma \rangle$ are described by gamma distribution of the form (11) of the variable $1/\tau$. According to our philosophy such fluctuations of parameter γ in the exponential distribution (1) lead therefore directly to the Lévy distribution of the form [5, 7]

$$L_q(t,\tau_0) = \int_0^\infty d\left(\frac{t}{\tau}\right) P_\nu\left(\frac{t}{\tau}\right) \exp\left(-\frac{t}{\tau}\right) = \frac{2-q}{\tau_0} \left[1 - (1-q)\frac{t}{\tau_0}\right]^{\frac{1}{1-q}}$$
(13)

with nonextensivity parameter q equal to $q = 1 + \frac{2}{\nu}$.

We shall close with some remarks on chaotic systems and mixing of configurations. Notice first that distribution $P_{\nu}(\gamma)d\gamma$ is known for $\nu = 1$ in the nuclear physics as the so called Porter-Thomas distribution $[12]^7$ and describes in a satisfactory way the widths of resonances observed experimentally [14]. On the other hand the many channel case of $\nu > 2$ corresponds, for example, to the case of total widths for gamma emision or for overbarrier fission. Model using random matrices corresponds to the extreme case of total mixing between allowed degrees of freedom. Such mixing of configurations is realized in the case of compound nucleus (where energy of excitation is divided among many degrees of freedom). It means that component representing one-particle motion in reference to the whole nucleus is decomposed into wave function of a large number of different resonance states. In the Fermi gas model the compound nucleus corresponds to a particularly strong mixing of different configurations, both corresponding to the stationary and to the resonance states. Such emergence of the mixing of configurations, which is leading to the compound nucleus, can be understood as effect of an increase of importance (parallel to the increase of the excitation energy of the system) of small perturbations in the one-particle motion. It happens because in this case the number of exciting particles increases (as increases the number of possible interactions) and excitation energy per single particle increases as well (the density of final states accessible in the collisions increases) 8 .

One should stress also that random matrix approach is applied in the chaotic quantum systems [15] and it seems to be the characteristic feature of such quantum

⁷Deviations from this distribution, which can be attributed to small N involved, have been discussed recently in [13].

⁸In this respect there is a whole list of problems which could possibly be considered, for example, the problem of mixed states in the quantum physics, connection with the composite biochemical systems for which fluorescence time distributions are known to be non-exponential (and are usually described by a composition of a number of exponents).

systems, which in the classical limit show chaotic behaviour. It is connected with the fact that in such systems there is no symmetry, i.e., there is no degeneracy and no selection rules which would exclude interactions of some particular states ⁹.

We are grateful for fruitful discussions with Prof. C. Tsallis. The partial support of Polish Committee for Scientific Research (grants 2P03B 011 18 and 621/E-78/SPUB/CERN/P-03/DZ4/99) is acknowledged.

⁹For application of random matrices to description of chaos in quantum systems see [16]

References

- L.Fonda, G.C.Ghirardi and A.Rimini, Rep. Prog. Phys. 41 (1978) 587;
 C.A.Nicolides and D.R.Beck, Int. J. Quantum Chem. 14 (1978) 457; P.Exner, Open Quantum Systems and Feynman Integrals (Reidel, Dordrecht, 1985);
 A.Sudbery, Ann. Phys. 157 (1984) 512.
- [2] P.T.Greenland, Nature 335 (1988) 298; S.R.Wilkinson et al., Nature 387 (1997) 575.
- [3] J.Maddox, Nature **306** (1983) 111; P.Knight, Nature (1990) 493; T.Petrosky,
 S.Tasaki and I.Prigogine, Phys. Rev. A41 (1990) 2295.
- [4] C.Tsallis, J. Stat. Phys. 52 (1988) 479; cf. also C.Tsallis, cond-mat/0010150 (to appear in Chaos, Solitons and Fractals (2001) and in Nonextensive Statistical Mechanics and its Applications, S.Abe and Y.Okamoto (Eds.), Lecture Notes in Physics LPN560, Springer (2000). For updated bibliography on this subject see http://tsallis.cat.cbpf.br/biblio.htm. Its recent summary is provided in the special issue of Braz. J. Phys. 29 (No 1) (1999) (available also at http://sbf.if.usp.br/ WWW-pages/ Journals/BJP/ Vol29/Num1/index.htm).
- [5] G.Wilk and Z.Włodarczyk, Phys. Rev. Lett. 84 (2000) 2770 and Chaos, Solitons and Fractals 13/3 (2001) 581.
- [6] C.Tsallis, G.Bemski and R.S.Mendes, *Phys. Lett.* A257 (1999) 93.
- [7] C.Beck, Non-additivity of Tsallis entropies and fluctuations of temperature, cond-mat/0105371 and Dynamical foundations of nonextensive statistical mechanics, cond-mat/0105374.
- [8] M.L.Mehta, Random Matrices and the Statistical Theory of Energy Level, Academic Press, New York, 1967; M.Carmeli, Statistical Theory of Random Matrices, Dekker, New York, 1083; M.L.Mehta and A.Pandey, J. Phys. A16 (1983) 2655.
- [9] N.Rosenzweig, *Statistical Physics*, Vol. 3, ed. K.W.Ford, Benjamin, New York.
- [10] E.P.Wigner, Ann. Math. 67 (1958) 325.
- [11] C.E.Porter and N.Rosenzweig, Ann. Acad. Sci. Finland A6 (1960) No. 44.
- [12] C.E.Porter and R.G.Thomas, *Phys. Rev.* **104** (1956) 483.

- [13] M.S.Hussein and M.P.Prato, Phys. Rev. Lett. 84 (2000) 3783.
- [14] J.B.Garg, J.Rainwater, J.S.Petersen and W.W.Havens, Phys. Rev. B134 (1964) 985.
- [15] H.Friedrich and D.Wintgen, Phys. Rep. 183 (1989) 37; T.Guhr, A.MÜler-Groeling and H.A. Weidenmüller, Phys. Rep. 299 (1998) 189 and A.D.Mirlin, Phys. Rep. 326 (2000) 260.
- [16] G.M.Zaslawski, Phys. Rep. 80 (1981) 151; M.V.Berry, Chaotic Behaviour of Deterministic Systems, Les Houches Summer School 1981, North Holland, Amsterdam (1983); O.Bohigas, M.J.Giannoni and C.Schmit, Phys. Rev. Lett. 52 (1984) 1, cf. also [15].